

This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

+ *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.

+ *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.

+ *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.

+ *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at `http://books.google.com/`



6000043285S

G.96.a.19.



E. BIBL. RADCL.

199611



Ornithological Dictionary.


Cirl Bunting.

Ornithological Dictionary;

OR,

Alphabetical Synopsis

OF

BRITISH BIRDS.

BY

GEORGE MONTAGU, F.L.S.

IN TWO VOLUMES.

VOL. I.

LONDON:

PRINTED FOR J. WHITE, FLEET STREET,

BY T. BENSLEY, BOLT COURT.

1802.

INTRODUCTION.

ERRATA, VOL. I.

Introduction, page x, marginal note, ove-viparous, *read* oviviparous
Avoſet, Scooping Avocet, *read* Avoſet
Bean-crake, vide Gallinula-crake, *read* Gallinule-crake
Bunting-Cirl, Emberona, *read* Emberiza
Bunting-reed, Ortolan de Voſeaux, *read* Roſeaux
Diver-black-throated, Speckled Zoon, *read* Loon
Duck-long-tailed, and elſewhere, Trachia, *read* Trachea
Duck tufted, line 5, black yellow, *read* dark yellow; line 49, great ſpeculum, *read* green
Dun-Diver, line 44, aquatic birds, in particular, *read* aquatic birds in particular,
Eagle-Golden, Falco Cryſactos, *read* Cryſaetos
Eagle-Sea, line 54, Area, *read* Aerie
Flycatcher, Bean Bird, *read* Beam
Finch-bul, Bouvreuil Briſ. *read* Buf.
Gadwall, line 25, ſuppoſed to bred, *read* breed
Gallinule-common, line 1, this birds, *read* bird
Gambet, line 1, the ſpecies, *read* this
Hoopoe, line 33, of bluiſh white, *read* bluiſh white
Jay, line 37, ſometime, *read* ſometimes
Knot, line 57, Lewis's, *read* Lewin's
Lark-tit, line 25, Pipip, *read* Pipit

Of the various writers on Britiſh Ornithology the public are particularly indebted to Mr. Pennant, who has not only been diffuſe on the ſubject, but has given a great variety of excellent figures. The

INTRODUCTION.

THE variety of publications on Ornithology are as extensive as inſtructive; but the difficulty of diſcovering any particular ſpecies in the works of an author, where the bird is called by a different name from which it may be provincially denominated, has, we conceive, been a preventive to a more enlarged inveſtigation of that part of the natural hiſtory of our country. To perſons well acquainted with the generic characters the difficulty may not be ſo great; but the general part of mankind might wade through volumes before they would find the object of their inquiry. For ſuch perſons, therefore, who are deſirous of being better acquainted with the moſt beautiful part of the animal creation, we have publiſhed the following Dictionary of Britiſh Ornithology, in hopes of advancing knowledge on the ſubject.

Of the various writers on Britiſh Ornithology the public are particularly indebted to Mr. Pennant, who has not only been diffuſe on the ſubject, but has given a great variety of excellent figures. The

laſt edition of the *Britiſh Zoology* is replete with information, and far exceeds the bounds of any thing written before on the ſubject of Engliſh birds. Since which the *General Synopſis of Birds*, publiſhed by Dr. Latham, has added much new light to this branch of hiſtory; which, together with his *Index Ornithologicus*, is a maſterpiece of erudition and knowledge. But theſe are too extenſive and too complex for the more confined part of the ſcience or ſtudy of Britiſh birds, and too expenſive for general uſe.

Theſe conſiderations have induced the author to lay before the public this work, which is at once calculated to give every information on the ſubject, to render the ſtudy more eaſy, and to be not only within the reach of moſt perſons pockets who are capable of ſcientific reſearches, but that it will be found an index to all the Engliſh works of credit on the ſubject in general uſe, as well as to many of the beſt authors in other languages. It has been our intention to confine this work within as ſmall a compaſs as poſſible, in order to render it more portable; at the ſame time we have been ſufficiently diffuſe in the deſcription of each ſpecies, as well as in the natural hiſtory belonging thereto. There are, however, ſome general remarks, which could not be ſo well placed in the alphabetical order of this work; and as they are ſuch as ought not to be omitted, we have thought proper to throw them into the form of an introduction.

In

In reſpect to the anatomy of birds we confeſs ourſelves not ſufficiently qualified to enlarge upon the ſubject; we ſhall, however, point out ſome ſingularities in the formation of particular ſpecies, which may add to the illuſtration of Ornithology.

Firſt, then, the *ſtomach* of birds forms them into two natural diſtinct zootomic claſſes: thoſe with *cartilaginous ſtomachs*, covered with very ſtrong muſcles, called a *gizzard*; and thoſe with *membranaceous ſtomachs*, more reſembling that of carnivorous quadrupeds. The former is given to ſuch whoſe principal food is grain and ſeeds of various kinds, or other hard ſubſtances that require much friction to comminute; to aſſiſt which gravel is neceſſary: the latter is given to thoſe who are carnivorous or piſcivorous, and their digeſtion accelerated more by the *gaſtric juice* than by the action of the ſtomach. Thoſe of the firſt claſs digeſt or retain every ſubſtance taken in; and thoſe who eject or diſgorge innutritious matter unavoidably taken in, ſuch as feathers, fur, bones, &c. belong to the ſecond claſs; conſpicuous in the Falcon and Owl genus, and others that feed on fiſh. Granivorous birds ſeem to poſſeſs a power of retaining the ſmall ſtones taken into the *gizzard*, or evacuating them when they become poliſhed and leſs uſeful, but cannot diſgorge them. In a ſtate of nature the quantity of gravel taken in muſt be regulated, no doubt, by the ſenſation of the *ſtomach*; but, extraordinary as it may ſeem, in domeſticated animals

thoſe inſtinctive faculties are deranged. We have known inſtances where the whole cavity of the *gizzard* has been filled with gravel ſtones. We once remember part of a brood of young Ducks half grown taking in ſuch a quantity of gravel as not only filled the *gizzard* but the *craw*, and even the *gullet*, and ſoon after died.

We ſhall now conſider *parturition* and *incubation*. And as the Cuckow appears to be an exception to the general rule, it may not be amiſs to make ſome inquiries into the natural hiſtory of that extraordinary bird. But firſt we ſhall remark, that by experiment it appears that birds do not inſtinctively know the neceſſary time of incubation; for we have repeatedly taken the eggs of a bird unincubated, and placed them under another of the ſame ſpecies who was on the point of hatching, and *vice verſa* thoſe on the point of hatching into the neſt of ſuch who had only began to ſit; and in both caſes the young were brought to maturity. Birds will ſometimes diſcriminate the egg of another ſpecies put into their neſt, and will turn it out; but they will frequently breed up the young of another when exchanged, provided they are of the ſame age, and not very large when the experiment is made.

It has long been a received opinion, that the Cuckow depoſited its egg in the neſt of ſome other bird; that it never ſat on its own egg, but left it to be incubated by the bird in whoſe neſt it was depoſited

depoſited; that it ſeldom laid more than one egg, becauſe that number is moſt commonly found in a neſt; and that as no other young but that of the Cuckow is found in a neſt, it was imagined that the old Cuckow either deſtroyed the eggs or young of the bird whoſe neſt it dropped its own egg into.

In this ſtate had the natural hiſtory of the Cuckow remained till the ingenious Mr. Genner illuſtrated it in a letter to Mr. John Hunter, publiſhed in the *Philoſophical Tranſactions*. To the light this gentleman has thrown on the ſubject every naturaliſt muſt feel himſelf obliged. But, knowledge arriving by ſlow degrees, and the fallibility of human power being certain, will, we hope, plead an excuſe for further attempting to elucidate the hiſtory of that ſingular bird. The opinion of different perſons coinciding cannot fail to ſtrengthen an aſſertion; and we are happy to declare, from perſonal experience, that we agree with that gentleman in reſpect to the incubation and nutrition of the Cuckow, and the phenomenon of the infant bird throwing the eggs or young of the foſter parent out of the neſt; and only differ ſomewhat in reſpect to the parturition of that and other birds. It muſt be underſtood we do not mean to contradict that author abſolutely in any point, but only to offer ſuch ideas to the friends of ſcience as may further ſtimulate to a more minute inveſtigation. Many years ago a Cuckow's egg was brought to

me taken out of a Reed Bunting's neſt; I immediately put it into the neſt of a Swallow, in an out-building, who had juſt begun to ſit. On viſiting it about the time I expected the excluſion of the young, was ſurpriſed to find the young Cuckow ſole poſſeſſor of the neſt. This and ſimilar inſtances convinced me that the eggs of the foſter parent could not be deſtroyed by the old Cuckow. It was difficult to ſuggeſt what became of the bird's eggs in whoſe neſt the Cuckow choſe to depoſit hers. It was unnatural to ſuppoſe the bird would throw out or deſtroy its own in preference to that of the intruder; and the circumſtance of young Cuckows never being found with companions in the neſt, could not admit of a ſuppoſition that they were deſtroyed by accident. In this very unenlightened ſtate, then, was the hiſtory of this bird, till Mr. Genner proved, by a variety of experiments, exhibited under his own inſpection, the extraordinary circumſtance of the young Cuckow's turning out its neſtling mates, or the eggs. But notwithſtanding this undoubted authority, many incredulous perſons, to our knowledge, diſbelieve the poſſibility of the facts related. Too apt are ſome perſons to wonder at and diſbelieve any thing ſeemingly out of the ordinary courſe of nature; whereas it ſhould only excite our admiration. That Being who alone can give animation, can moſt aſſuredly endow it with what powers and properties ſeems beſt to his unerring wiſdom. To diſallow things

things becauſe our contracted comprehenſions and confined ideas are limited within ſo ſmall a compaſs, ſavours of atheiſm. It is more than probable the all-wiſe Creator of all things never intended we ſhould penetrate into the reaſons for all his actions; and ſhall we mortals of fallible nature deny well-atteſted facts becauſe we are not able to ſee into the wiſdom of an infallible Being?

But to return to the Cuckow. Previous to the abovementioned publication I had taken much pains towards inveſtigating the ſeveral phenomena I had noticed in this bird, and was ſo fortunate to have ocular proof of the fact related by Mr. Genner, of a young Cuckow turning out of a Hedge-Sparrow's neſt a young Swallow I had put in for the purpoſe of experiment. It is needleſs to recite all the circumſtances attending this extraordinary bird, as that gentleman has ſo amply explained it; I ſhall therefore only add, that I firſt ſaw it when a few days old in the Hedge-Sparrow's neſt, in a garden cloſe to a cottage, the owner of which aſſured me the Hedge-Sparrow had four eggs when the Cuckow dropped in a fifth; that on the morning the young Cuckow was hatched, two young Hedge-Sparrows were alſo excluded, and that on his return from work in the evening nothing was left in the neſt but the Cuckow. At five or ſix days old I took it to my houſe, where I frequently ſaw it throw out the young Swallow for four or five days after. This ſingular action was performed by inſinuating

itſelf under the Swallow, and with its rump forced it out of the neſt with a ſort of jerk. Sometimes, indeed, it failed after much ſtruggle, by reaſon of the ſtrength of the Swallow, which was nearly full feathered; but after a ſmall reſpite from the ſeeming fatigue it renewed its efforts, and ſeemed continually reſtleſs till it ſucceeded. At the end of the fifth day this diſpoſition ceaſed, and it ſuffered the Swallow to remain in the neſt unmoleſted.

In reſpect to the Cuckow being able to continue laying from the production of the firſt egg till ſhe leaves us, we are doubtful, that is, in regular ſucceſſion; but we ſuſpect, contrary to all other birds, it has the property of retaining its egg, of which we ſhall ſpeak hereafter. It is poſſible ſome may lay two lots of eggs during their ſtay with us; but then we have reaſon to believe there is a conſiderable ſpace of time between the firſt and ſecond lot. In two or three females diſſected at the time they firſt began to lay, we could only diſcover four or five eggs that could poſſibly be laid ſucceſſively; from the ſmalleſt of which to what may be termed the ſecondary eggs, there was a ſudden break off, not a gradual decreaſe in ſize. Now if this bird was able to continue laying for a month or ſix weeks together daily, all the eggs to be protruded in that time would appear in regular gradation. If any bird is killed at the time of laying the firſt egg, the number of eggs deſtined to be excluded ſucceſſively may be pretty eaſily aſcertained; and we

are

are confident no bird we are yet acquainted with, the Cuckow excepted, has the power of keeping back its eggs after it has received the ſtimulus of the male fecundity; and after the firſt egg is laid the others muſt ſucceſſively follow, one after the other, each in twenty-four hours, with a few exceptions in the larger undomeſticated birds; and we know the force of nature is often ſufficiently powerful to ſeparate eggs from the *ovarium* without male contact*. Cuckows, on the contrary, we conceive, are capable of retaining their egg after it is arrived at maturity; and we think there is much reaſon for ſuppoſing this bird is endowed with ſo extraordinary a property, conſidering it makes no neſt, and that it is obliged to ſeek the neſt of another bird to depoſit it in, and that neſt fit to receive it, into which one egg only is dropped †, and other neſts to ſeek for the like purpoſe, it becomes not only a work of labour, but of hazard and uncertainty; therefore, did it not poſſeſs ſuch a qualification, we ſhould conceive the difficulties it had to encounter were too numerous to enſure a continuation of its ſpecies. Suppoſe this bird was obliged, like others, to lay its eggs ſucceſſively day after day, which we believe to be

* Pullets often lay eggs in a coop without impregnation, and ſmall birds often in cages: many inſtances of which we have ſeen. Theſe are of courſe unprolific.

† Where two have been found in one nest they certainly were laid by different birds.

five or ſix, it is hardly probable that five or ſix neſts ſhould be found in a ſtate proper to receive them, much leſs if ſhe laid a greater number, as has been ſuggeſted.

We think it reaſonable to conceive the female Cuckow receives the male's embraces, from a certain period after her arrival, till ſhe is near laying her firſt egg; and that ſhe does not admit the male again till after ſhe has laid her uſual number. If ſhe has been fortunate in diſcovering ſuch neſts as anſwer her purpoſe, and have laid all her firſt forwarded eggs early in the ſeaſon, then, after a ſhort interval, it is poſſible nature may ſtimulate again, and a ſecond lot of eggs be produced; ſo that, in either caſe, Cuckows may lay eggs about the time they depart from us; but we cannot help doubting whether more than one lot of eggs is laid during its continuation in this country; and that alone may be a work of time, for the reaſons before mentioned.

If a Cuckow continued laying each day ſucceſſively from the time of her excluding the firſt egg till near the time of migrating, ſurely a greater number of eggs or young of that bird would be found, eſpecially as they are ſo diſperſed; but, on the contrary, although we have been anxious to procure the eggs or young for ſeveral years together, we have not been able to ſucceed; yet the old birds have been in plenty about us.

Thoſe who ſuppoſe a bird is capable of producing

ing eggs at will, or that any bird is excited to lay more eggs than uſual by daily robbing their neſt, are certainly miſtaken. In a domeſticated fowl it is probable the deſire of incubation may be prolonged by leaving little or nothing in the neſt to ſit on. It will therefore lay the number allotted by nature, which is determined before the firſt egg is produced. If it is prevented from incubation by any means whatever, it may begin again to lay in five or ſix days; but there is always an interval of a few days, and ſometimes as many weeks, which muſt wholly depend on the age and vigour of the bird. When it happens that a freſh lot of eggs is laid with only a few days interval, and that perhaps in the ſame neſt, it is deemed a continuation, for want of nice obſervation; but we are not to look to domeſticated animals for natural cauſes, for thoſe are taken from their ſtate of nature. Let us look to birds in their natural wild ſtate, and ſee if any well-atteſted inſtances are to be found where they have laid more eggs ſucceſſively by taking one from the neſt daily: for inſtance; the number laid by a Hedge-Sparrow is commonly five, ſometimes only four, and rarely ſix; will the taking away the daily-laid egg produce a ſeventh or an eighth? No: we believe there never was an inſtance; at leaſt we have never been fortunate enough to diſcover one in the great variety of experiments we have tried on various birds, amongſt which was the Swallow, which has been declared to lay as

many as nineteen*. A bird will only lay the uſual number peculiar to the ſpecies; and if, at the period of incubation, it perceives the neſt emptied, it is deſerted. The link of nature having been broken, the female ſtimulates to love again, and ſoon brings forward by that *ſtimulus*, aided by the *male fecundity*, a new lot of eggs; never more than the former, and uſually leſs, becauſe this is properly a forced production, at the additional expence of the vigour of the bird, and loſs of animal parts; which is the cauſe of great variation as to the number of eggs laid by domeſtic fowls, depending entirely on the ſtrength of conſtitution, and the nouriſhment of the food. In all animals taken immediately under the care of man, the dictates of nature are partly ſuppreſſed, their food changed, habits and manners altered, and diſeaſe often enſues, which is the origin of the great variety of colours in reclaimed animals.

Nature purſues invariably one courſe; therefore to draw a general rule of her actions we muſt ſtrictly adhere to her in an unmoleſted, uncultivated ſtate; for if we deviate from that we muſt infallibly err. We do not mean to ſay accidental varieties do not take place in animals unreclaimed, but ſuch *luſus* are by no means common; and when we ſee a bird materially deviate in colour from its ſpecies, we may conſider it as a conſtitutional defect; that

* An aſſertion of Dr. Liſter. Vide Br. Zool. p. 337.

the

the natural ſecretions are changed or ſuppreſſed with which the feathers are dyed. To enter minutely into a diſcuſſion on this head would ſwell this Introduction beyond its limits; all we wiſh is, to point out the neceſſity of ſtrictly adhering to nature for obſervations on natural cauſes. A domeſtic fowl who will ſit for ſix weeks upon an empty neſt is not to be produced as a proof of the actions of nature. Will any bird, in its natural wild ſtate, continue to ſit on its neſt after the eggs are taken out? One egg, indeed, is ſometimes ſufficient to produce the act of incubation; but what is it then prevents the ſecondary eggs from coming forward, when it is well known if a bird is prevented from ſitting ſhe ſoon reſumes her deſire of propagation natural to every animated being? Becauſe the very act of incubation is the effective cauſe; the line which nature has drawn, and which the animal by inſtinct feels. We conceive the production of a ſecond lot of eggs to be an extraordinary exertion of nature; a wonderful proof of the affection of the all-wiſe Creator for the preſervation and continuation of his creatures, and the reſources he has furniſhed ſome animals with, in caſe of neceſſity, to prevent the total extinction of the breed. It is but few birds, if any, that would produce a ſecond lot of eggs in the ſame ſeaſon if unmoleſted; but if their neſts are deſtroyed, it is probable three or four ſeparate lots may be protruded.

We have never been able to diſcover with certainty,

tainty, either in the Redbreaſt or Hedge-Sparrow, who are the earlieſt breeders, the production of a ſecond brood after the firſt has been brought to maturity. Their attention to their young continues long after they leave the neſt. The great exertion to collect food for ſo many muſt exhauſt the animal ſpirits, to recruit which is a work of time; ſo that the ſeaſon is too far advanced for a ſecond production. The ſecondary eggs being brought forward is not effected by the will of the bird, but is cauſed by the dictates of nature, the impulſe of love.

We ſhall here take notice of the eggs of a Hedge-Sparrow being found in a neſt of that bird in a prolific ſtate, and a young Cuckow of a fortnight old covering them, as related by Mr. Genner, in order to ſtrengthen a ſuppoſition we ſhall hereafter mention. May not this be owing to a very different cauſe than what has been ſuſpected? If the Cuckow had dropped an egg into the Hedge-Sparrow's neſt before it was finiſhed, it is unlikely it would have been ſuffered to remain; but even ſuppoſe it had, it would not have been ſat on till the Hedge-Sparrow began to incubate her own eggs; conſequently the Cuckow's egg, which is largeſt, could not be hatched before the others in the common courſe of nature. Again: if we ſuppoſe the Hedge-Sparrow had previouſly laid her eggs, which ſhe ſat on together with the Cuckow's, and that the young Cuckow, ſoon after it was hatched, had

turned

turned out of the neſt the eggs or young of the Hedge-Sparrow, which is the natural conſequence, how ſhould theſe eggs come into the neſt containing a living *fœtus*? It is very improbable a bird ſhould lay eggs in a neſt where ſhe had young, for ſuch we may call the young Cuckow, the Hedge-Sparrow not knowing the difference. If birds were capable of this, it would be ſimilar to ſuperfetation in viviparous animals, of which there are but few inſtances. We can therefore only account for this ſingular circumſtance by ſuppoſing the Cuckow is actually endowed with the property of retaining its egg in the *uterus*, after it is matured, till it has diſcovered a neſt in a ſtate fit to depoſit it. The conſequence of this retention would be a dilatation of the *embryo* by the internal heat of the body*, and the *fœtus* advanced towards perfection in proportion to the time the egg remained in that ſtate. Of courſe, after ſuch a previous enlargement of the *fœtus*, was the egg dropped into the neſt of a bird on the point of ſitting, it would moſt certainly be hatched as long before the eggs of the bird whoſe neſt it was depoſited in as it had been forwarded in the *uterus*. It has frequently been obſerved, that where the egg of a Cuckow has been found in the neſt of a bird together with ſome of its own, that the Cuckow's egg is hatched

* The Viper is oviparous, or rather ove-viparous, hatching its young by the internal heat of its body.

firſt.

firſt *. This ſeems difficult to account for, unleſs upon the principle we have ſuggeſted, as the egg of that bird is rather ſuperior in ſize to that of any bird whoſe neſt it makes choice of to depoſit it in, amongſt which the Yellowhammer's is the largeſt, weighing in general from thirty-ſix to forty-ſix grains; whereas that of the Cuckow weighs from forty-four to fifty-four grains. The other birds which the Cuckow more generally chooſes to incubate its egg ſeldom produce eggs above forty grains in weight, and moſtly from thirty to thirty-ſix; if, therefore, the *embryo* of the Cuckow was not ſometimes enlarged before the egg was laid, is it reaſonable to ſuppoſe it would be firſt hatched?

In reſpect to the young Cuckow ſuffering the Hedge-Sparrow's eggs to remain in the neſt, as above mentioned, which is contrary to the general rule, we can only ſuppoſe ſome defect exiſted in the young Cuckow, or that theſe eggs were ſo placed in ſome ſmall cavity in the bottom of the neſt that its utmoſt efforts could not diſengage them. Theſe and other accidental cauſes are much more likely than that the Hedge-Sparrow ſhould lay theſe additional eggs at the time ſhe was performing the act of nutrition to her ſuppoſed offſpring.

From late obſervation we conclude the female

* Page 3, Genner, Nat. Hiſt. of the Cuckow.

Cuckow

Cuckow retires from the male ſo ſoon as ſhe feels an inclination to depoſit an egg, and does not admit him again while ſhe is capable of laying, all the eggs of the firſt ſet being fecundated at the ſame time. This, indeed, is probable with reſpect to all birds, though it is difficult to aſcertain how the prolific quality is regulated ſo as only to affect ſuch a portion of eggs in the *ovarium**; and that every particular ſpecies ſhould almoſt invariably lay the ſame quantity.

The extraordinary growth of a young Cuckow is no more than what happens with others produced from eggs of a diſproportionate ſize; for inſtance, the egg of a Raven is not half the ſize of that of the Kite, and yet the young of the former are not longer arriving at maturity than the latter. Some birds are hatched blind, and entirely naked; others poſſeſs the organs of ſight, and are covered with down. Thoſe birds who feed and breed on the ground, and never perch on trees, are generally provided with warm covering, and have the uſe of their eyes and legs ſoon after they are excluded from the egg †. Theſe are not fed by the parent bird, being more perfectly formed when hatched, but ſoon diſcover by inſtinct their proper food; on the

* Domeſtic fowls, ſeparated from the male after the firſt impregnation, are known to lay many prolific eggs. *Young on Turkeys, in Nat. Hiſt. of Norfolk.*

† With a few exceptions. Some that occaſionally perch have theſe properties.

contrary, predacious birds are totally helpleſs at firſt, though warmly clothed with down.

The extraordinary and rapid growth of the egg is alſo worthy notice. We before obſerved there is a line drawn by nature to prevent more than a certain quantity of eggs, peculiar to each ſpecies, being fecundated at once; otherwiſe a bird in one ſeaſon might produce all the eggs ſhe poſſeſſed, and afterwards become uſeleſs in propagation.

But notwithſtanding only a portion of the *ovaries* are impregnated at the ſame time, yet the ſtimulus to love conſiderably increaſes the ſize of all the eggs in the *ovarium*. When this ſtimulus ceaſes, be it from what cauſe it may, the organic particles ceaſe to flow to that part; therefore, as a redundance of organic particles excite love by diſtending thoſe parts neceſſary for production; ſo, on the contrary, the want of a ſufficient quantity reduces them to their primitive ſtate. It ſeems abſolutely neceſſary that a dilatation of the *ovaries* ſhould take place before they can be fecundated; and that by the law of nature only a certain number ſhould be ſufficiently large to be impregnated at the ſame time; and that no others can poſſibly receive the *male ſtimulus* till after the firſt ſet are produced. We have before remarked, a ſuperabundance of organic particles is cauſe ſufficient to ſeparate an egg from the *ovarium* without *male contact* *.

The-

* It is poſſible a bird, in its natural ſtate, may lay an egg unimpregnated, which may account for addled eggs being found

The growth of an egg after impregnation is exceedingly rapid: the yolk only is formed in the *ovarium*, where it remains till within twenty-four hours of its being produced; when that part is fully matured, it ſeparates, and falls down the *oviduct* into the *uterus*, where the egg is perfectly formed; firſt the *vitellus*, or yolk, is ſurrounded by the *albumen*, or white; and laſtly is covered with a calcareous ſhell *. The very expeditious growth of theſe laſt appear to be an extraordinary exertion of nature. The calcareous covering of an egg is concreted and formed in a moſt expeditious manner; a few hours only ſeems neceſſary for this work. Only one *vitellus* ſeparates from the *ovarium* at a time (except as we ſhall hereafter mention), till the excluſion of which no other ſucceeds. But as this is a daily production, with few exceptions, there is no more time allowed for perfecting the *albumen* and ſhell than twenty-four hours.

As the courſe of nature is ſometimes obſtructed by remote cauſes, there are few general rules without ſome exceptions; but an individual deviation by no means perverts the law of nature, but is ſimply an individual defect. We ſhall here inſtance

found in neſts. We once found a Whitethroat ſitting on four eggs perfectly dried up. This alſo ſeems to prove birds do not know the neceſſary time of incubation.

* We do not mean to enter minutely into the phyſiological deſcription of an egg, which may be found in various publications.

imperfect eggs ſometimes produced, ſuch as want the *vitellus* *, and others containing two yolks; and although there are probably very few inſtances of ſuch productions from birds in their wild ſtate †, yet it is no uncommon thing in domeſtic fowls. We are alſo told of eggs with double ſhells; and we have frequently ſeen eggs without any calcareous covering, but wrapped in a ſoft pliable ſkin like vellum.

To account for theſe extraordinary productions we muſt conceive a defect exiſts at the time in ſome part of the animal body; and as the *vitellus* and *albumen* derive their origin from different parts, it is natural to conclude, in thoſe preternatural eggs deſtitute of yolk, the cauſe proceeds from ſome defect in the *ovarium*. May it not be occaſioned by an unequal ſtimulus in the parts neceſſary to perfect the egg, and that the growth of the *vitellus* is not in proportion to the *albumen?* Thus while the veſicles appointed for collecting, preparing, and uniting, the organic particles of the *albumen* in the *uterus* have all their effective powers, thoſe of the *ovarium* are weak; of courſe the one goes on with the operations appointed by nature, while

* The *Centininum Ovum* of naturaliſts, vulgarly called a Cock's egg. This name has been given it from a ſuppoſition it was the hundredth egg, or laſt the bird could lay.

† The egg of an Hedge-Sparrow and of the Common Tern are the only inſtances, in a ſtate of nature, we ever remember finding yolkleſs, and were of courſe not half the uſual ſize.

the

the other is ſtopped in its progreſs for want of a ſufficient quantity of organic matter to bring it to perfection at the ſame time. Hence the reaſon of the common fowl ſometimes producing three or four imperfect eggs following, by which time the *vitellus* becomes ſufficiently large, and a regular ſucceſſion of perfect eggs are produced. On the contrary, when, from the ſame cauſes, the growth of the *vitellus* in the *ovarium* is too luxuriant, two yolks paſs the *oviduct* together, which being ſurrounded with the uſual quantity of the *albumen*, is brought forth in the form of a ſingle egg, of an extraordinary ſize. From theſe it is poſſible twins are produced, but more frequently deformities, which are ſeldom hatched.

We ſhall now proceed to the ſubject of incubation. It is probable birds are endowed with an inſtinctive power of regulating the neceſſary heat for this purpoſe; of courſe, ſhould the heat of the air, together with the natural warmth of the body on the cloſe contact of the bird to the eggs, be too great, her feelings would dictate the neceſſity of leaving them for a time to cool. At the early part of incubation birds quit their eggs more frequently than at the time the *fœtus* is more perfect. Yet, in the advanced ſtate, the *embryo* young is not in more danger of being deſtroyed, if ſo much; for we have frequently found a living *fœtus* in an egg that has been taken from the neſt two days. If, however, the young is within a few hours of being

excluded, and the egg is ſuffered to be ſome time cold, it either dies, or becomes ſo weak as not to be able to extricate itſelf from the ſhell. Various degrees of heat will enlarge the *embryo* young, but regular heat ſeems neceſſary to its production; and yet artificial heat, regulated by the heat of a brooding bird, will not produce young with ſuch certainty *. Birds frequently turn and change the ſituation of their eggs in the neſt; beſides this, it is poſſible the moiſture of the bird's body may aſſiſt the natural growth of the *fætus* and production of the young.

The male birds of ſome ſpecies ſupply the place of the female on the neſt; but then it is of ſhort duration, and rarely, if ever, when the eggs are near hatching: at that time the female is frequently fed by the male †. Many ſpecies of birds poſſeſs a *reſervoir* for food called a *craw*, or *crop*; this ſeems to anſwer the ſame purpoſe as the firſt *ſtomach* in ruminating animals. Here it is the food is ſoftened and prepared for the ſtomach; from this *reſervoir* it is by ſome ejected for the purpoſe of feeding their young; conſpicuous in the Pigeon.

* In Egypt a vaſt quantity of eggs are hatched by artificial heat in ſtoves. It is probable, however, one third or one fourth miſcarry. The neceſſary heat for this purpoſe is about 96 degrees of Fahrenheit's thermometer, or 32 of Reaumur's ſcale.

† This is not common to all ſpecies, but very conſpicuous in the Rook, the Pigeon, and many others.

The

The Rook has a ſmall pouch under the tongue, in which it carries food to its young. It is probable the uſe of the *craw* may be extended further than is generally imagined; for, beſides the common preparation of the food to aſſiſt its digeſtion in the *ſtomach*, there are ſome ſpecies who actually ſecrete a *lacteal* ſubſtance in the breeding ſeaſon, which, mixing with the half-digeſted food, is ejected to feed and nouriſh the young. The *mammæ*, from which this milky liquor is produced, is ſituated on each ſide the upper part of the breaſt, immediately under the *craw*. In the female Turtle Dove we have met with theſe glands tumid with milky ſecretion; and we believe it common to both ſexes of the Dove genus. The Corvorant or Pelican genus poſſeſs no *craw*; but, to ſupply its place, they have a looſe ſkin at the baſe of the under mandible capable of great diſtention, in which they carry fiſh to their young. The Buſtard is ſaid to poſſeſs a bag of an aſtoniſhing ſize, for the purpoſe of retaining water. But the moſt unaccountable and extraordinary formation in the *trachea* of many of the males of the Duck genus, called a *labyrinth*, is beyond our reach to diſcover the uſe of; as well as the ſingular flexure in the *windpipe* of the Hooping Swan and Crane *.

We

* It is much to be wiſhed naturaliſts would pay more attention to theſe ſingular internal formations of birds; it would no doubt throw light on many obſcurities in the diviſion of

We ſhall here mention the difficulty naturaliſts labour under in the deſcription of ſome birds, their eggs, and neſt. From a variation in plumage in ſome birds we doubt not naturaliſts have enumerated a greater variety in ſome genera than there really are; in others a ſimilitude of colour in diſtinct ſpecies have occaſioned their being confounded and blended together for one and the ſame. We do not ſo much wonder at it in exotics, who come to us in an imperfect ſtate. But this has frequently happened to thoſe of our own country.

If the ornithologiſt attended more to the habits and manners of birds he would not be ſo liable to be led into theſe errors, being invariably diſtinct in ſome particulars, either in their notes, their neſt, eggs, place of reſort, and various other circumſtances neceſſary to be conſidered. It is true, the eggs, and ſometimes the neſt, of diſtinct ſpecies are ſo nearly allied *, that it would puzzle the moſt ſcrutinizing eye to determine; but other concomitant circumſtances in the habits and manners of birds, ſuch as the place of *nidification*, &c. &c. would determine. The length of time ſome birds are arriving at maturity in their plumage is cauſe

ſpecies. The number of tail-feathers ſhould alſo be well aſcertained, for this alone is a mark of diſtinction, as in the Corvorant and Shag.

* That of the Marſh and Cole Titmouſe, which are ſtill ſuppoſed to be only a variety of the ſame ſpecies by ſome perſons who have never minutely attended them in their natural haunts.

of

of very great difficulty in determining the ſpecies. The Gulls are not maturely feathered till the return of the breeding ſeaſon, and many not till the ſecond, or perhaps the third year *. It has been obſerved, by an author of great ability, and for whom we have the higheſt reſpect, that the quill and tail feathers are permanent; but in this we cannot agree, as we could produce innumerable inſtances to the contrary †. Beſides the markings of the feathers, there are other markings which change by age and ſeaſon, ſuch as the colour of the bill, legs, irides, and ſkin, particularly the orbits of the eye. Some ſpecies of birds ſeem to change their winter and ſummer feathers, or at leaſt in part: in ſome this is performed by moulting twice a year, as in the Ptarmigan; in others only additional feathers are thrown out. But we have no conception of the feathers themſelves changing colour, although we have been informed of ſuch happening in the courſe of one night. That confined birds do ſometimes become wholly

* There is very little doubt the Winter Mew is no other than the young of the Common Gull; as no doubt the Wagel of authors is the young of the Great Black-back Gull of this work.

† For inſtance: the male Keſtrel, which, from being barred on the tail feathers, becomes wholly aſh-coloured, except at the end; alſo the Black-headed Gull, whoſe tail is at firſt tipped black, and the quills without white at their ends, change in the firſt moulting. The Herring Gull, and others whoſe tail is white when matured, are for the firſt two years mottled.

black,

black, is unqueſtionable; but this is occaſioned by a ſucceſſion of feathers; and we have more than once ſeen the Bulfinch in the ſtate of changing, in which the black on the breaſt firſt appeared. The colouring of feathers is perhaps one of the moſt wonderful productions of nature. How the colouring ſecretions are diſpoſed in ſuch regular order, far ſurpaſſes human abilities to determine. It is probable, however, this is performed in an *embryo* ſtate. The circulation in a feather ſeems to ſtop as faſt as the parts are matured, and when it is fully grown, there certainly is no circulatory fluid whatever; of courſe the colours cannot change except in brilliancy. It might naturally be ſuppoſed for want of ſome ſecretory matter the feathers might become more brittle, or be injured, by wet or otherwiſe; but Nature, ever provident in all her ways, has taken care to ſupply every bird, more or leſs, with an external ſecretion of an unctuous nature, ſituated in a glandular bag upon the rump, which they by inſtinct make uſe of for oiling and dreſſing their feathers, as occaſion requires. In water-fowl this bag is moſt conſpicuous; and it is remarkable that birds moſt frequently uſe it after waſhing, previous to their feathers becoming perfectly dry.

Migration of birds is a curious fact that no one denies, though we have not wanted for incredulous perſons who believed the Nightingale was to be found in every hedge during winter. That an accidental

cidental ſummer bird of paſſage may be by diſeaſe prevented from returning to its natural winter quarters, we can admit; becauſe there are variety of inſtances of the Swallow and Martin having been ſeen flying in the months of November and December; rouſed probably from a ſtate of torpidity by an unuſual warmth of the air. So alſo there are inſtances of ſome of our winter migrants remaining with us the whole ſummer. The Woodcock's eggs and young have many times been taken in our woods; but theſe are individual occurrences only, occaſioned by accident. If all the migrative ſpecies did actually reſide with us the whole year, whether in a torpid ſtate or not, we ſhould have daily productions of the fact; and yet, in the various hiſtoric pages of this country, very few inſtances have been related, that had the appearance of authenticity, of the torpidity of any of our ſummer migrants; and ſuch appear to be only that of the Swallow tribe *. That our ſummer migrants come from the ſouth, or warmer climate, and our winter migrants from the north, or colder climate, there is no doubt; but by what rule this is performed is difficult to determine, eſpecially as

* Torpidity is probably the ſtate of thoſe ſummer birds of paſſage which accident may have detained with us during winter; ſimilar to the Hedgehog, the Dormouſe, and Bat. Theſe become inanimate when the thermometer ſinks within ten degrees of the freezing point, as their animal heat keeps pace with the temperature of the air at that time.

we

we have reaſon to believe the ſame bird will return for many years together to the ſame ſpot; and yet, why is this more to be wondered at than the performance of a carrier Pigeon, or the labouring Bee; who returns with certainty to its well-known hive? But what is the more extraordinary part in the hiſtory of our ſummer migrants is, that the males always precede the other ſex in their vernal flight; from what cauſe this is produced we cannot determine, though it is certain the male is no ſooner arrived than he feels the impulſe of love, for, if the weather is warm, he is inceſſant in his call, or ſong.

But as the ſong of birds is not allowed to be the effect of love by an honourable author on the ſubject of ſinging birds, we ſhall endeavour to elucidate this matter from experiments on birds in their natural wild ſtate; and alſo endeavour to prove that their notes are innate, contrary to the ſame author's opinion.

That confined birds will learn the ſong of others they are conſtantly kept with, there is no doubt; but then it is generally blended with that peculiar to the ſpecies. In the ſpring the very great exertion of the male birds in their vociferous notes are certainly the calls to love; and the peculiar notes of each is an unerring mark for each to diſcover its own ſpecies. If a confined bird had learned the ſong of another, without retaining any part of its natural notes, and was ſet at liberty, it is probable

it

it would never find a mate of its own ſpecies; and even ſuppoſe it did, there is no reaſon to believe the young of that bird would be deſtitute of its native notes; for if neſtling birds have no innate notes peculiar to the ſpecies, and that their ſong is only learned from the parent bird, how are we to account for the invariable note each ſpecies poſſeſs, when it happens two different ſpecies are bred up in the ſame buſh or very contiguous, or when hatched and foſtered by a different ſpecies*. There is every reaſon to believe it is neceſſary there ſhould be native notes peculiar to each ſpecies, or the ſexes might have ſome difficulty in diſcovering each other, the ſpecies be intermixed, and a variety of *mules* produced†; for we cannot ſuppoſe birds diſcriminate colours by which they know their ſpecies, becauſe ſome diſtinct ſpecies are ſo exactly alike that a mixture might take place‡. The males of ſong birds, and many others, do not in general ſearch for the female, but, on the contrary, their buſineſs in the ſpring is to perch on ſome conſpicuous ſpot, breathing out their full and amorous notes, which by inſtinct the female knows, and repairs to the ſpot to chooſe her mate. This is particularly verified with reſpect to the ſummer birds of paſſage. The Nightingale, and

* A Goldfinch hatched and foſtered by a Chaffinch retained its native notes.

† This we believe never happens in a ſtate of nature.

‡ The Rook and Crow.

moſt of its genus, although timid and ſhy to a great degree, mount aloft to pour forth their amorous ſtrains inceſſant, each ſeemingly vieing in their love-laboured ſong before the females arrive *. No ſooner do they make their appearance than dreadful battles enſue, and their notes are conſiderably changed; ſometimes their ſong is hurried through without the uſual grace and elegance; at other times modulated into a ſoothing melody. The firſt we conceive to be a provocation to battle on the ſight of another male; the laſt an amorous cadence, a courting addreſs. This variety of ſong laſts no longer than till the female is fixed in her choice, which is in general in a few days after her arrival; and, if the ſeaſon is favourable, ſhe ſoon begins the taſk allotted to her ſex †.

The male now no more expoſes himſelf to ſing as before, nor are his ſongs heard ſo frequent, or ſo loud; but while ſhe is ſearching for a ſecure place to nidificate in, he is no leſs aſſiduous in attending her with ridiculous geſtures, accompanied with notes peculiarly ſoft ‡. When the female has choſen a ſpot for nidification the male conſtantly

* The females of the migrative part of this genus come to us later than the males; ſome indeed not till three weeks after.

† The females make their neſt without much aſſiſtance from the males, with few exceptions.

‡ When we have diſturbed their courting, and ſeparated the ſexes from the ſight of each other, the male aſſumes his uſual vociferous notes.

attends

attends her flight to and from the place, and ſits upon ſome branch near, while his inſtinctive mate places the ſmall portion of material ſhe each time brings to rear a commodious fabric for her intended brood. When the building is complete, and ſhe has laid her portion of eggs, incubation immediately takes place. The male is now heard loud again, but not near ſo frequent as at firſt; he never rambles from her hearing, and ſeldom from her ſight; if ſhe leaves her neſt he ſoon perceives it, and purſues her, ſometimes accompanied with ſoft notes of love. When the callow brood appears he is inſtantly appriſed of it, either by inſtinct, or by the female carrying away the fragment ſhells to ſome diſtant place. The male is now no more heard in tuneful glee (unleſs a ſecond brood ſhould force the amorous ſong again); his whole care and attention is now taken up in ſatisfying the nutrimental calls of his tender infant race, which he does with no leſs aſſiduity than his mate, carrying them food, and returning frequently with the *muting* of the young in his beak, which is dropped at a diſtance from the neſt*. Here we muſt beg leave to digreſs for a moment to remark, that with the utmoſt attention we have never been able to diſ-

* The ſagacity of this, as alſo the diſpoſal of the egg ſhells, is a providential inſtinctive power implanted in theſe little creatures for the ſecurity of their young; to aſſiſt which Nature has given a ſkin, or covering, in which the *fæces* is enveloped.

cover

cover the parent birds giving their young a muſical leſſon; and much queſtion if the late brood of many ſpecies ever hear the ſong of their parents till they join chorus the enſuing ſpring, when they alſo feel the impulſe of love, the great dictate of nature.

The continuation of ſong in caged birds by no means prove it is not occaſioned by a ſtimulus to love; indeed it is likely the redundancy of animal matter from plenty of food, and artificial heat, may produce it; and this is ſufficient reaſon for continuing their ſong longer than birds in their natural wild ſtate, becauſe they have a conſtant ſtimulus; whereas wild birds have it abated by a commerce with the other ſex, by which, and other cauſes, it is prevented. It is true wild birds are heard to ſing ſometimes in the middle of winter when the air is mild, animated by the genial warmth of the ſun, which acts as a ſtimulus. Birds in ſong are generally found by diſſection to have the *teſtes* ſomewhat dilated *. But we ſhall now proceed to ſhew, by experiments, that birds in their natural ſtate may be forced to continue their ſong much longer than uſual. A male Redſtart made its appearance near my houſe early in the ſpring, and ſoon commenced his love-tuned ſong. In two days after a female arrived, which for ſeveral days the male was continually chacing, emitting ſoft inter-

* We believe a Capon never crows.

rupted

rupted notes, accompanied by a chattering noiſe. This ſort of courting laſted for ſeveral days, ſoon after which the female took poſſeſſion of a hole in a wall cloſe to my houſe, where it prepared a neſt, and depoſited ſix eggs. The male kept at a diſtance from the neſt, and ſometimes ſung, but not ſo loud or ſo frequent as at firſt, and never when he approached nearer his mate. When the eggs had been ſat on a few days I endeavoured to catch the female on the neſt, but ſhe eſcaped through my hand. However, ſhe ſoon returned, and I caught her. The male did not immediately miſs his mate; but on the next day he renewed his vociferous calls, and his ſong became inceſſant for a week, when I diſcovered a ſecond female; his note immediately changed, and all his actions as before returned. This experiment has been repeated with the Nightingale with the ſame ſucceſs; and a Golden-creſted Wren, who never found another mate, continued his ſong from the month of May till the latter end of Auguſt. On the contrary, another of the ſame ſpecies, who took poſſeſſion of a fir-tree in my garden, ceaſed its notes ſo ſoon as the young were hatched, and as this beautiful little family cauſed me much delight and amuſement, ſome obſervations thereon may not be unacceptable to the curious reader. When firſt I diſcovered the neſt I thought it a favourable opportunity to become acquainted with ſome of the manners of this minute ſpecies, and to endeavour to diſcover whe-

ther the male ever ſung by way of inſtructing the young ones. Accordingly I took the neſt when the young were about ſix days old, placed it in a ſmall baſket, and by degrees enticed the old ones to my ſtudy window; and after they became familiar with that ſituation the baſket was placed within the window; then at the oppoſite ſide of the room. It is remarkable, that although the female ſeemed regardleſs of danger from her affection to her young, yet the male never once ventured within the room; and yet would conſtantly feed them while they remained at the outſide of the window; on the contrary, the female would feed them at the table at which I ſat, and even when I held the neſt in my hand *, provided I remained motionleſs. But on moving my head one day, while ſhe was on the edge of the neſt, which I held in my hand, ſhe made a precipitate retreat, miſtook the open part of the window, knocked herſelf againſt the glaſs, and laid breathleſs on the floor for ſome time. However, recovering a little, ſhe made her eſcape, and in about an hour after I was agreeably ſurpriſed by her return, and would afterwards frequently feed the young while I held the neſt in my hand. The male bird conſtantly attended the female in her flight to and fro, but never ventured beyond the

* It is probable the focal diſtance of ſuch minute animals eyes is very near, and that large objects are not repreſented perfect on the *retina*; that they do not ſeem to ſee ſuch diſtinctly is certain, unleſs in motion.

window-

window-frame; nor did he latterly ever appear with food in his bill. He never uttered any note but when the female was out of ſight, and then only a ſmall chirp. At firſt there were ten young in the neſt, but probably for want of the male's aſſiſtance in providing food two died. The viſits of the female were generally repeated in the ſpace of a minute and an half or two minutes; or, upon an average, thirty-ſix times in an hour; and this continued full ſixteen hours in a day, which, if equally divided between the eight young ones, each would receive ſeventy-two feeds in the day; the whole amounting to five hundred and ſeventy-ſix. From examination of the food, which by accident now and then dropped into the neſt, I judged from thoſe weighed that each feed was a quarter of a grain upon a medium; ſo that each young one was ſupplied with eighteen grains weight in a day; and as the young birds weighed about ſeventy-ſeven grains at the time they began to perch, they conſumed nearly their weight of food in four days at that time *. I could always perceive by the animation of the young brood when the old one was coming; probably ſome low note indicated her near approach, and in an inſtant every mouth was open to receive the inſect morſel. But there appeared no regularity in the ſupply given by the

* This extraordinary conſumption ſeems abſolutely requiſite in animals of ſuch rapid growth. The old birds of this ſpecies weigh from eighty to ninety grains.

parent bird; ſometimes the ſame was fed two or three times ſucceſſively; and I generally obſerved that the ſtrongeſt got moſt, being able to reach fartheſt, the old one delivering it to the mouth neareſt to her, and after each feed ſhe waited a while to ſee if any *muted* *.

We ſhall now take notice of the external formation of ſome birds, ſo admirably adapted to their mode of life, and other ſingularities not ſo eaſily accounted for. It is well known carnivorous quadrupeds cannot exiſt without drink, and that they take liquid by means of their tongue; rapacious birds ſeldom or ever drink; Eagles, Hawks, and Owls, we have kept for years without ever taſting water. The large crooked talons of theſe birds, and their hooked beaks, are well formed for ſecuring and tearing their prey.

The formation of the feet of the Woodpeckers, the toes being placed two forward and two backward, is well calculated for climbing; and the cuneiform ſhape of the extremity of their bills are ſuited to the purpoſe of cutting holes in decayed trees; their tongue is alſo wonderfully contrived to ſearch out inſects beyond the reach of their bill by its great length, which is double that of the bill at leaſt. The Wryneck has this ſingularity, as well

* The leſſer ſpecies of birds, who are ſo frequently fed, ſeldom or ever mute but immediately after they are fed, by which means the *fæces* are never left on the neſt, but are inſtantly carried away by the parent bird.

as the like formation of the toes; which laſt is alſo found in the Cuckow. But neither of theſe birds is ſeen to climb up the bodies of trees; nor have either the ſtiff ſharp-pointed feathers in the tail, as in the Woodpecker, ſo admirably adapted to ſupport them when in the act of hewing. The call of moſt birds is effected by the lungs and *larynx*; but ſome ſpecies of Woodpeckers, in the breeding ſeaſon, have a very extraordinary and loud call to each other, by ſtrong reiterated ſtrokes of their bill againſt a dead ſonorous branch of a tree, peculiar to this genus. The flight of birds differs exceedingly: ſome fly by jerks, cloſing their wings every three or four ſtrokes, which gives them an undulated motion, very conſpicuous in the Woodpeckers and Wagtails, and in moſt ſmall birds; others fly ſmooth and even; and ſome appear to buoy themſelves in the air without perceptible motion of the wings, ſuch as the Kite and Keſtril Hawk. Moſt birds fly with their legs contracted and neck extended; but there are ſome whoſe length and weight of neck makes it neceſſary to contract it in flight, in order to bring the centre of gravity on the wings; to aid which the legs are alſo extended behind, as in the Heron and Bittern; others fly with extended neck, but are obliged to throw out their legs behind, as in the Duck, Gooſe, and other water-fowl. All aquatic birds and waders walk or run in the ordinary manner, placing one leg before the other alternate; but the greater part of the ſmaller

wings, which are ſometimes uſed as fins to accelerate the motion, but is occaſioned by the preſſure of the water above. In ſwimming on the ſurface a bird has two motions; one upwards, the other forward, at every ſtroke of the feet; ſo that when covered with water that force which was loſt by the upward motion is all directed to the progreſſive, by which it is enabled to purſue its prey or to eſcape an enemy with incredible ſpeed*.

Having now gone through the greater part of the ſingularities which are peculiar to each particular ſpecies or genus, we ſhall cloſe this introduction with ſome remarks on conſummation. The *teſtes* of the male lie within the body cloſe to the ſpine, and require minute inſpection to diſcover them in the winter, even in large birds; at the approach of ſpring theſe gradually increaſe till they become vaſtly larger than their former ſize. In land birds the male has no viſible external parts of generation; the impregnation therefore of the female can only be accompliſhed by coheſion and abſorption; but it is otherwiſe in aquatic birds, who moſt frequently copulate in the water; their genital parts are very conſpicuous at the time of *coition*; by ſuch formation the prolific matter is ſecured. Had this claſs of birds been formed like thoſe of the land their prolificacy would have been uncertain by reaſon of the interpoſition of the water.

* The Otter and Water Rat ſwim much faſter under water than upon the ſurface.

The

The following ſheets have been entirely drawn from our own obſervations, and compiled from the notes of twenty years ſearch and attention to the habits of this beautiful part of the creation in moſt parts of this kingdom. The wood, the mountain, and the barren waſte, the craggy rock, the river, and the lake, are never ſearched in vain; each have their peculiar inhabitants, that enliven the ſcene and pleaſe the philoſophic eye.

Ten thouſand warblers cheer the day, and one
The live-long night; nor theſe alone, whoſe notes
Nice-finger'd art muſt emulate in vain;
But cawing Rooks, and Kites that ſwim ſublime
In ſtill repeated circles, ſcreaming loud;
The Jay, the Pie, and e'en the boding Owl,
That hails the riſing moon, have charms for me.

COWPER.

To the numerous friends who have aſſiſted to enrich my muſeum, and enabled me to deſcribe many rare birds, I beg leave to acknowledge my public obligations; and in particular to my much-eſteemed friends Dr. Latham and Mr. Boys. To the publications of the former on Ornithology this work is greatly indebted for the deſcription of, and remarks on, ſome few of the ſcarcer birds, and whoſe excellent arrangement of birds into genera we have followed.

REMARKS.

THE characteriſtic diſtinctions of each genus will be found annexed to the firſt of the claſs; and if the name of a bird ſought for terminates with the name of a genus, look for that; thus, *Finch-Gold*, inſtead of *Goldfinch*.

If a bird conſiſts of two names, neither of which are the names of a genus, ſuch as *Wood-pigeon*, it will be found under the heads of *Wood* or *Pigeon*, which will refer to *Dove-ring* for *Ring-dove*.

If the generic characters of a bird, ſuch as the *Stint*, is wanted, it will refer to *Purre* as the proper name, where it will be found to be of the *Sandpiper* genus; then look for *Sandpiper* for the generic characters.

Ornithological Dictionary.

A.

ABERDEVINE. Vide *Siſkin.*

ALP. Vide *Finch-bul.*

AMZEL. Vide *Ouzel-ring*, and *Black-bird.*

ARCTIC-BIRD. Vide *Gull-arctic.*

ARSFOOT. Vide *Grebe-tippet*, and *little.*

AVOSET. A genus of birds, the characters of which are,

Bill long, ſlender, very thin, depreſſed, bending upwards.

Noſtrils narrow, pervious.

Tongue ſhort.

Legs very long.

Feet palmated.

Back toe very ſmall.

AVOSET-SCOOPING.

Recurveroſtra Avocetta. *Linn. Syſt.* 156. 1. *Gmel. Syſt.* ii. p. 693. *Briſ.* vi. p 538. t. 47. —*Ib.* 8vo. ii. p. 504. *Raii Syn.* p. 117. A. 1. *Will.* p. 240. t. 60. *Will. Angl.* p. 321.

L'Avocette. *Buf.* viii. p. 466. t. 38.

Scooping Avocet. *Br. Zool.* ii. N° 228. t. 80. *Lath. Syn.* v. p. 293. 1.—*Ib. Sup.* p. 263.

Avocetta. *Ind. Orn.* ii. p. 786. 1. *Don. Br. Birds*, t. 66. *Lewin, Br. Birds*, vi. t. 202. *Walc. Syn.* ii. t. 165. *Pult. Cat. Dorſet.* p. 16.

PROVINCIAL.

Butter-flip. Scooper. Yelper. Picarini. Crooked-bill. Cobler's-awl.

The length of this ſpecies to the end of the tail is eighteen inches; to the end of the toes twenty-two: weight thirteen ounces. Bill black, recurved at the point, flexible like whalebone; irides duſky. The upper part of the head, and half the hind part of the neck, black; the cheeks and whole under parts of the bird are pure white; outer ſcapulars, middle coverts of the wings, and greater quill-feathers, black; the ridge of the wings, greater coverts, back, and tail, white; legs bluiſh grey; toes webbed about half their length.

The Scooping Avoſet is the only ſpecies found in England. It breeds in the fens of Lincolnſhire, and on Romney Marſh in Kent. In winter they aſſemble in ſmall flocks of ſix or ſeven, and frequent our ſhores, particularly the mouths of large rivers, in ſearch of worms and marine inſects, which they ſcoop out of the mud or ſand. It lays two eggs about the ſize of thoſe of a pigeon, white, tinged

tinged with green, and marked with large black ſpots; is ſaid to be very tenacious of its young; when diſturbed at this ſeaſon will fly round in repeated circles, uttering a note that reſembles the word *twit* twice repeated.

The feet of this bird ſeem calculated for ſwimming, but it has never been obſerved to take the water for that purpoſe. We remember one of this ſpecies being wounded in the wing, and floating with the tide for near a mile, when it was taken up alive without ever attempting to ſwim; ſo that the palmated feet ſeem only intended to ſupport it on the mud.

AUK. A genus of birds, the characters of which are,

Bill ſtrong, thick, compreſſed.

Noſtrils linear, placed near the edge of the mandible.

Tongue almoſt as long as the bill.

Toes, three forward webbed; none behind.

AUK-BLACK-BILLED.

Alca Pica. *Lin. Syſt.* i. p. 210. 2. *Gmel. Syſt.* ii. p. 551.

Mergus Bellonii. *Raii Syn.* p. 119. 2.

Utamania. *Will.* p. 243. t. 64.—*Ib. Angl.* p. 324.

Le Petit Pengoin. *Buf.* ix. p. 396.

Black-billed Auk. *Br. Zool.* ii. N° 231.—*Ib.*

fol. 137. *Arct. Zool.* ii. N° 426. *Lath. Syn.* v. p. 320. 6.

Alca Torda. β. *Ind. Orn.* ii. p. 793. 5.

Alca minor. *Brif.* vi. p. 923. t. 8. f. 2.—*Ib.* 8vo. ii. p. 383. *Walc. Syn.* i. t. 85.

The weight of this bird is about eighteen ounces; length fifteen inches.

The bill is black, of the ſhape and ſize of that of the Razor-bill; ſmooth, or void of furrows, in ſome; others poſſeſs three diſtinct furrows, as in a ſpecimen now before us; that neareſt the baſe white; irides hazel. The top of the head, taking in the eyes, part of the neck, back, wings, and tail, are black; the leſſer quill-feathers tipped with white; the ſide of the head, fore part of the neck, and whole under ſide of the body, white; legs brown black.

Dr. Latham, in his Synopſis, mentions his ſuſpicion, that this is no other than the Razor-bill immatured; and, in his Index Ornithologicus, gives it as the young of that bird. We cannot, however, but differ in opinion from that excellent ornithologiſt, by late obſervations on the young of the Razor-bill, before and after they could fly, which differ only from the parent bird in having no furrows in the bill; being deſtitute of the white line from the bill to the eye, and no white on the ſecondary quill-feathers; but the whole head and upper part of the neck is black; which is the eſſential

tial difference between that bird and the Black-billed Auk. This is only found on our coaſts in winter. The Razor-bill breeds with us, and retires in the autumn, at which time none are to be found with the white cheeks and throat amongſt the myriads that haunt our cliffs. It is hardly poſſible, then, to conceive that the young, who are in their firſt feathers ſo like the old ones, ſhould become more unlike them in winter, which is contrary to nature; for obſervation has taught us, that all birds become more like their parents at every moulting; ſo that to make theſe birds one and the ſame ſpecies, we muſt conclude both old and young change their plumage in winter: but this we have no reaſon at preſent to believe. And indeed it is probable, that whatever change may take place, that ſingular line in the matured Razor-bill, running from the bill to the eye, never varies; and which conſtitutes one of the greateſt characteriſtic diſtinctions. The black part of the plumage of this bird is invariably deeper-coloured than in the Razor-bill, particularly about the head. From theſe obſervations we are inclined to give each a ſeparate place in this work, as diſtinct ſpecies.

For further remarks, ſee Razor-bill.

AUK-COMMON. Vide *Razor-bill.*

AUK-GREAT.

Alca impennis. *Lin. Syſt.* p. 210. 3. *Gmel. Syſt.* p. 550.

Penguin. *Raii Syn.* p. 118. *Will.* p. 242. t. 65. —*Ib. Angl.* p. 322. t. 65. *Edw.* t. 147.

Alca major. *Brif.* vi. p. 85. 1. t. 7.—*Ib.* 8vo. ii. p. 382.

Le grand Pengoin. *Buf.* ix. p. 393. t. 29.

Great Auk. *Br. Zool.* ii. N° 229. t. 81.—*Ib.* fol. 136. *Arct. Zool.* ii. N° 424. *Lath. Syn.* v. p. 311. 1. *Ind. Orn.* ii. p. 791. N° 1. *Lewin, Br. Birds,* vi. t. 222. *Walc. Syn.* i. t. 86.

The length of this ſpecies is three feet. The bill is black, very ſtrong, compreſſed, and marked with ſeveral furrows. The baſe of the upper mandible is covered with ſhort velvet-like feathers; between the bill and the eye is a large patch of white; the head, neck, back, and wings, gloſſy black; leſſer quill-feathers tipped with white; legs black.

The ſmallneſs of the wings renders them uſeleſs for flight, the longeſt quill-feather not exceeding four inches in length. Theſe, however, are admirably adapted to its mode of life, and are of peculiar uſe in diving under water, where they act as fins; by which means it purſues its prey with aſtoniſhing velocity.

This bird is only found in the moſt northern parts of the kingdom; is ſaid to breed in the iſle of St. Kilda. Like the reſt of this genus, it lays only one egg, white, ſometimes irregularly marked with

with purpliſh lines, or blotched with ferruginous and black at the larger end; length ſix inches.

AUK-LITTLE.

Alca Alle. *Lin. Syſt.* i. p. 211. 5. *Gmel. Syſt.* ii. p. 554.

Mergulus melanoleucos roſtro acuto brevi. *Raii Syn.* p. 125. A. 5. *Will.* p. 261. t. 59.

Small black and white Diver. *Will. Angl.* p. 343. *Edw.* t. 91.

Greenland Dove. *Albin.* i. t. 85.

Le petit Guillemot. *Buf.* ix. p. 354.

Little Auk. *Br. Zool.* ii. N° 233. t. 82.—*Ibid.* fol. 137. t. H. 4. f. 1. *Arct. Zool.* ii. t. 429. *Lath. Syn.* v. p. 327. 11. *Ind. Orn.* ii. p. 795. 19. *Lewin, Br. Birds,* vi. t. 223. *Walc. Syn.* i. t. 88. *Pult. Cat. Dorſet.* p. 17.

Uria minor. *Briſ.* vi. p. 73. 2.—*Ib.* 8vo. ii. p. 378.

In ſize this bird rather exceeds the blackbird; length near nine inches.

Bill black, ſhort, ſtrong, and ſomewhat rounded in ſhape. The plumage is generally black above and white beneath, with the tips of the ſecondary quill-feathers white; ſometimes the ſcapulars are mixed with white; in others the head and neck is black. Whether any of theſe varieties are owing to ſex or age is not aſcertained: it is ſaid the throat of the male is black. It is not frequently

met with in England, but is common in Greenland, where it breeds, and is ſaid to lay two eggs; if ſo, it differs from others of the genus, which uſually lay but one; theſe are larger than thoſe of a pigeon, of a bluiſh white.

AWL. Vide *Woodpecker-green.*

B.

BALD-BUZZARD. Vide *Oſprey.*
BALD-COOT. Vide *Coot.*
BARGANDER. Vide *Shieldrake.*
BARKER. Vide *Snipe-ſpotted.*
BARLEY-BIRD. Vide *Siſkin.*
BEAM-BIRD.

We are at a loſs to find what bird Mr. Willughby means under this denomination. He ſays it is found in Yorkſhire, and called the Beam-bird, from its neſting under beams in out-buildings. This gentleman, however, calls it Beccafigo, or Fig-eater. Mr. Pennant has conſidered it as the *Motacilla Hippolais* of Linnæus; but his deſcription, which is as follows, by no means anſwers to that bird. "Leſs than the Blackcap. The inſide of the mouth is red; the head, neck, back, and wings, are of an olivaceous aſh colour; the quill-feathers darker, edged with olive; the inner coverts of the wings yellow; breaſt white, tinged with

with yellow; the belly ſilvery white; the tail duſky; the legs bluiſh." Pettychaps, Br. Zool. 149. Moſt certainly this deſcription does not correſpond with either the Greater or Leſſer Pettychaps. From the habit we ſhould be led to believe it might be the Spotted Flycatcher, which in ſome places is called by the name of Rafter, from its neſting on, or under, rafters in old buildings; whereas the Pettychaps invariably repairs to woods and hedges for the purpoſe of nidification.

BEAN-CRAKE. Vide *Gallinula-crake.*

BEE-BIRD. Vide *Flycatcher-ſpotted.*

BEE-EATER. A genus of birds, the characters of which are,

Bill quadrangular, a little incurvated, ſharp-pointed.

Noſtrils ſmall, placed near the baſe.

Tongue ſlender, in ſome ſpecies fringed at the end.

Toes three forward, one backward; the outer toe ſomewhat connected with the middle one.

BEE-EATER-COMMON.

Merops apiaſter. *Lin. Syſt.* i. p. 182. 1. *Gmel. Syſt.* i. p. 460. *Raii Syn.* p. 49. 3. *Will.* p. 110. 10. *Ind. Orn.* i. p. 269. 1. *Briſ.* iv. p. 532.—*Ib.* 8vo. ii. p. 194.

Le Guépier. *Buf.* vi. p. 480. t. 23.

BEE

Bee-eater. *Will. Angl.* p. 147. *Albin.* ii. t. 44. *Lath. Syn.* ii. p. 667. 5.—*Ib. Sup.* p. 119. *Lin. Tranſ.* iii. p. 333.

The Common Bee-eater is the only one, out of twenty-three known ſpecies, that has ever made its appearance in England.

As we never had an opportunity of examining more than one which was killed in this country, which varied but little from the deſcription given by Latham in his *General Synopſis*, we ſhall take the liberty of borrowing it from that author.

It meaſures from bill to tail ten inches. The bill is an inch and three quarters in length, and black; the baſe of the upper mandible covered with dirty-white feathers; the irides are red; the forehead is of a blue-green colour, behind it green; the top of the head cheſnut, tinged with green; hind-head and upper part of the neck cheſnut, growing paler towards the back; from the bill to the hind-head is a black ſtripe, paſſing through the eyes; the back and ſcapulars are very pale yellow, tinged with both cheſnut and green; rump and upper-tail coverts blue-green, with a yellow tinge; the throat is yellow; the under part of the body blue-green, growing paler towards the belly; the leſſer wing coverts are dull green; the quills, for the moſt part, ſea-green without, and many of the inner ones rufous; the firſt very ſhort, the ſecond longeſt of all; the tail is wedge-ſhaped, and con-

ſiſts

ſifts of twelve feathers, the ſhafts of which are brown above, and whitiſh beneath; the two middle feathers are ſea-green with a ſhade of rufous; the reſt the ſame, but margined with cinereous within; the two middle feathers exceed the outer ones by three quarters of an inch; the legs are of a reddiſh brown, claws reddiſh black.

This bird does not appear to have been noticed in England till within theſe few years. In the third volume of the *Linnean Tranſactions* an account is given of one of this ſpecies having been ſhot (for the firſt time in Great Britain) in July 1794, near Mattiſhall in the county of Norfolk; which ſpecimen was exhibited before the Linnean Society. A flight of about twenty was ſeen in June; and the ſame flight probably (much diminiſhed in number) was ſeen paſſing over the ſame ſpot in October following. Since the above period we have been credibly informed more than one have been killed in England.

The *Merops apiaſter* is an inhabitant of various parts of the European continent. They are not uncommon in the ſouth of France and in Italy, as well as in the iſlands of the Mediterranean. It has alſo been ſeen in Germany, and in Sweden, but no where ſo plentiful as in the ſouthern parts of Ruſſia, particularly about the rivers Don and Wolga, in the banks of which they build their neſts, perforating holes to the depth of half a foot for that purpoſe. Are ſaid to be gregarious as well in the breeding ſeaſon as in their migrations, excavating the clayey banks

banks ſo near to each other as to appear like a *honeycomb*. In the autumn they migrate in large flocks to the more ſouthern latitudes.

Theſe birds appear in ſmall flocks at Gibraltar in March, but are ſaid to remain there only a few hours. The neſt is compoſed of moſs; the eggs are ſix or ſeven in number, perfectly white, about the ſize of thoſe of a Stare. Its name has doubtleſs been taken from that inſect of which it is partially fond; but it alſo feeds on moſt winged inſects, which it takes on wing, like the Swallow.

BERNACLE. Vide *Gooſe Brent*, and *Gooſe Bernacle.*

BILCOCK. Vide *Rail-water.*

BITTERN.

Ardea ſtellaris. *Lin. Syſt.* i. p. 239. 21. *Gmel. Syſt.* ii. p. 646. *Raii Syn.* p. 100. A. 11. *Will.* p. 207. t. 50. 52.

Le Butor. *Buf.* vii. p. 411. t. 21.

Bittern. *Br. Zool.* ii. N° 174.—*Ib.* fol. 711. t. A. 1. *Will. Angl.* p. 282. *Albin.* i. t. 68. *Hayes, Br. Birds*, t. 19. *Lath. Syn.* v. p. 56. 17. —*Ib. Sup.* p. 234. *Ind. Orn.* ii. p. 680. 18. *Lewin, Br. Birds*, iv. t. 146. *Walc. Syn.* ii. t. 127.—*Pult. Cat. Dorſet.* p. 14.

Botaurus. *Briſ.* v. p. 444. 24. t. 37. f. 1.—*Ib.* 8vo. ii. p. 327.

PROVINCIAL.

Bittour. Bumpy-coſs. Butter-bump. Miredrum.

This

This ſpecies of Heron is rather leſs than the common Heron; length about two feet and a half.

The bill is four inches long, of a brown horn colour above; the lower mandible, and baſe of the upper, greeniſh; irides yellow. The feathers on the top of the head are black; thoſe on the hind head, neck, and breaſt, are long and looſe; the plumage, in general, is of a dull pale yellow, elegantly variegated with ſpots, and bars of black; the greater coverts and quill-feathers ferruginous, regularly barred with black; tail ſhort; legs pale green; toes and claws very long and ſlender; the middle claw ſerrated on the inner edge. The female is rather leſs, the plumage not ſo bright, and the feathers on the neck not ſo long and flowing as in the male.

The Bittern is by no means a plentiful ſpecies. In the breeding ſeaſon it is only found in the leſs-frequented reedy marſhes, and ſwampy moors well clothed with ruſhes, where it forms a neſt on ſome tump, by collecting a quantity of ſedge or other coarſe plants together. It lays four or five eggs of a light olive-green colour, inclining to cinereous. At this ſeaſon the male makes a ſingular bellowing noiſe, vulgarly ſuppoſed to be produced by the bird putting his bill into a reed. It is rouſed with difficulty from its lurking-place, flies heavily, and frequently lights again at a ſmall diſtance; ſo that it becomes an eaſy prey to the ſportſman. We

are

are informed, however, that ſometimes it ſoars to a prodigious height in the air, with a ſpiral aſcent, making at the ſame time a ſingular noiſe. In the winter theſe birds leave the more mountainous ſwamps, where it is probable the greater part breed, and become more ſcattered in the low moiſt ſituations, and in ſevere weather are found in the ſedgy banks of rivers and ſtreams of water. It is, however, become much more ſcarce than formerly, ſince its fleſh has been accounted a delicacy; and the poulterers value it at not leſs than half a guinea.

The principal food of this bird is ſmall fiſh, frogs, and inſects; the warty lizzard alſo becomes its prey, as we have found by diſſection.

BITTERN-LITTLE.

Ardea minuta. *Lin. Syſt.* i. p. 240. β. *Gmel. Syſt.* ii. p. 646. *Ind. Orn.* ii. p. 683. 27. *Edw.* t. 275.

Little Bittern. *Br. Zool. App.* p. 537. t. 8. *Arct. Zool.* ii. N° 359. *Lath. Syn.* v. p. 65. 27. —*Ib. Sup.* p. 235. *Lewin, Br. Birds,* iv. t. 147. *Walc. Syn.* ii. t. 128. *Don. Br. Birds,* iii. t. 54. *Pult. Cat. Dorſet.* p. 14.

Le Blongios. *Buf.* vii p. 395.

Ardeola nævia. *Briſ.* v. p. 500. 47. t. 40. f. 2. —*Ib.* 8vo. ii. p. 342.

PROVINCIAL.

Boonk. Long-neck.

This

This beautiful ſpecies of Heron is ſcarce larger than a Fieldfare in the body.

The length, to the end of the tail, is fifteen inches. The bill two inches long, duſky at the point, yellow on the ſides. The top of the head, back, and tail, black, gloſſed with green; the fore part of the neck, breaſt, and thighs, buff colour; belly and vent white; at the ſhoulders a large cheſnut ſpot; the larger coverts of the wings whitiſh; the leſſer coverts yellowiſh buff; quill-feathers black; legs and toes duſky green; thighs feathered to the knees; middle claw ſerrated on the inner ſide.

The above is the deſcription of the male. The female is ſuppoſed to be the *Ardea minuta*, Lin. Syſt. 240. 26. It differs from the other ſex in the plumage of the upper part of the body being brown, margined with pale rufous; beneath, the feathers are the ſame, but paler, and more deeply margined.

The eggs are ſaid to be white, about the ſize of thoſe of the blackbird, four or five in number, which are placed on the ground upon a few dried flags.

The Little Bittern is a very rare bird with us. Few inſtances only are recorded of its being killed in England. A male was ſhot near Bath in the autumn of 1789, perched on the ſtump of a tree on the bank of the Avon. It is more frequent in

ſome parts of the European continent, particularly in Switzerland.

BITTOUR. Vide *Bittern.*

BLACKBIRD.

Turdus merula. *Lin. Syſt.* i. p. 295. 22. *Gmel. Syſt.* ii. p. 831. *Raii Syn.* p. 65. A. 1. *Will.* p. 140. t. 37. *Ind. Orn.* i. p. 340. 50.

Merle. *Buf.* iii. p. 330. t. 20. *Briſ.* ii. p. 227. 10. —*Ib.* 8vo. i. p. 221.

Blackbird. *Br. Zool.* N° 109. t. 47.—*Ib.* fol. 92. *Arct. Zool.* ii. p. 345. *Albin.* i. t. 37. *Will. Angl.* p. 190. *Lath. Syn.* iii. p. 43. 46.—*Ib. Sup.* p. 141. *Lewin, Br. Birds,* ii. t. 61. *Walc. Syn.* ii. t. 210.—*Pult. Cat. Dorſet.* p. 10.

PROVINCIAL.

Amſel.

This well-known ſpecies of Thruſh needs little deſcription. The male is wholly of a deep black, when it has attained maturity, which is not till towards the return of ſpring, at which time the bill and orbits of the eyes are yellow: the young and females are of a dark ruſty brown; the bill and eyelids duſky.

The ſong of the blackbird is a ſhrill kind of whiſtle of various notes which it commences early in the ſpring. It is an early breeder, and prepares a neſt compoſed

compoſed externally of green moſs, fibrous roots, and other like materials; the inſide is plaſtered with earth and afterwards lined with fine dry graſs; the neſt is uſually placed in a thick buſh, or againſt the ſide of a tree, or on a ſtump in the ſide of a bank. It lays four or five light blue eggs, thickly covered with pale ferruginous brown ſpots, moſtly at the larger end; which are hatched in about fourteen days incubation.

Its food is principally worms and ſhelled ſnails; theſe laſt are moſt dexterouſly broken againſt a ſtone in order to get at the animal; all kinds of inſects as well as fruit are eagerly ſought after. In confinement they readily eat crumbs of bread, and fleſh either raw or otherwiſe. It is never obſerved to migrate with us, or to congregate; but is at all ſeaſons a ſolitary ſpecies, preferring woods and encloſed ſituations.

BLACKBIRD-MICHAELMAS. Vide *Ouzel-ring*.

BLACKCAP. Vide *Titmouſe-great*, *Titmouſe-marſh*, and *Gull-black-headed*.

BLACKCAP.

Motacilla atricapilla. *Lin. Syſt.* i. p. 332. 18. *Gmel. Syſt.* ii. p. 970.

Atricapilla ſeu Ficedula Aldrov. *Raii Syſt.* p. 79. A. 8. *Will.* p. 162. t. 41.

Sylvia Atricapilla. *Ind. Orn.* ii. p. 508. 6.

La Fauvette à tête noire. *Buf.* v. p. 125. t. 8. f. 1.

Blackcap. *Br. Zool.* i. N° 148.—*Ib.* fol. 101. t. 8. f. 5. *Arct. Zool.* ii. p. 418. F. *Will. Angl.* p. 226. *Lath. Syn.* iv. p. 415. 5. *Lewin, Br. Birds,* iii. t. 116. *Walc. Syn.* ii. t. 234. *Pult. Cat. Dorſet.* p. 9.

Curruca atricapilla. *Briſ.* iii. p. 380. 6.—*Ib.* 8vo. i. p. 416.

PROVINCIAL.

Mock-nightingale. Nettle-creeper. Nettle-monger.

This ſpecies of warbler weighs about four drams and a half: length near ſix inches. The bill is brown; irides dark hazel. The upper part of the head in the male is black, the hind part of the neck cinereous brown; back greyiſh brown with a tinge of green: the quill-feathers and tail duſky, edged with dull green; breaſt and upper part of the belly light aſh colour; legs lead colour.

The female is diſtinguiſhed from the other ſex by the crown of the head, which is of a dull ruſt colour; ſhe is alſo ſuperior in ſize.

The blackcap is a migrative ſpecies viſiting us early in the ſpring, and retiring in September; it frequents woods and thick hedges, and ſeems particularly partial to orchards and gardens, where it delights us with its charming melodious ſong, which is very little inferior to that of the nightingale, except in variety of notes.

It makes a neſt in ſome low buſh or ſhrub, compoſed of dried ſtalks, generally of gooſe-graſs, put together

together with a little wool, and ſometimes a little green moſs on the outſide; the inſide is lined with fibrous roots, upon which are frequently placed a few long hairs. The eggs are four or five in number, of a pale reddiſh brown, mottled with a deeper colour, and ſometimes ſprinkled with a few aſh-coloured ſpots; their weight about thirty-five grains. On the firſt arrival of this bird it feeds greedily on ivy-berries, but forſakes that food as ſoon as the vernal ſun has rouſed the inſect tribe.

BLACKCOCK. Vide *Grous-black.*

BLACKCOCK-SPOTTED. Vide *Grous-black-ſpotted.*

BLACK-GAME. Vide *Grous-black.*

BOONK. Vide *Bittern-little.*

BOTTLETOM. Vide *Titmouſe long-tailed.*

BOTTLE-NOSE. Vide *Puffin.*

BOUGER. Vide *Puffin.*

BRAMBLING. Vide *Bunting-mounting.*

BRAMBLING, or BRAMBLE.

Fringilla montifringilla. *Lin. Syſt.* i. p. 318. 4. *Gmel. Syſt.* ii. p. 902. *Raii Syn.* p. 88. *Briſ.* iii. p. 155.—*Ib.* 8vo. i. p. 349. *Will.* p. 187. t. 45. *Ind. Orn.* i. p. 439. 7.

Le Pinçon d'Ardennes. *Buf.* iv. p. 124.

Brambling, *or* Mountain Finch. *Br. Zool.* N° 126. —*Ib.* fol. 108. t. V. f. 4. *Arct. Zool.* ii. p. 381. E. *Albin,* iii. t. 64. *Will. Angl.* p. 254. t. 45. *Lath. Syn.* iii. p. 261. 13.

Hayes, Br. Birds, t. 36. *Lewin, Br. Birds,* ii. t. 80. *Walc. Syn.* ii. t. 218. *Don. Br. Birds,* iv. t. 85. *Pult. Cat. Dorſet.* p. 12.

PROVINCIAL.

Kate.

This ſpecies of Finch is rather larger than the Chaffinch. Length about ſix inches.

The bill is yellow, tip black; irides duſky. The head, hind part of the neck, and back, black; ſome of the feathers edged with ruſty brown and aſh-colour, which laſt predominates on the back of the head and ſide of the neck; the throat, breaſt, and upper coverts of the wings, ferruginous orange; middle coverts yellowiſh white; the greater coverts black, tipped with orange; quill-feathers black, edged with yellow on their exterior webs; on three or four of the primores a ſpot of white runs through the whole of the exterior web, and forms an oblique bar when the wing is cloſed; the belly and rump white; on the ſides above the thighs are a few round black ſpots; the tail is a little forked, the exterior feather white on the outer web, the others black, except the two middle, which are edged and tipped with aſh-colour; legs greyiſh brown.

The female is much leſs bright in the colour of the plumage: the ſide of the head and back of the neck is grey; on the latter are two duſky lines paſſing

paſſing from the head downwards; the top of the head and back are duſky, each feather deeply margined with grey, which gives thoſe parts a pretty mottled appearance; the rufous on the breaſt and wings is very faint, but the markings correſpond with thoſe of the other ſex.

We have not been able to diſcover that this bird has ever bred with us, but they are frequently ſeen in the winter, in large flocks, upon the coaſt of Kent and Suſſex when the weather is ſevere, and have been ſo exhauſted as to ſuffer themſelves to be taken up. They are alſo found in the interior parts of the kingdom at that ſeaſon, flying in company with Chaffinches and Yellowhammers. By obſervation on thoſe we have kept in confinement, it is a hardy, bold bird, feeding on moſt kinds of ſeed uſually given to caged birds; is ſaid to be particularly fond of beech maſt. It is common in many parts of the European continent, but moſt probably breeds only in the northern parts; is ſaid to build in fir-trees, and form a neſt with moſs, lined with wool and feathers, and to lay four or five yellowiſh ſpotted eggs.

BRANTAIL. Vide *Redſtart.*

BROOK-OUZEL. Vide *Rail-water.*

BUMPYCOSS. Vide *Bittern.*

BUNTING. A genus of birds, the characters of which are,

Bill ſtrong and conic, the ſides of each mandible bending inwards; a hard knob in the roof of the upper mandible.

BUNTING-CIRL.

Emberona Cirlus. *Lin. Syſt.* i. p. 311. 12. *Gmel. Syſt.* ii. p. 879. *Raii Syn.* p. 93. 4. *Will.* p. 196. *Ind. Orn.* i. p. 401. 10.

Le Bruant de haye. *Buf.* iv. p. 347. *Plan. Enl.* 653. f. 1. 2.

Cirl Bunting. *Lath. Syn.* iii. p. 190 26.

Emberiza ſepiaria. *Brif.* iii. p. 263. 2.—*Ib.* 8vo. i. p. 383.

The length of this ſpecies is ſix inches and a half; weight about ſeven drams. The bill is of a bluiſh lead-colour above, paler beneath; irides hazel. The crown of the head, nape of the neck, and upper part of the breaſt, is of a fine olive-green, the firſt ſtreaked with duſky; from the upper mandible through the eye a duſky ſtroke; above and beneath the eye a bright yellow one; throat black, ſlightly tipped with brown, running into a bar under the yellow on the cheek; beneath this is a gorget of beautiful bright yellow; the back and ſcapulars are of a fine cheſnut-brown, the former marked with duſky ſtreaks, the margin of the feathers with olive; rump olive-brown; upper tail coverts inclining to cheſnut; the ſmaller wing coverts olive-green; greater coverts duſky, their outer webs cheſnut-brown; greater quills duſky, edged with green on the exterior webs; ſmaller quills cheſnut, duſky down their middle; acroſs the breaſt is a band of cheſnut, mixed with yellow; belly and under-tail coverts yellow, the latter duſky on the ſhafts;

ſhafts; ſides more inclining to brown; the under-wing coverts bright yellow; the two middle feathers of the tail cheſnut-brown, the reſt black, except the two exterior on each ſide, which have an oblique bar of white from the tip half way; and the outmoſt feather is white throughout the whole of the exterior web; the legs are brown; claws duſky.

The female in general weighs half a dram leſs than the male.

The upper part of the head is olivaceous brown, ſtreaked with duſky; over the eye a dull yellow ſtreak, paſſing down the ſide of the head; cheeks brown, on which is a yellowiſh ſpot; on each ſide the lower mandible is a broken ſtreak of duſky, paſſing downwards; chin and throat dull yellow, the latter ſtreaked with duſky; the back-part and ſides of the neck and breaſt olivaceous-brown, with duſky ſtreaks; belly and ſides pale yellow, with large duſky ſtreaks on the latter; the upper part of the body and wings like the other ſex, but the colours leſs bright.

We firſt diſcovered this ſpecies near Kingſbridge in the winter of 1800, not uncommon amongſt flocks of Yellow Buntings and Chaffinches, and procured ſeveral ſpecimens of both ſexes, killed in different places ſix or ſeven miles from that place. They are indigenous to Devonſhire, but ſeem to be confined to the ſouthern parts of that county contiguous to the coaſt, having found

them extending as far as Teignmouth, at both of which places we found their neſts; but have never obſerved them far inland. It generally builds in furze, or ſome low buſh; the neſt is compoſed of dry ſtalks, roots, and a little moſs, and lined with long hair and fibrous roots. The eggs are four or five in number, cinereous white, with irregular long and ſhort curved duſky lines, terminating frequently with a ſpot at one end; ſize rather inferior to thoſe of the Yellow Bunting, to which it bears great reſemblance. Theſe birds pair in April, and begin laying early in May.

The female might readily be miſtaken for that ſex of the Yellow Bunting at a little diſtance, but is materially different when compared, eſpecially in the cheſnut-colour of the upper parts of this bird. The note is alſo ſimilar to that of the Yellow Bunting, but ſhorter, not ſo ſhrill, and the latter part not drawn out to ſuch a length.

It is remarkable that ſo common a bird as the Cirl Bunting ſeems to be in the weſt of England, ſhould have ſo long eſcaped the notice of Britiſh naturaliſts; but in all probability this has been occaſioned by their locality. It is ſaid to be only found on the continent in the warmer parts of France and Italy; ſo with us it ſeems confined to the mildeſt part of England; but the winter of 1800, which was ſevere in Devonſhire, did not force them to ſeek a warmer climate, but, on the contrary, they continued gregarious with other ſmall birds, ſearching their food amongſt the ploughed lands.

BUNTING-COMMON.

Emberiza miliaria. *Lin. Syſt.* i. p. 308. 3. *Gmel. Syſt.* ii. p. 868. *Raii Syn.* p. 93. A. 1. *Will.* p. 195. t. 40. *Ind. Orn.* i. p. 402. 12.

Le Proyer. *Buf.* iv. p. 355. t. 16.

Bunting. *Br. Zool.* N° 118.—*Ib.* fol. iii. t. W. f. 7. *Arct. Zool.* ii. p. 366. B. *Albin. Br. Birds,* ii. t. 50. *Will. Angl.* p. 267. t. 40. *Lewin, Br. Birds,* ii. t. 74. *Lath. Syn.* iii. p. 171. *Walc. Syn.* ii. t. 213. *Don. Br. Birds,* iii. t. 50. *Pult. Cat. Dorſet.* p. 11.

Cynchramus. *Briſ.* iii. p. 292. 10.—*Ib.* 8vo. i. p. 390.

PROVINCIAL.

Bunting-lark. Ebb.

The weight of this ſpecies is nearly two ounces; length ſeven inches and a half. Bill brown; irides dark hazel; head and upper parts light brown, inclining to olive; the under parts yellowiſh white, with a duſky ſtroke down the ſhaft of each feather, except on the belly, which is quite plain; quill-feathers duſky, with lighter edges; tail ſomewhat forked and duſky; the legs are of a dull light yellow.

Male and female alike.

The Common Bunting ſeems to delight in champaign countries abounding with corn, and is rarely found in uncultivated parts, or graſs fields, diſtant from arable land. In the winter theſe birds become

become gregarious, and ſometimes viſit the farmers yards. When it ſings (if it can be called a ſong) it generally aſſumes the loftieſt branches of a buſh, or ſome low tree; or even perched on the top of a dock it cheers its patient incubating mate with a ſcreaming, inharmonious note. The neſt is placed on the ground, formed externally of ſtraw, lined with fibrous roots or dry graſs, ſometimes finiſhed with long hairs. The eggs are generally four in number, of a dirty white, ſpotted and veined with reddiſh brown and aſh-colour; their weight about one dram.

Theſe birds are ſometimes brought to market, and ſold for larks, to which they are little or nothing inferior, but are eaſily diſtinguiſhed by the form of the bill, and the tooth-like knob in the roof of the mouth, by the moſt common obſerver.

BUNTING-GREEN-HEADED.

Emberiza chlorocephala. *Gmel. Syſt.* ii. p. 887.
Emberiza Tunſtalli. *Ind. Orn.* i. p. 418. 69.
Green-headed Bunting. *Lath. Syn.* iii. p. 211. 61. *Lewin, Br. Birds,* ii. t. 76. *Brown, Ill.* p. 74. t. 30.

This ſpecies has the head and neck dull olive-green; back and wing-coverts duſky brown, mixed with black; paler on the rump; the reſt of the wings, breaſt, and belly, deep brown; tail brown, forked; legs yellowiſh.

A bird

A bird of this deſcription was caught in London, a figure of which is given in Brown's Illuſtration of Zoology.

Dr. Latham mentions another in the collection of Mr. Tunſtal; but, in his Index Ornithologicus, he ſeems to doubt its being a diſtinct ſpecies, in which we moſt readily agree, as no other inſtances appear to be recorded than thoſe above mentioned. From which circumſtance we are inclined to believe this bird is no other than an accidental variety of the female Yellow Bunting; one of which we have, with the whole head and neck greeniſh yellow.

BUNTING-MOUNTAIN.

Emberiza Montana. *Gmel. Syſt.* ii. p. 867. *Ind. Orn.* i. p. 398. 3.

Leſſer Mountain Finch, *or* Brambling. *Will. Angl.* p. 255. *Morton's North.* p. 423. t. 13. f. 1.

Mountain Bunting. *Br. Zool.* N° 123.—*Ib.* fol. 113. *Lath. Syn.* iii. p. 165. 3.

This ſpecies is deſcribed to have the bill ſhort, ſtrong, and yellow, the point black; the forehead is dark cheſnut, lighter on the hind part of the head and cheeks; the hind part of the neck and back aſh coloured; the latter more ſpotted with black; throat white; breaſt and belly waved with flame colour; at the ſetting-on of the wing grey; the five firſt feathers blackiſh brown, the reſt white, the point of each daſhed with brown; the three

outer tail feathers white, the reſt dark brown; feet black; the hind claw as long again as the others: the breaſt of the female is of a darker colour than that of the male.

Linnæus ſeems to have conſidered this bird as a variety of the Snow Bunting; others have given it as a diſtinct ſpecies: on this ſubject we dare not hazard an opinion, never having had an opportunity of examining the bird in queſtion.

It is ſaid to be found in Yorkſhire and Northamptonſhire, as alſo in Lincolnſhire.

BUNTING-REED.

Emberiza Schœniclus. *Lin. Syſt.* i. p. 311. 17. *Gmel. Syſt.* ii. p. 881.

Paſſer torquatus. *Raii Syn.* p. 93. A. 3.

S. Arundinaceus. *Will.* p. 196. *Ind. Orn.* i. p. 402. 13.

Reed Bunting. *Br. Zool.* N° 120.—*Ib.* fol. 112. t. w. *Arct. Zool.* ii. p. 368. E. *Albin.* ii. t. 51. *Hayes, Br. Birds,* t. 35. *Lewin, Br. Birds,* ii. t. 75. *Lath. Syn.* iii. p. 173. 9.—*Ib. Sup.* p. 157. *Walc. Br. Birds,* t. 214.

Ortolan de voſeaux. *Buf.* iv. p. 315. *Briſ.* iii. p. 274. 5.—*Ib.* 8vo. i. p. 386.

PROVINCIAL.

Reed Sparrow. Water Sparrow.

The weight of this ſpecies is near five drams and a half; length ſix inches.

The

The bill is duſky; irides hazel; the head, chin, and throat black; at the corner of the mouth commences a white ring which grows broader behind the ears, and encircles the head; the breaſt and belly white; the ſides grey, marked with a few dark brown ſtrokes; the back is black, deeply bordered with reddiſh brown, interſperſed with grey, which grows more conſpicuous towards the rump; quill-feathers, and coverts of the primores duſky, edged with tawny red; the tail is black; the two middle feathers deeply bordered with rufous, the two exterior on each ſide marked obliquely with white towards the end; the ſhafts and tips black.

The female is rather leſs; the head is rufous brown ſtreaked with duſky; from each ſide of the under mandible a duſky line paſſes under the neck, where it joins and forms a bed of that colour; behind the eye a light coloured ſtroke; the breaſt is ſtreaked with reddiſh brown; the rump plain olive brown; it has no white ring round the head as in the male.

The young male birds do not aſſume their full black head till the enſuing ſpring; nor is the white ring ſo conſpicuous.

It is ſomewhat extraordinary that the manners and habits of ſo common a bird ſhould remain ſo long in obſcurity; even modern authors tell us it is a ſong bird, that it ſings after ſunſet; and deſcribe its neſt to be ſuſpended over the water faſtened between three or four reeds. There can be no doubt

doubt however, that the neſt, as well as the ſong of the Sedge Warbler, have been taken and confounded for thoſe of this bird; for as they both frequent the ſame places in the breeding ſeaſon, that elegant little warbler is pouring forth its varied notes concealed in the thickeſt part of a buſh; while this is conſpicuouſly perched above, whoſe tune is not deſerving the name of ſong; conſiſting only of two notes, the firſt repeated three or four times, the laſt ſingle and more ſharp.

This inharmonious tune it continues to deliver with ſmall intervals from the ſame ſpray, for a great while together when the female is ſitting.

The neſt is moſt commonly placed on the ground near water; ſometimes it builds in a buſh ſome diſtance from the ground; at other times in high graſs, reeds, ſedge, or the like, and even in furze at a conſiderable diſtance from any water; in all theſe ſituations we have met with it, but never faſtened or ſuſpended as authors have related. The neſt is compoſed of ſtalks of graſs, or other dry vegetable ſubſtances; ſometimes partly moſs, and lined with fine graſs; frequently finiſhed with long hair.

The eggs, which are four or five in number, weigh about thirty-ſix grains, and are of a dirty bluiſh white, or purpliſh brown, with numerous dark-coloured ſpots and veins, much reſembling thoſe of the Chaffinch. We ſhall here remark that the eggs of very diſtinct ſpecies of birds are ſometimes very ſimilar, and not eaſily aſcertained; they are alſo ſubject to conſiderable variation: the neſt is a much

greater

greater mark of diſtinction; the materials with which they are compoſed ſeldom vary.

BUNTING-SNOW.

Emberiza nivalis. *Lin. Syſt.* i. p. 308. 1. *Gmel. Syſt.* ii. p. 866. *Ind. Orn.* i. p. 397. 1.

Hortulanus nivalis. *Briſ.* iii. p. 285. 9.—*Ib.* 8vo. i. p. 388.

L'Ortolan de neige. *Buf.* iv. p. 329.

Pied Mountain Finch. *Albin.* iii. t. 71.

Snow Bunting. *Br. Zool.* i. N° 122. t. 50. *Arct. Zool.* ii. N° 222. *Edw.* t. 126. *Walc. Br. Birds,* ii. t. 210. *Lath. Syn.* iii. p. 161. A. B. C. *Ib. Sup.* p. 157. *Lewin, Br. Birds,* ii. t. 71.

Pied Chaffinch. *Albin.* iii. t. 54.

PROVINCIAL.

Snow-bird. Snow-flake.

This ſpecies is ſuperior in ſize to the Chaffinch; weight about one ounce and a quarter. The bill is black; the forehead and crown white, mixed with black on the hind head; back black; rump white; greater quill-feathers black, at the baſe white; ſecondaries white, with a black ſpot on their inner webs; baſtard wing and ends of the greater coverts white; the whole under parts, from chin to tail, pure white; the middle feathers of the tail black; the three outer ones white, with a duſky ſpot near their ends; legs black.

The

The plumage of this ſpecies is ſubject to very great variety; in ſome the white on the upper parts is more predominant.

Theſe birds appear in the north of Scotland in large flocks during the winter. Some few are ſaid to breed upon the higheſt mountains with the Ptarmigans, and, like thoſe birds, probably change their plumage with the ſeaſons.

It is rarely ſeen in the ſouth of England.

The eggs are ſaid to be reddiſh white, ſpotted with brown.

BUNTING-TAWNY.

Emberiza muſtelina. *Gmel. Syſt.* ii. p. 867.

Emberiza glacialis. *Ind. Orn.* i. p. 398. 2.

Montifringilla calcaribus Alaudæ major. *Raii Syn.* p. 88. A. *Will.* p. 187. t. 77.

Great pied Mountain Finch. *Will. Angl.* i. p. 255. t. 77. *Albin.* iii. t. 71.

Tawny Bunting. *Br. Zool.* i. N° 121.—*Ib.* fol. 112. f. 6. *Lath. Syn.* iii. p. 164. 2. *Lewin, Br. Birds,* ii. t. 72. *Walc. Syn.* ii. t. 211. *Don. Br. Birds,* ii. t. 27.

This bird has by ſome been called Sea Lark and Brambling.

The length of this ſpecies is ſix inches and three quarters. Bill ſhort, yellow, point black; crown of the head tawny; the neck the ſame, but lighter; under part white, in ſome daſhed with yellowiſh marks,

marks, and the breaſt dull yellow; the back and ſcapulars black-edged, with pale reddiſh brown; rump and upper-tail coverts part white and part yellow; the ſix firſt quill-feathers duſky, white at the baſe; the next ſeven become gradually whiter; the two next wholly white; the reſt of the quills and baſtard wings are black, edged with rufous brown; the greater coverts above the ſecondaries are white, forming a bed of that colour on the wing; the four middle tail-feathers black, edged with white; the three outer ones white, the two firſt with a ſmall duſky ſpot on the outer web; the third on both webs at the tip; legs black; hind claw very long.

This appears to be a rare ſpecies, but has been ſometimes met with in the northern parts of England.

BUNTING-YELLOW.

Emberiza Citrinella. *Lin. Syſt.* i. p. 309. 5. *Gmel. Syſt.* ii. p. 870. *Raii Syn.* p. 93. A. 2. *Will.* p. 196. t. 40. *Ind. Orn.* i. p. 400. 7.

Le Bruant. *Buf.* iv. p. 342. t. 8.

Yellow Bunting. *Br. Zool.* N° 119. t. 50.—*Ib.* fol. p. 112. *Arct. Zool.* ii. p. 367. C. *Albin.* i. t. 66. *Will. Angl.* p. 268. t. 40. *Lath. Syn.* iii. p. 170. *Lewin, Br. Birds,* ii. t. 73. *Walt. Syn.* ii. t. 212. *Don. Br. Birds,* v. t. 119.

Emberiza flava. *Brif.* iii. p. 258. 1.—*Ib.* 8vo. i. p. 382.

PROVINCIAL.

Yellow-hammer.

The weight of this ſpecies is about ſeven drams; length ſix inches.

Bill duſky bluiſh; irides hazel; the crown of the head, throat, and belly, are of a beautiful bright yellow; the back part and ſides of the head tinged with green; the breaſt in ſome is marked with reddiſh brown; quill-feathers duſky, the primores edged on their exterior webs with greeniſh yellow, the ſecondaries with ruſty brown, thoſe next the body, the greater coverts, and back, duſky, deeply margined with the ſame, the latter daſhed with green; the rump and upper-tail coverts tawny red; the tail a little forked, duſky, edged with greeniſh yellow; the two outer feathers marked with white on the exterior webs; legs yellow brown.

The female has much leſs yellow about the head, and the colours in general leſs vivid. The Yellow Bunting is ſubject to ſome variety in plumage. We have a ſpecimen in which the whole head and neck is of a light yellow; ſome of the quill-feathers and ſcapulars white; and the under parts and rump pale yellow. The young birds have no yellow about them when firſt they leave the neſt.

This is one of the moſt common indigenous

birds of this country; if it was more rare its beauty would be leſs diſregarded. Its ſong, however, is as little attractive as that of the Common Bunting, poſſeſſing only a repetition of the ſame note five or ſix times ſucceſſively, terminating in one more lengthened and ſhrill.

In winter it aſſembles in flocks with other granivorous birds, and picks up the ſcattered grain diſperſed by the bountiful flail, and not by the maſter of the hoarded ſheaves, who knows too well the value of his auriferous ſtore.

It does not breed till late in the ſpring. The neſt is generally placed near the ground, in ſome low buſh or hedge. It is compoſed of ſtraw and various dried ſtalks, and lined with fine dry graſs, finiſhed with long hair.

The eggs differ ſomewhat in colour and ſize; ſome are nearly white, others have a purpliſh hue, but are more or leſs marked with hair-like ſtreaks, terminating with a roundiſh ſpeck; the number from three to five, but uſually four; their weight from thirty to forty-ſeven grains.

BUSTARD. A genus of birds, the characters of which are,

Bill ſtrong, a little incurvated.

Toes, three before, none behind.

Legs long, and naked above the knees.

BUSTARD-GREAT.

Otis Tarda. *Lin. Syſt.* i. p. 264. 1. *Gmel Syſt.*

ii. p. 726. *Raii Syn.* p. 58. A. 1. *Will.* p. 129. t. 32. *Brif.* v. p. 18. 1. *Ind. Orn.* ii. p. 658. 1.

Outarde. *Buf.* ii. p. 1. t. 1.

Great Buſtard. *Br. Zool.* i. N° 98. t. 44.—*Id.* fol. 87. t. N. *Artt. Zool.* ii. N° 186. *Will. Angl.* p. 178. t. 32. *Edw.* t. 79. 80. *Albin.* iii. t. 38. 39. *Lath. Syn.* iv. p. 796. *Oſb. Men.* p. 33. 34. M. F. *Lewin, Br. Birds,* iv. t. 139. *Walc. Syn.* ii. t. 173. *Pult. Cat. Dorſet.* p. 6.

This is the largeſt of the Britiſh birds, ſometimes weighing as much as thirty pounds. The bill is duſky; irides light hazel; head and neck aſh-coloured, inclining to brown on the top of the head; the lower part of the neck behind almoſt bare of feathers; the back and leſſer coverts of the wings elegantly barred with black and light ruſt colour; greater coverts pale cinereous; quills black, ſlightly tipped with white; belly white; the tail conſiſts of twenty feathers, the middle ones are of a light ruſt colour, barred with black; the two outer ones almoſt white, with two or three ſmall bars of black; legs duſky brown.

The female weighs about ten or twelve pounds. The crown of the head is deep orange-brown, croſſed with tranſverſe black lines; the reſt of the head brown; lower part of the neck before aſh-coloured, in other reſpects like the male, except in

not

not having the long tuft of feathers growing from each ſide the lower mandible, ſo conſpicuous in that ſex when arrived at maturity.

The male has alſo a very capacious pouch, ſituated along the fore part of the neck, the entrance of which is under the tongue, capable of holding ſeveral quarts of water; it is ſaid not leſs than ſeven. The ſize, however, of this pouch, or bag, ſeems to be ſomewhat exaggerated, for we think it impoſſible the bird could fly with ſuch an addition of weight before its wings, which would throw it out of the centre of gravity. We ſee the Heron, and many other birds, obliged to extend their legs behind, and contract their necks when flying, in order to balance themſelves on wing. Seven quarts of water is nearly equal to fourteen pounds weight, and certainly more than the bird could carry in that ſituation. The pouch, however, is large, as may be ſeen in the Leverian Muſeum. This, however, is only diſcoverable in adults, as it is moſt likely intended for the purpoſe of furniſhing the female and young, in the breeding ſeaſon, with water, which, in general, is only to be procured at a diſtance, upon the dry and extended downs they inhabit.

The Buſtard is only found upon the large extenſive plains, and are almoſt extinct, except upon thoſe of Wiltſhire, where they are become very ſcarce within theſe few years. It is an extremely ſhy bird, and difficult to be ſhot. Young ones

have frequently been taken by the ſhepherds dogs before they are capable of flight; and their eggs are eagerly ſought after for the purpoſe of hatching under hens. Half-a-guinea is no unuſual price for an egg, and ten or twelve guineas a pair for young birds not full grown. A perſon at Tilſhead, contiguous to the downs in Wiltſhire, has reared a great many in this way; and the conſequence will be a total extinction in a few years.

Theſe birds pair early in the ſpring. The female lays two eggs on the bare ground, which are about four weeks hatching. The young follow the dam ſoon after they are excluded from the egg, but are not capable of flying for ſome time. The egg is larger than that of a Turkey, of an olive-brown colour, blotched with pale ferruginous and aſh-coloured ſpots. Their food is green corn, the tops of turnips, and various other vegetables, as well as worms; but they have been known to eat frogs, mice, and young birds of the ſmaller kind, which they can ſwallow whole.

The Buſtard is not properly migratory with us; it only leaves its uſual haunts in very ſevere winters, when the downs are covered with ſnow for ſome time; preſſed by hunger it repairs to the more encloſed and ſheltered ſituations in ſmall flocks, and even ſtray to a great diſtance. In the winter of 1798 one was killed near Plymouth in Devonſhire, and two others the following year in the ſame county, as well as in other places, to our knowledge.

ledge. Theſe occaſional migrations always prove fatal; ſo large an object ſoon attracting notice, it rarely eſcapes the number of purſuers Theſe birds were formerly found in the Wolds of Yorkſhire, and even as far north as Scotland. It is common in ſome parts of Ruſſia and the deſerts of Tartary, as alſo in ſome parts of Germany.

BUSTARD-LITTLE.

Otis Tetrax. *Lin. Syſt.* i. p. 264. 3. *Gmel. Syſt.* ii. p. 725. *Ind. Orn.* ii. p. 659. 3.
Otis Minor. *Raii Syn.* p. 59. 2. *Will.* p. 129. t. 32. *Briſ.* v. p. 24. 2. t. 2. f. 1. 2.
Petite Outard. *Buf.* ii. p. 40.
Field Duck. *Albin.* iii. t. 41.
Little Buſtard. *Br. Zool.* i. N° 99. *Arct. Zool.* ii. p. 321. A. *Edw.* t. 251. *Lath. Syn.* iv. p. 799. 2.—*Ib. Sup.* p. 226. *Lewin, Br. Birds,* iv. t. 140. *Walc. Syn.* ii. t. 174.

This ſpecies is about the ſize of a Pheaſant; length ſeventeen inches.

The bill is light brown; the crown of the head black, mixed with ferruginous; ſides of the head, chin, and throat, rufous white; the neck black, encircled with an irregular band of white on the upper part, and another near the bottom; the back, ſcapulars, and wing coverts, ferruginous, mottled with brown, and croſſed with fine irregular black lines; greater quills black, white at the baſe; ſe-

condaries white; breaſt and belly white; the tail conſiſts of eighteen feathers, the four middle ones are tawny, barred with black, the reſt white; legs grey.

The neck of the female is of the ſame colour as the back; the breaſt rufous white, ſtreaked with duſky; in other reſpects like the other ſex.

The Little Buſtard has been found rarely in England, three or four inſtances only, and thoſe all females, except one, which ſeems to be accounted for by its being polygamous, and that there are more of that ſex. It is found in many parts of the European continent, particularly in France, where it is frequently taken for the table, and eſteemed as a delicacy. The eggs are ſaid to be of a green colour, four or five in number.

In the Supplement to the *General Synopſis* mention is made of one having been ſhot in Suſſex that had the plumage of a female, but proved otherwiſe on diſſection; probably a bird of the firſt year.

BUSTARD-THICK-KNEED.

Charadrius Œdicnemus. *Lin. Syſt.* i. p. 255. 10. *Gmel. Syſt.* ii. p. 689. *Raii Syn.* p. 108. A. 4—105. A. 6. *Will.* p. 216—227. t. 58.

Otis Œdicnemus. *Ind. Orn.* ii. p. 661. 11.

Le grand Pluvier. *Buf.* viii. p. 105. A. 6. *Brif.* v. p. 76. 12. t. 7. f. 1.

Stone Curlew. *Albin.* i. t. 69. *Will. Angl.* p. 306. 293. t. 58. 57.

Thick-

Thick-kneed Buſtard. *Br. Zool.* i. N° 100.—*Ib.* fol. 127. *White's Selb.* 4to. p. 43—88. *Lath. Syn.* iv. p. 806. 9. *Lewin, Br. Birds,* iv. t. 141. *Walc. Syn.* ii. t. 163.

PROVINCIAL.

Norfolk Plover.

The weight of this ſpecies is about ſeventeen ounces; length eighteen inches.

The bill is almoſt two inches long, duſky at the point, yellow at the baſe; irides and orbits pale yellow. Behind the eye a ſmall ſpace bare of feathers, of a yellowiſh green, moſtly concealed by the ear coverts; the feathers on the head, neck, and whole upper parts, duſky down their middle, deeply bordered with pale tawny-brown; above and beneath the eye is a pale ſtroke; a band of the ſame acroſs the coverts of the wings; the quills black; the two firſt marked with a broad bar of white acroſs each web; the ſeventh and eighth ſlightly tipped with white; breaſt and belly yellowiſh white, the former marked with longitudinal duſky ſtreaks; the tail conſiſts of twelve feathers deeply tipped with black, except the two middle ones; the three outer are barred with black and white, the others with brown; legs long, yellow; toes ſhort; the outer toe connected to the middle one, as far as the firſt joint, by a membrane; claws black.

This

This is a migrative ſpecies, making its firſt appearance with us the latter end of April, or beginning of May, when the male is heard to make a very loud ſhrill note, particularly in the duſk of the evening.

It frequents open, hilly ſituations, moſtly large corn-fields, heaths, or warrens.

It makes no neſt, but lays two eggs on the bare ground; theſe are of a light-brown colour, blotched and ſtreaked with duſky; their weight about an ounce and a half.

Their food is principally inſects and worms, but are ſaid to devour mice, frogs, and toads.

In the autumn they aſſemble in ſmall flocks, preparatory to their departure, and are ſeldom ſeen with us after the beginning of October. We do not recollect having ſeen this bird in the north of England, nor in the weſtern counties farther than Dorſetſhire; but is not uncommon to many of the ſouthern and eaſtern parts.

BUTCHER-BIRD CINEREOUS. Vide *Shrike cinereous.*

BUTCHER-BIRD GREATER. Vide *Shrike cinereous.*

BUTCHER-BIRD LEAST. Vide *Titmouſe bearded.*

BUTCHER-BIRD RED-BACK. Vide *Skrike red-back.*

BUTTER-BUMP. Vide *Bittern.*

BUTTERFLIP. Vide *Avoſet.*

BUZZARD. Several ſpecies of the Falcon genus go under this denomination. Vide *Buzzard-common, Oſprey, Honey Buzzard, Moor Buzzard.*

BUZZARD-COMMON.

Falco Buteo. *Lin. Syſt.* i. p. 127. *Gmel. Syſt.* i. p. 265. *Raii Syn.* p. 16. A. 1. *Will.* p. 38. t. 6. 1. *Ind. Orn.* i. p. 23. 47. *Briſ.* i. p. 406. —*Ib.* 8vo. p. 206.

La Buſe. *Buf.* i. p. 206. t. 8.

Buzzard. *Br. Zool.* i. N° 54. t. 25.—*Ib.* fol. t. A. 3. *Arct. Zool.* ii p. 224. I. *Will. Angl.* p. 70. *Albin.* i. t. 1. *Lath. Syn.* i. p. 48.—*Ib. Sup.* p. 14. *Lewin, Br. Birds,* i. t. 6. *Walc. Br. Birds,* i. t. 6. *Pult. Cat. Dorſet.* p. 3.

PROVINCIAL.

Puttock.

This ſpecies of Falcon is ſubject to great variety in plumage; the males are generally of a lighter colour than the other ſex, and the irides ſeem to correſpond, having ſeen ſome almoſt grey upon the back and wings whoſe irides were light grey. The one now before us is a female; weight thirty-eight ounces, length one foot nine inches, breadth four feet one inch. The bill is black, bluiſh towards the baſe: cere yellow; irides yellow hazel; the upper part of the head, cheeks, and upper part of the

the breaſt, light ruſt-colour; the middle of the feathers dark brown; the lower breaſt blotched with yellowiſh white; the throat and back of the head white, ſtreaked with yellowiſh brown; the back of neck, ſcapulars, and coverts of the wings, tipped and edged on the two laſt with pale ruſt-colour and dirty white; belly white, ſparingly ſpotted with ruſt-colour; the thighs ruſty brown, barred with a deeper ſhade; the greater quills duſky black towards their ends, with a ſhade of aſh-colour on the exterior webs; the interior webs of the primores white towards the baſe, the others barred with dark aſh-colour; the tail is marked alternately with ſix or ſeven bars of duſky and pale ruſt-colour, that near the end is broad and duſky; the whole are tipped with white, and have a daſh of cinereous; legs yellow.

The Buzzard is one of the moſt common ſpecies of Falcon we have; it is a ſluggiſh, inactive bird, ſlow in flight, and ſeldom remains long on wing, except in the breeding ſeaſon, when it will ſoar to a prodigious height, ſpirally aſcending. It makes a neſt in the fork of a tree with large ſticks, and lines it with wood, hair, and other ſubſtances; ſometimes takes poſſeſſion of a deſerted crow's neſt. The eggs are two, and not unfrequently three in number, rather larger than thoſe of a hen, of a dirty white, moſt commonly ſpotted with ruſt-colour, chiefly at the larger end.

This bird never purſues its prey on wing, but

is

is contented with young hares, rabbits, and feathered game, of which it is a great deſtroyer; or if old ones are wounded they become an eaſy prey. In defect of ſuch delicious food it will eat carrion, and even worms and beetles.

C.

CAPERKALLY. Vide *Grous-wood.*

CARGOOSE. Vide *Grebe, great-creſted.*

CHANCHIDER. Vide *Flycatcher, ſpotted.*

CHATTERER. A genus of birds, the characters of which are,

Bill ſtrait, convex, bending towards the point; near the end of the upper mandible a ſmall notch.

Noſtrils hid in reflexed briſtles.

Middle toe connected to the outer at the baſe.

CHATTERER-BOHEMIAN.

Ampelis garrulus. *Lin. Syſt.* i. p. 297. 1. *Gmel. Syſt.* ii. p. 838. *Ind. Orn.* i. p. 363. 1.

Garrulus bohemicus, *and* Silk-tail. *Raii Syn.* p. 85. A.—*Ib.* p. 174. *Will.* p. 90. t. 20.

Le Jaſeur de Boheme. *Buf.* iii. p. 429. t. 26.

Bombycilla bohemica. *Briſ.* ii. p. 333. 63.—*Ib.* 8vo. i. p. 250.

Waxen Chatterer. *Br. Zool.* i. N° 112. t. 48.—*Ib.* fol. 7. t. 1. C. *Arct. Zool.* N° 207. *Edw.* t. 242. *Lath. Syn.* iii. p. 91. 1.—p. 93. 1. A.

Lewin,

Lewin, Br. Birds, ii. t. 65. *Oſt. Men.* t. 15. *Walc. Syn.* ii. t. 204. *Don. Br. Birds*, i. t. 11. *Pult. Cat. Dorſet.* p. 11.

This is the only ſpecies of the genus ever met with in this country, and that not very frequently.

The length of this bird is about eight inches; the ſize near that of a Starling.

The bill is black; irides purpliſh; the head and upper parts vinaceous brown, daſhed with aſh-colour, lighteſt on the rump; the feathers on the crown of the head are long, and form a pointed creſt reclining backwards, of a cheſnut colour; over the eye is a black ſtreak paſſing from the bill to the hind head; chin and throat black; breaſt and belly pale purpliſh aſh-coloured brown, lighteſt towards the vent; the greater coverts of the primores black, tipped with white; greater quills black, the three firſt tipped white, the others with yellow on their outer margins; the ſecondaries tipped on the outer web with white, terminating in flat horny appendages the colour of red ſealing-wax, the number of which varies in different ſpecimens; in that now before us there are five on one ſide, and ſix on the other; but are ſometimes found as many as eight; the tail is black, tipped with yellow, and daſhed with aſh-colour at the baſe; the under coverts of the tail cheſnut; legs black.

The female is ſaid to want the red appendages at the end of the quill-feathers, as alſo the yellow on the wings: all thoſe, however, which have come under

under our inſpection, killed in England, had thoſe characters; but it was not aſcertained whether any of them were females. Dr. Latham, who has conſidered the American ſpecies, or Carolina Chatterer, as only a variety of this, ſays both ſexes have the wings of a plain colour, and the female has no appendages to the quill-feathers. Mr. Pennant informs us theſe birds appear annually about Edinburgh, and feed on the berries of the mountain-aſh. We have received it out of Staffordſhire, and have known others killed in the more ſouthern counties in the autumn or winter.

It is ſuppoſed theſe birds breed in the more northern parts, and are ſaid to neſt in the holes of rocks.

CHOUGH. Vide *Crow red-legged.*

CLAKIS. Vide *Gooſe-bernacle.*

CLATTERGOOSE. Vide *Gooſe-brent.*

COB. Vide *Gull great-black-backed.*

COBBLE. Vide *Diver imber* and *ſpeckled.*

COBLER'S-AWL. Vide *Avoſet.*

COBWEB. Vide *Flycatcher-ſpotted.*

COCK OF THE WOOD, or MOUNTAIN. Vide *Grous-wood.*

CODDYMODDY. Vide *Gull-winter.*

COLEMOUSE. Vide *Titmouſe-cole.*

COLK. Vide *Duck-eider.*

COOT. A genus of birds, the characters of which are,

Bill

Bill ſhort, thick, with a callous extending up the forehead.

Noſtrils narrow and pervious.

Toes furniſhed with broad ſcallopped membranes.

COOT-COMMON.

Fulica atra. *Lin. Syſt.* i. p. 257. 2. *Gmel. Syſt.* ii. p. 702.

Fulica leucorex. *Ib.* ii. p. 703. 21.

Fulica æthiops. *Ib.* ii. p. 704. 22.

——— atra. *Raii Syn.* p. 116. A. 1. *Will.* p. 239. t. 59. *Briſ.* vi. p. 23. t. 2. f. 2.—*Ib.* 8vo. ii. p. 365. *Ind. Orn.* ii. p. 77. N° 1.

Le Foulque ou morelle. *Buf.* viii. p. 211. t. 18.

Common Coot. *Br. Zool.* ii. N° 220. t. 77. —*Ib.* fol. 132. t. F. *Arct. Zool.* ii. N° 416. *Will. Angl.* p. 319. t. 59. *Albin.* i. t. 83. *Walc. Syn.* ii. p. 167. *Lath. Syn.* v. p. 275. 1. —*Ib. Sup.* p. 259. 1. A and B. *Lewin, Br. Birds,* v. t. 195. *Don. Br. Birds,* v. t. 106. *Pult. Cat. Dorſet.* p. 416.

PROVINCIAL.

Bald coot.

This ſpecies ſometimes weighs as much as thir ounces; length eighteen inches.

The bill and callous on the forehead pale fle colour; irides hazel. The whole plumage is bla except on the belly, which is of a duſky aſh-colc

and a line of white on the ridge of the wing; legs and toes dark green.

The Coot is a very common bird in this country; it breeds in many of our lakes, rivers, and large ponds, forming a neſt of flags amongſt the reeds and other aquatic plants cloſe to the ſurface of the water.

The eggs are ſix or ſeven in number, of a dirty white, ſprinkled over with ſmall deep ruſt-coloured ſpots. Many of theſe birds never forſake their breeding-places, even in ſmall pieces of water; and notwithſtanding are frequently rouſed, they are not to be compelled to fly farther than from one ſide to the other. The vaſt flocks which are ſeen in Southampton river, and other ſalt-water inlets, in winter, moſt probably breed farther north; at leaſt a great part of them. At this ſeaſon of the year it is commonly ſold in our markets, frequently ready picked. They look exceedingly white, but the flavour is rather fiſhy. Moſt authors give as a ſpecific character a yellow band, or garter, on the bare part above the knee. This, however, does not always hold good, and may depend on the ſeaſon or ſex.

COOT-FOOT TRINGA. Vide *Phallarope.*

COOT-GREATER.

Fulica aterrima. *Lin. Syſt.* i. p. 258. 3. *Gmel. Syſt.* ii. p. 703. 3. *Ind. Orn.* ii. p. 778. 2.

Fulica major. *Raii Syn.* p. 117. 2. *Will.* p. 239. t. 51.

La grande Foulque, ou la Macroule. *Briſ.* v. p. 28. 2. t. 2. f. 2.—*Ib.* 8vo. ii. p. 366. *Buf.* viii. p. 220.

Greater Coot. *Br. Zool.* ii. N° 221. *Will. Angl.* p. 320. *Lath. Syn.* v. p. 277. 2. *Walc. Syn.* ii. t. 168.

This is larger than the Common Coot, and the plumage of a deeper black; in other reſpects it ſo much reſembles that bird, that ſome doubts may be entertained of its being a diſtinct ſpecies. Said to differ in the colour of the callous on the forehead, and in the band or garter above the knee, the former being white, the latter red. It is rarely met with in England, but is ſaid to be found in Lancaſhire and in Scotland; not uncommon in ſome of the northern parts of the continent.

COOT-MOOR. Vide *Gallinule-common.*

COOT-WEESEL. Vide *Smew red-headed.*

CORMORANT. Vide *Corvorant.*

CORNCREAK, *or* CREK. Vide *Gallinule-crake.*

CORNCRACKER. Vide *Gallinule-crake.*

CORNDRAKE. Vide *Gallinule-crake.*

CORNISH-DAW. Vide *Crow red-legged.*

CORVORANT. A ſpecies of the Pelican genus, the characters of which are,

Bill ſtrong, ſtrait, and either hooked or ſloping at the point.

Noſtrils

Noſtrils either wanting or placed in a longitudinal furrow.

Face naked.

Gullet naked, capable of great diſtention.

Toes four, all webbed together.

Pelicanus Corbo. *Lyn. Syſt.* i. p. 216. 3. *Gmel. Syſt.* ii. p. 573. *Ind. Orn.* ii. p. 886. 14.

Corvus aquaticus. *Raii Syn.* p. 122. A. 3. *Will.* p. 248. t. 63.

Le Cormoran. *Buf.* viii. p. 310. t. 26.

Cormorant. *Will. Angl.* p. 329. t. 63. *Albin.* ii. t. 81.

Corvorant. *Br. Zool.* ii. N° 291.—*Ib.* fol. 159. t. I. 1. *Lath. Syn.* vi. p. 593. 13. *Lewin, Br. Birds*, vii. t. 263. *Walc. Syn.* i. t. 92. *Pult. Cat. Dorſet.* p. 21.

Phalacrocorax. *Briſ.* vi. p. 511. 1. t. 45.—*Ib.* 8vo. ii. p. 495.

PROVINCIAL.

Sea Crow.

The Corvorant is ſubject to much variety, both in ſize and colour. The ſpecimen from which the following deſcription is taken was the largeſt we ever ſaw.

It weighed eight pounds; length three feet three inches; breadth four feet eleven inches; the length of the bill five inches, hooked at the point; irides green.

The chin, and round the baſe of the bill to the eyes, bare and yellow; the head and neck black; back greeniſh black, gloſſed with purple, each feather bordered with fine deep black; ſcapulars and coverts of the wings the ſame, daſhed with aſh-colour; the whole under parts black, except a ſmall patch of white on the throat; quills duſky black; the tail is rounded, and conſiſts of fourteen feathers, which is the great characteriſtic diſtinction of this ſpecies from the Shag.

We have ſeen twenty or thirty Corvorants together: in ſome the upper parts are duſky, daſhed over with aſh-colour; the under parts of the neck and breaſt light brown; the belly dirty white; in ſome the feathers of the head are a little elongated, forming a ſhort pendant creſt, and a patch of white on the thighs.

This variety of markings are, perhaps, common to this ſpecies, without any regard to age or ſex. We have, however, upon diſſection, generally found the females lighteſt coloured.

The Corvorant breeds on moſt of our rocky coaſts, ſometimes in conſort with the Shag. We have ſeen, on the coaſt of Wales, an inſulated rock covered with their neſts, which are compoſed of ſticks and ſea-weed.

The eggs are generally three in number, colour white, weight about two ounces.

It frequently builds on the very ſummit of the higheſt

higheſt rocks impending the ſea; the Shag rarely ſo high.

In the winter it is frequently found in freſh-water rivers, at a conſiderable diſtance from the ſea. We have ſeen eight or ten together far up the river Uſk, perched upon a tree. The Shag is never obſerveed to quit the neighbourhood of the ſea.

It is a great deſtroyer of fiſh; and by frequent diving the wings become ſo wet as to incapacitate its flying, when it walks out of the water, and extends the wings to the wind in order to dry them.

It is ſaid this bird has been trained and uſed for fiſhing, in the manner hawks have for fowling; and in order to prevent their ſwallowing the prey, a collar was put round the neck.

COULTERNEB. Vide *Puffin.*

CRACKER. Vide *Duck-pintailed.*

CRANE.

Ardea Grus. *Lin. Syſt.* i. p. 234. 4. *Gmel. Syſt.* ii. p. 620. *Briſ.* v. p. 374. 6. t. 33.—*Ib.* 8vo. ii. p. 307. *Raii Syn.* p. 95. A. 1. *Will.* p. 200. t. 48. *Ind. Orn.* ii. p. 674. 5.

La Grue. *Buf.* vii. p. 287. t. 14.

Common Crane. *Br. Zool.* ii. *App.* p. 534. t. 6. *Arct. Zool.* ii. p. 453. *Will. Angl.* p. 274. t. 48. *Albin.* ii. t. 65. *Lath. Syn.* v. p. 40. 5. *Lewin, Br. Birds,* iv. t. 143. *Walc. Syn.* ii. t. 124.

This ſpecies of Heron weighs near ten pounds; length five feet.

The bill is above four inches long, of a greeniſh black; the forehead is covered with black down, or hairs; the hind part of the head bare and red, with a few ſcattered hairs; on the nape is a bare place aſh-coloured; the ſides of the head and hind part of the neck white; between the bill and eyes, beneath them, and the fore part of the neck, blackiſh aſh-colour; the lower part of the neck, and the reſt of the body, fine aſh-colour, darkeſt on the tail coverts; the quill-feathers and greater coverts black; baſtard wing the ſame; from the pinion of each wing, neareſt the body, ſprings a large tuft of unwebbed feathers, curled at the ends, which can be erected at will, but moſt frequently hang over and cover the tail.

There ſeems no doubt that this bird was by no means uncommon formerly in our fens; but only one or two inſtances are recorded by our later writers of its having been killed in this kingdom.

It is common in many parts of the European continent and Aſia, migrating with the ſeaſon. It breeds in the more northern parts. It is ſaid to fly remarkably high in its paſſage, even out of ſight; but that their paſſing is known by the noiſe they make, which is louder than any other bird: this is attributed to the formation of the windpipe, which is of a very ſingular ſtructure: it enters the keel of

the ſternum, or breaſt-bone, which is formed with a large cavity to receive it; from thence it returns, after being twice reflected. But we refer our readers to Dr. Latham's Eſſay on the *Tracheæ* of Birds, publiſhed in the fourth volume of the *Linnean Tranſactions*.

This bird is ſaid to make its neſt in the marſhes, and to lay two bluiſh eggs.

CRANE. Vide *Shag*.

CRANK-BIRD. Vide *Woodpecker leaſt-ſpotted*.

CREEPER. A genus of birds, the characters of which are,

Bill ſlender, incurvated.
Toes three before, one behind.
Claws hooked and long.
Tail of twelve ſtiff ſharp-pointed feathers.

CREEPER-COMMON.

Certhia familiaris. *Lin. Syſt.* i. p. 184. 1. *Gmel. Syſt.* i. p. 469. *Ind. Orn.* i. p. 280. 1.
Certhia. *Briſ.* iii. p. 603. 1.—*Ib.* 8vo. ii. p. 2. *Raii Syn.* p. 47. A. 5. *Will.* p. 100. t. 23.
Grimpereau. *Buf.* v. p. 481. t. 21. f. 1.
Common Creeper. *Br. Zool.* i. N° 92. t. 39. —*Ib.* fol. p. 82. t. K. *Arct. Zool.* ii. N° 174. *Albin.* iii. t. 25. *Lath. Syn.* ii. N° 701.—*Ib. Sup.* p. 126. *Lewin, Br. Birds*, ii. t. 53. *Walc. Syn.* i. t. 54. *Pult. Cat. Dorſet.* p. 5.

PROVINCIAL.

Tree-creeper. Tree-climber.

This is the only ſpecies of the genus in England; its weight about two drams; length five inches; bill half an inch long, ſlender and curved; irides hazel.

The upper part of the head and neck are prettily marked with ſtreaks of black and yellow brown; above each eye is a ſtroke of white; back, rump, and ſcapulars, inclining to tawny; quills duſky, moſtly tipped and edged with white, or very light brown; the coverts are varied with duſky brown and yellowiſh white, the laſt of which forms a ſort of bar acroſs the wing; the breaſt and belly are of a ſilvery white; the tail conſiſts of twelve ſharp-pointed ſtiff feathers, of a tawny brown.

Some authors have deſcribed this bird as poſſeſſing only ten feathers in the tail, which is a miſtake.

This bird is perpetually climbing up the body and limbs of trees in ſearch of inſects, its only food.

It makes a neſt in ſome hole, or behind the bark of ſome decayed tree, compoſed of dry graſs and the inner bark of wood, looſely put together, and lined with ſmall feathers. The eggs are from ſix to eight in number (not twenty, as ſome aſſert), weight about eighteen grains; theſe are white, minutely ſpeckled with bright ruſt-colour. During the time of incubation the female is fed by the other ſex, whenever ſhe quits her neſt in ſearch of food. The note of the Creeper is monotonous and weak, ſeveral times repeated in a deliberate manner; but

but is rarely heard in winter. At this ſeaſon it is conſtantly active in ſearch of food, which is chiefly the *larvæ* of inſects, found under moſs, and in the crevices of the bark; which is procured in ſufficient abundance to ſubſiſt it during that ſeaſon.

CROCKER. Vide *Gull black-headed.*

CROOKEDBILL. Vide *Avoſet.*

CROSSBILL.

Loxia curviroſtra. *Lin. Syſt.* i. p. 299. 1. *Gmel. Syſt.* ii. p. 843. *Ind. Orn.* i. p. 370. 1.

Loxia. *Briſ.* iii. p. 329. t. 17. f. 3.—*Ib.* 8vo. p. 401. *Raii Syn.* p. 86. A. *Will.* p. 181. t. 44.

Le Bec croiſé. *Buf.* iii. p. 449. t. 27. f. 2.

Croſsbill, *or* Sheld-apple. *Br. Zool.* i. N° 115. t. 49.—*Ib.* fol. 106. t. U. f. 2. *Arct. Zool.* ii. N° 208. *Will. Angl.* p. 248. t. 44. *Edw.* t. 303. *Albin.* i. t. 61. *Lath. Syn.* iii. p. 106. 1. *Lewin, Br. Birds,* ii. t. 66. *Walc. Syn.* ii. t. 205. *Pult. Cat. Dorſet.* p. 11. *Don. Br. Birds,* ii. t. 39.

This ſpecies of Groſbeak weighs about an ounce and a half; length near ſix inches and a half; bill ſtrong, both mandibles convex, and croſſing each other at the points, which are hooked; colour brown; eyes ſmall; irides duſky.

The plumage of the male varies from a beautiful red to orange-colour on the head, neck, breaſt, back,

back, and rump; the wing coverts rufous-brown; quills and tail duſky; vent almoſt white; under-tail coverts ſpotted duſky; tail forked; legs ſhort; claws ſtrong.

The females alſo vary ſomewhat in colour. In general they are of a dull olive-green on thoſe parts where the male is red; but the feathers on the back are mottled with duſky; the wings and tail ſimilar to that of the male, but not ſo dark. We have obſerved the croſſing of the mandibles are not conſtantly on the ſame ſide.

The Croſsbill is not known to breed with us, but is more or leſs found, amongſt our fir plantations, from June to the latter end of the year, feeding on the ſeed, by dexterouſly dividing the ſcales of the cone, for which purpoſe the bill is admirably adapted. It is ſometimes found in our orchards in autumn, and will readily divide an apple to get at the kernels. In the year 1791 we were informed, by a birdcatcher at Bath, that he had taken an hundred pair in the months of June and July: the greater part were males, which were generally ſold for five ſhillings each. Many are taken with a call-bird and birdlime; others are caught by a horſe-hair nooſe fixed to a long fiſhing-rod. So intent are theſe birds when picking out the ſeeds of the cone, that they will ſuffer themſelves to be taken by the nooſe being put over the head. They are diſcovered by the twittering noiſe they make while feeding.

This

This bird is ſaid to be a conſtant inhabitant of the northern parts of Europe; breeds early in the month of March, on the tops of the pine-trees.

In confinement it ſeems to have much the actions of a Parrot, climbing up the wires of a cage by the claws and beak.

Mr. Pennant mentions a variety he received out of Shropſhire, ſuperior in ſize, with the bill remarkably thick and ſhort, and more incurvated than in the common kind; the ends more blunt.

CROW. A genus of birds, the characters of which are,

Bill ſtrait, ſtrong.

Noſtrils covered with reflected briſtles.

Tongue bifid at the end.

Toes, the outer connected to the middle one as far as the firſt joint.

CROW-CARRION.

Corvus corone. *Lin. Syſt.* i. p. 155. 3. *Gmel. Syſt.* i. p. 365. *Ind. Orn.* i. p. 152. 4.

Cornix. *Raii Syn.* p. 39. A. 2. *Will.* p. 83. *Briſ.* ii. p. 12. 2.—*Ib.* 8vo. p. 157.

La Corneille. *Buf.* iii. p. 45. t. 3.

Carrion Crow. *Br. Zool.* i. N° 75. t. 34.—*Ib.* fol. p. 75. *Arct. Zool.* ii. N° 135. *Albin.* ii. t. 21. *Will. Angl.* p. 122. t. 18. *Lath. Syn.* i. p. 370. 3. *Lewin, Br. Birds,* i. t. 34. *Walc. Syn.* i. t. 33. *Pult. Cat. Dorſet.* p. 4.

PROVINCIAL.

Gor-crow.

This

This ſpecies weighs about nineteen ounces; length eighteen inches; bill black; irides duſky.

The whole plumage black, gloſſed above with a purpliſh blue; legs black.

Diſtinguiſhed from the Rook by the bill, which is rather more convex towards the end; and the nares, or reflected briſtles, at the baſe being always perfect: but theſe are only obvious in adults. In young birds the note is the only criterion of diſtinction, which in this is much more hoarſe than that of the Rook.

The Crow feeds on fleſh, inſects, and grain; but is particularly fond of carrion. It frequently attacks the eyes of dying animals, and deſtroys weakly lambs; it will alſo purſue birds on wing when preſſed by hunger. We once ſaw this bird in purſuit of a pigeon, at which it made ſeveral pounces, like a Hawk; but the Pigeon eſcaped by flying in at the door of a houſe. We have alſo ſeen it ſtrike a pigeon dead from the top of a barn. It is alſo a great deſtroyer of young game and poultry.

It is a bold bird, always at war with the leſſer ſpecies of Hawks; nor does the Kite, the Buzzard, or the Raven, approach its neſt without being attacked, and driven away. Even the Perigrine Falcon is inſulted at that ſeaſon, who frequently at one pounce brings it to the ground.

Theſe birds keep in pairs all the year; ſeldom congregate but to regale on ſome carcaſe, or in winter

winter to rooſt; will frequently hide their food till hunger is more preſſing.

The neſt is generally placed in the forked branch of a tree; is compoſed of ſticks plaſtered with earth, on which are laid various ſoft materials, ſuch as wool and hair. The eggs are four or five in number, of a greeniſh colour, ſpotted with duſky and aſh-colour; their weight about five drams.

CROW-BUNTING. Vide *Crow-hooded.*

CROW-DUN. Vide *Crow-hooded.*

CROW-GOR. Vide *Crow-carrion.*

CROW-HOODED.

Corvus Cornix. *Lin. Syſt.* i. p. 156. 5. *Gmel. Syſt.* i. p. 366. 5. *Ind. Orn.* i. p. 153. 7.

Cornix cinerea. *Raii Syn.* p. 39. A. 4. *Will.* p. 88. t. 18 & 77. *Briſ.* ii. p. 19. 4.

La Corneille mantelée. *Buf.* iii. p. 61. t 4.

Royſton Crow. *Will. Angl.* p. 124. t. 18. 77. *Albin.* ii. t. 23.

Hooded Crow. *Br. Zool.* i. N° 77.—*Ib.* fol. p. 76. t. D. 1. *Arct. Zool.* ii. p. 251. D. *Lath. Syn.* i. p. 374. 5.—*Ib. Sup.* p. 77. *Lewin, Br. Birds,* i. t. 36. *Walc. Syn.* i. t. 35. *Don. Br. Birds,* v. t. 117. *Pult. Cat. Dorſet.* p. 5.

PROVINCIAL.

Dun-Crow. Scare-Crow. Bunting-Crow.

This ſpecies weighs about twenty-two ounces; length

ſength twenty-one inches; bill black, in ſize and ſhape like the Carrion Crow; irides duſky.

The head, underſide of the neck, the point of the breaſt, wings, and tail, are of a gloſſy bluiſh-black; belly, back, ſcapulars, and upper ſide of the neck, are of a light aſh-colour; legs and claws black.

The Hooded Crow viſits the ſouthern parts of this kingdom in October, and retires northward to breed in the beginning of April; is found in ſome parts of Scotland during the ſummer; and we have received its eggs from Ireland, in ſome parts of which it continues the whole year.

The neſt and eggs are ſimilar to thoſe of the Common Crow. In England theſe birds are principally found near the ſea-coaſt, where the various animal matter thrown up by the tide afford a conſtant ſupply of food. It is alſo ſeen in conſiderable abundance upon ſome of our downs, or in the open champaign parts, where it feeds on grain, worms, and carrion. Ten or a dozen are not unuſually ſeen perched together in the neighbourhood of a dead ſheep, patiently watching till the ſhepherd's dog has filled his paunch, and, retiring to his maſter's coat, is ſlunk in deep repoſe. Like the laſt ſpecies, it attacks the eyes of ſickly animals. We have ſeen it make repeated attacks upon the eyes of weakly ſheep, and generally ſucceed if the animal is incapable of riſing. Mr. Pennant informs us this is the only ſpecies of genuine Crow in all the Hebrides,

brides, Orknies, and Shetlands, the Carrion and the Rook being unknown there.

It is not uncommon in Germany; is alſo found in Ruſſia and Siberia.

CROW-MIRE. Vide *Gull black-headed.*

CROW-RED-LEGGED.

Corvus graculus. *Lin. Syſt.* i. p. 158. 18. *Gmel. Syſt.* i. p. 377. *Ind. Orn.* i. p. 165. 41.

Coracias, ſeu Pyrrhocorax. *Raii Syn.* p. 40. A. 6. *Will.* p. 86. t. 19. *Brif.* ii. p. 3. t. 1. f. 1.

Le Coracias. *Buf.* iii. p. 1. t. 1.

Corniſh Chough. *Albin.* ii. t. 24. *Borlas, Cornw.* p. 249. t. 24. *Will. Angl.* p. 126. t. 19. *Hayes, Br. Birds,* t. 6.

Red-legged Crow. *Br. Zool.* i. N° 80. t. 35.— *Ib.* fol. p. 83. t. L. *Lewin, Br. Birds,* i. t. 41. *Lath. Syn.* i. p. 401. 39. *Walc. Syn.* i. t. 40. *Pult. Cat. Dorſet.* p. 5.

PROVINCIAL.

Corniſh-daw. Cornwall-kae. Killigrew.

This ſpecies weighs about fourteen ounces; length near ſeventeen inches.

The bill is longer and more ſlender than in any of the genus, a little curved, of a deep orange-red, much reſembling red coral, and is remarkably brittle; irides hazel.

The

The plumage is wholly black, gloſſed with purple; legs and feet red; claws black, ſtrong, and much hooked

The female differs in not being ſo large, and in the bill being ſhorter; the plumage in both ſexes is alike.

This bird with us ſeems to be chiefly confined to Devonſhire, Cornwall, and Wales, where it is found on moſt of the bold rocky ſhores. It has been ſeen on the cliffs of Dover, ſuppoſed to have eſcaped from confinement, and ſtocked thoſe rocks. But we believe the breed in thoſe parts is again loſt.

Mr. Pennant obſerves that it is found in ſome parts of Scotland, and in the Hebrides. It is ſeldom ſeen at any great diſtance from the ſea-coaſt, where it breeds in the rocks and caverns; and not unfrequently in ruined towers. A pair of theſe birds had for many years bred in the ruins of Crow Caſtle, in the vale of Llangollen in Denbighſhire; by accident one of them was killed, and the other continued to haunt the ſame place for two or three years, without finding another mate.

The neſt is compoſed of ſticks, and lined with a great quantity of wool and hair. The eggs are generally five in number, of a dull white, ſprinkled with light-brown and aſh-coloured ſpots, moſt at the larger end; their weight about three drams and a half.

The note of this bird is ſomewhat like that of the

the common Jackdaw, but more ſhrill. Its food is grain and inſects, but in confinement will greedily feed on fleſh; becomes extremely tame and docile, active, and crafty; will hide part of its food, and not unfrequently things of value. We are told houſes have been ſet on fire by its carrying away lighted ſticks in its bill. Some are ſaid to acquire black legs in autumn. This, however, does not appear to be the caſe in England. With us they remain all the year, and rarely ſhift their quarters; in other parts they ſhould ſeem migratory, as it has been obſerved in Egypt to attend the inundation of the Nile in the months of September and October; is found alſo upon the Alps; and is ſaid to be met with in Perſia.

CROW-ROYSTON. Vide *Crow-hooded.*

CROW-SCARE. Vide *Crow-hooded*, and *Tern-black.*

CROW-SEA. Vide *Corvorant*, *Gull-ſkua*, and *Gull-black-headed.*

CUCKOW, or CUCKOO. A genus of birds, the characters of which are,

Bill ſomewhat arched.

Tongue ſhort.

Tail compoſed of ten feathers.

Toes, two backward, two forward.

CUCKOW-COMMON.

Cuculus canorus. *Lin. Syſt.* i. p. 168. 1. *Gmel. Syſt.* i. p. 409. *Raii Syn.* p. 23. *Will.* p. 6.

t. 10. 27. *Brif.* iv. p. 105. 1. *Ind. Orn.* i. p. 207. 1.

Coucou. *Buf.* vi. p. 305.

Common Cuckow. *Br. Zool.* i. N° 82. t. 36. —*Ib.* fol. 80. t G. G. 1. *Arct. Zool.* ii. t. 266. A. *Will. Angl.* p. 97. t. 10. 77. *Albin.* i. t. 8. *Hayes, Br. Birds,* t. 17. 18. *Lewin, Br. Birds,* ii. t. 42. *Lath. Syn.* ii. p. 509. 1. —*Ib. Sup.* p. 98. *Walc. Syn.* i. t. 43. *Don. Br. Birds,* ii. t. 41. *Pult. Cat. Dorſet.* p. 5.

The Common Cuckow is the only Britiſh ſpecies; it weighs about four ounces and a half; length fourteen inches.

The bill is black, yellowiſh at the baſe of the under mandible; inſide of the mouth red; irides yellow; the head and whole upper part of the bird dark aſh-colour; throat, under ſide of the neck, and upper breaſt, of a pale aſh-colour, the latter in ſome inclining to rufous-brown; lower breaſt and belly white, marked with tranſverſe undulated black lines; the quill-feathers are duſky, the inner webs barred with oval white ſpots; the tail conſiſts of ten feathers of unequal length, the two middle ones black, daſhed with aſh-colour, and tipped with white; the reſt are black, marked with white ſpots on each ſide the ſhaft; in ſome the lateral feathers have white ſpots only on their interior webs, but are all tipped with white.

The female is rather leſs, and in general differs from

from the other ſex in the neck and breaſt being of a tawnyiſh brown, barred with duſky, and the coverts of the wings marked with light ferruginous ſpots; the markings on the tail and quill-feathers much like the male, only the edges of the ſpots are inclining to reddiſh brown; the legs of both ſexes ſhort and yellow.

The outer-tail feather and the firſt quill-feather are remarkably ſhort.

We believe this bird does not entirely throw off its neſtling feathers till the ſecond year's moulting; for in three ſpecimens before us, killed the ſame ſeaſon (two males and a female), the thirteenth and three ſucceeding quill-feathers, and the three greater coverts impending them, are barred with brown and ferruginous.

This bird comes to us early in the ſpring, and almoſt invariably leaves us by the firſt of July; the females may ſometimes remain ſomewhat later, not having depoſited all their eggs. We have killed this ſex as late as the twenty-ſixth of June, from which we took a matured egg, which weighed forty-four grains, the colour white, ſprinkled with two ſhades of aſh-coloured ſpots, moſtly at the larger end.

The male generally comes to us earlier, and leaves us again before the other ſex.

The food of the Cuckow is chiefly inſects, particularly caterpillars, or the *larvæ* of the *lepidopterous* inſects of all kinds, and not only the ſmooth

ſort, as have been imagined, for we have ſeen the ſtomach of this bird more than once lined with the hairs of the rougher ſort.

It is well known this ſingular bird makes no neſt, but depoſits its eggs in the neſts of ſome ſmall bird. For the illuſtration of the natural hiſtory of this ſpecies we are much indebted to Mr. Genner, who, in a letter to Mr. John Hunter, publiſhed in the Tranſactions of the Royal Society, has given much light on the ſubject. We muſt therefore refer our curious reader to that work; and alſo to the Introduction to this, where the curious phenomena belonging to this bird is more copiouſly treated of.

It has been aſſerted that the Cuckow does not leave us till the latter end of September, or beginning of October; but ſuch as are ſeen ſo late are the young of that year, or an accidental old one that may have been wounded. We have, indeed, been told it has been heard in winter; and we have known the young killed ſo late as November; but doubtleſs theſe were prevented remigrating by ſome defect. We have alſo been told its well-known cry has been heard in the ſummer at midnight, when the moon has ſhone, the note of the male; the female makes only a chattering noiſe.

The young Cuckow is not the leaſt like the old birds in plumage; it is brown, mixed with ferruginous and black.

The foſter-parent of this bird feeds it long after

it

it has left the neſt, and in order to receive its nouriſhment it uſually lays down on one ſide, in order that the ſmall bird may reach its mouth, and frequently extends one wing for the bird to pitch on to perform that office.

We have known theſe birds ſurvive the winter in confinement, fed on raw meat.

CUDDY. Vide *Gallinule-common.*

CURLEW. A genus of birds, the characters of which are,

Bill long, incurvated.
Face covered with feathers.
Noſtrils linear.
Tongue ſhort, ſharp-pointed.
Toes connected, as far as the firſt joint, by a membrane.

CURLEW-COMMON.

Scolopax Arquata. *Lin. Syſt.* i. p. 242. *Gmel. Syſt.* ii. p. 655.

Numenius Arquata. *Ind. Orn.* ii. p. 710. 1.

Numenius. *Raii Syn.* p. 103. A. 1. *Will.* p. 216. t. 54. *Briſ.* v. p. 311. 1.—*Ib.* 8vo. ii. p. 289.

Le Courlis. *Buf.* viii. p. 19.

Common Curlew. *Br. Zool.* ii. N° 176. t. 63.—*Ib.* fol. 118. *Arct. Zool.* p. 462. A. *Will. Angl.* p. 294. t. 54. *Albin.* i. t. 79. *Lath. Syn.* v. p. 119. 1.—*Ib. Sup.* p. 242. *Pult. Cat,*

Cat. Dorſet. p. 14. *Walc. Syn.* ii. t. 133. *Lewin, Br. Birds,* iv. t. 153.

This ſpecies is ſubject to vary conſiderably in ſize, weighing from twenty to upwards of thirty ounces; the length of the largeſt about twenty-five inches.

The bill is from ſix to ſeven inches long, duſky black; irides hazel.

The head, neck, ſcapulars, and coverts of the wings, pale brown, each feather black in the middle, and lighteſt at the edges; breaſt and belly white, marked with oblong black ſpots; lower part of the back white, with a few duſky ſpots; the tail and its upper coverts barred with yellowiſh white and black; quills black, ſpotted on the inner webs with white; the legs are long, of a bluiſh grey colour.

Male and female much alike.

The Curlew is common on moſt parts of our coaſt in winter, where it feeds on ſmall crabs and other marine inſects, and worms. At this ſeaſon it is gregareous.

In the ſpring theſe birds retire inland, and moſt commonly to the more northern parts of this kingdom, to breed. For this purpoſe the moſt retired ſituation is reſorted to, either on the mountains amongſt the heath, or in the extenſive, unfrequented marſhes. We have taken the young on the mountains in Northumberland, and in the low, ſwampy grounds

grounds in the iſle of Mull in Scotland. It makes no neſt, but depoſits its eggs amongſt the heath, ruſhes, or long graſs, generally four in number, of a pale olive colour, marked with browniſh ſpots.

The young make uſe of their legs as ſoon as they are hatched, but cannot fly for a conſiderable time.

CURLEW-JACK, or HALF CURLEW. Vide *Whimbrel.*

CURLEW-KNOT. Vide *Whimbrel.*

CURLEW-LAND. Vide *Buſtard-thick-kneed.*

CURLEW-PIGMY.

Scolopax pigmea. *Gmel. Syſt.* ii. p. 655.

Numenius pigmeus. *Ind. Orn.* ii. p. 713. 11. —*Ib. Sup.* p. 291. nota o. *Gen. of Birds*, p. 64. t. 11. *Boys, Sandwich*, t. p. 850. *Walc. Syn.* ii. t. 135. *Lewin, Br. Birds*, iv. t. 155.

This ſpecies is deſcribed to be the ſize of a Lark; weight near two ounces; length eight inches and a half; the bill one inch and a half long, bent and black; the head, back, and coverts of the wings, mixed with brown, ferruginous, and white; primores duſky, edged with white; breaſt, belly, and rump, white; tail duſky; the exterior feathers edged with white; legs black.

This is a very rare bird; only two ſeems to appear on record; one was killed in Holland, the

other near Sandwich in Kent. In this ſpecimen the edges of all the feathers were of a very pale oker inſtead of white, as Dr. Latham informs us, from whoſe works we have borrowed this deſcription.

CURLEW-STONE. Vide *Buſtard-thick-kneed*, and *Whimbrel.*

CUSHAT. Vide *Dove-ring.*

D.

DAB-CHICK. Vide *Grebe-little.*

DAKER-HEN. Vide *Gallinule-crake.*

DAW. Vide *Jackdaw.*

DIDAPPER. Vide *Grebe-little.*

DIVER. A genus of birds, the characters of which are,

Bill ſtrong, ſtrait, pointed; upper mandible the longeſt, edges of each bending inwards.

Noſtrils linear, the upper part divided by a ſmall cutaneous appendage.

Tongue long and pointed, ſerrated on each ſide near the baſe.

Legs thin and flat.

Toes four in number, the exterior longeſt; the back toe ſmall, joined to the inner one by a ſmall membrane.

Tail ſhort; conſiſts of twenty feathers, or more.

DIVER-BLACK. Vide *Scoter*, and *Duck-velvet*.

DIVER-BLACK-THROATED.

Colymbus arcticus. *Lin. Syſt.* i. p. 221. 4. *Gmel. Syſt.* ii. p. 587. *Ind. Orn.* ii. p. 800. 4. *Raii Syn.* p. 125. 7. *Will.* p. 259. t. 62.—*Ib. Angl.* p. 343. t. 62.

Mergus gutture nigro. *Briſ.* vi. p. 115. 4.—*Ib.* 8vo. ii. p. 391.

Lumme. *Buf.* viii. p. 262.

Black-throated Diver. *Br. Zool.* ii. N° 241. t. 85. f. 2. *Arct. Zool.* ii. N° 444. *Edw.* t. 146. *Lath. Syn.* vi. p. 343. 4. *Lewin, Br. Birds*, vi. t. 229.

PROVINCIAL.

Northern Doucker. Speckled Zoon.

The length of this ſpecies is two feet.

Bill near two inches long, ſlender, black.

The fore part of the head and throat black; hind part of the head and neck aſh-colour; ſides of the neck white, ſpotted black; on the fore part of the neck a large patch of black, changeable to purple and green in different lights; the back and upper parts black; ſcapulars marked with ſquare ſpots of white; wing coverts with round ſpots; breaſt and belly white; quills duſky; tail ſhort and black; legs black, with a reddiſh caſt on the inſide.

This bird is rarely found in England, but is not uncommon in the northern parts of Europe; ſaid

to

to inhabit the lakes of Siberia, eſpecially thoſe of the arctic regions, as well as North America, particularly Hudſon's Bay.

In ſome countries the ſkin is uſed for various ſorts of clothing, and other purpoſes, being warm and exceedingly tough, which is common to all the genus.

DIVER-DUN. Vide *Dun-diver.*

DIVER-DUN-LESSER. Vide *Merganſer-red-breaſted.*

DIVER-GREATEST-SPECKLED. Vide *Diver-northern*, and *Diver-ſpeckled.*

DIVER IMBER.

Colymbus Immer. *Lin. Syſt.* i. p. 222. 6. *Gmel. Syſt.* ii. p. 588. *Ind. Orn.* ii. p. 800. 2.

Colymbus maximus Geſneri. *Raii Syn.* p. 126. 8. *Will.* p. 260. III.

Mergus major. *Briſ.* vi. p. 105. 1. t. 10. f. 1. —*Ib.* 8vo. ii. p. 389.

Le grand Plongeon. *Buf.* viii. p. 251.

Ember Gooſe. *Sibb. Scot.* 21.

Imber Diver. *Br. Zool.* ii. N° 238. t. 84. *Arct. Zool.* ii. N° 440. *Will. Angl.* p. 342. *Lath. Syn.* vi. p. 340. 2. *Pult. Cat. Dorſet.* p. 17. *Walc. Syn.* i. t. 99. *Don. Br. Birds*, iv. t. 99. *Lewin, Br. Birds*, vi. t. 227.

PROVINCIAL.

Immer. Greater Doucker. Cobble.

This

This ſpecies is leſs than the Northern Diver; it meaſures two feet in length.

The bill is four inches long, of a duſky brown-colour. The top of the head and back part of the neck brown; the forehead, ſides of the neck, and cheeks, ſpeckled with brown; the feathers on the back and wings are brown, with paler margins; the throat and upper part of the neck before are ſpotted black and white; the quills and tail brown, in ſome duſky black, the laſt edged with white; the whole under parts are white, with a little brown about the vent; legs duſky.

The female is leſs bright in colour above, and duſky white beneath; the neck ſcarcely ſpeckled at all.

This is rather a ſcarce ſpecies in the ſouth of England, but is ſometimes met with in ſevere winters. It is more common in Scotland, and in the ſea about the Orkneys; is found alſo in moſt parts of the north of Europe; ſaid to inhabit Switzerland, particularly the lake Conſtance, where it is known by the name of Fluder.

It makes a neſt on the water, placed amongſt the reeds and flags.

It feeds on fiſh, after which it dives with great celerity, and is ſometimes taken under water by a baited hook.

DIVER-LESSER-TOOTHED. Vide *Merganſer redbreaſted.*

DIVER-LOUGH. Vide *Smew.*

DIVER-

DIVER-NORTHERN.

Colymbus glacialis. *Lin. Syſt.* i. p. 221. 5. *Gmel. Syſt.* ii p. 588. *Ind. Orn.* ii. p. 799. 1.

Colymbus maximus caudatus. *Raii Syn.* p. 125. A. 4. *Will.* p. 259.

Mergus major nævius. *Briſ.* vi. p. 120. 6. t. 11. f. 2.

Mergus nævius. *Briſ.* vi. p. 118. 5.—*Ib.* 8vo. ii. p. 392—391.

L'Imbrim. *Buf.* viii. p. 258. t. 22.

Greateſt ſpeckled Diver, *or* Loon. *Will. Angl.* p. 341. *Albin.* iii. t. 93.

Northern Diver. *Br. Zool.* ii. N° 327. t. 84. —*Ib.* fol. 139. t. K. 2. *Arct. Zool.* ii. N°439. *Don. Br. Birds,* iii. t. 58. *Lath. Syn.* vi. p. 337. 1. *Pult. Cat. Dorſet.* p. 17. *Walc. Syn.* i. t. 90. *Lewin, Br. Birds,* vi. t. 226.

This ſpecies is the largeſt of the genus, ſometimes weighing as much as fifteen or ſixteen pounds; length near three feet and a half.

The bill is black, four inches and a half long; irides purpliſh; the head and neck deep velvety black; on the throat are ſeveral parallel white lines, formed of raiſed feathers; on each ſide the neck a large portion of the ſame, almoſt uniting behind and before; the ſides of the breaſt ſtreaked with black and white lines; the back, ſcapulars, and wing coverts, are black, marked with white ſpots in a moſt elegant manner; thoſe on the back and

rump

rump are ſmall and round, the others are large and of a ſquare form, diſpoſed in rows; the quills and tail are black; the breaſt and under parts of the body white, with a few black ſtreaks under the wings; legs black.

The female is not ſo large, and the white markings on the neck are leſs diſtinct. The variation which has been obſerved in the plumage of theſe birds has very juſtly been conſidered to be owing to its not arriving at perfection till the ſecond, or perhaps the third year.

The Northern Diver is rarely met with in the ſouthern parts of England. It ſeldom leaves the water; but inſtances are recorded of its having been taken alive on land. In the ſpring of the year 1797 one of theſe birds was taken near Penzance in Cornwall, at ſome diſtance from water. It appeared incapable of raiſing itſelf from the ground; it did not ſeem to have any defect, as it lived for ſix weeks in a pond, and was ſupplied with fiſh; but for want of a ſufficient quantity was ſtarved, as was apparent when we diſſected it for preſervation.

It is not uncommon in Iceland and Greenland, where it breeds in the freſh waters, and is ſaid to lay two large eggs of a pale brown colour in the month of June. Is plentiful in Norway, and ſome parts of Ruſſia. In the latter country, as well as in ſome others, the ſkin is dreſſed and uſed for various ſorts of clothing. It is tough, and well covered with ſoft down.

DIVER-RED-THROATED.

Colymbus ſeptentrionalis. *Lin. Syſt.* i. p. 220. 3. *Gmel. Syſt.* ii. p. 586. *Ind. Orn.* ii. p. 801. 5.

Mergus gutture rubro. *Briſ.* vi. p. 111. 3. t. 11. f. 1.—*Ib.* 8vo. ii. p. 390.

Le Plongeon à gorge rouge. *Buf.* viii. p. 264.

Red-throated Diver, *or* Loon. *Br. Zool.* ii. N° 240. t. 85.—*Ib.* fol. 140. *Arct. Zool.* ii. N° 443. *Edw.* t. 97. *Lath. Syn.* vi. p. 344. 5. *Pult. Cat. Dorſet.* p. 17. *Walc. Syn.* i. t. 100. *Lewin, Br. Birds,* vi. t. 230. *Don. Br. Birds,* iv. t. 78.

This ſpecies weighs about three pounds; length near two feet and a half.

Bill black, three inches long, and ſlender; irides hazel. The head and upper part of the neck before cinereous; the reſt of the neck, running up behind almoſt to the head, is marked with longitudinal duſky and white lines; on the throat is a patch of cheſnut red; the upper part of the body, wings, and tail, duſky; the feathers of the back and ſcapulars ſlightly margined with brown; the under parts of the body white; the ſides under the wings and thighs ſtreaked with duſky; vent brown; legs duſky on the outſide, lighter within.

The bird from which this deſcription is taken was killed at Haſtings in Suſſex in the winter of 1795, at which time a great many were ſeen on that coaſt.

This,

This, like others of the genus, ſeems ſubject to ſome variety, as we are informed ſome have the head and chin dotted with brown, and the cheſnut mark on the throat extends farther on the neck. Whether this variety is occaſioned by age or ſex is not aſcertained.

This bird is ſaid to breed in the northern parts of Scotland, but rarely viſits the ſouthern parts of England, except in very ſevere ſeaſons In the breeding ſeaſon it frequents the lakes, making a neſt amongſt the reeds and flags, and lays two eggs of an aſh-colour, marked with a few black ſpots. Is found in moſt of the northern parts, common in Greenland and Iceland; in the latter it is ſaid to make its neſt amongſt the graſs on the ſhores contiguous to the water, compoſed of moſs and graſs, and lined with down.

It is frequently taken in nets, by diving after the fiſh which are entangled. In the General Synopſis mention is made of no leſs than fourteen having been taken by this means out of a ſingle net at Hudſon's Bay at one tide.

DIVER-SPECKLED.

Colymbus ſtellatus. *Gmel. Syſt.* ii. p. 587. 17. *Ind. Orn.* ii. p. 800. 3.

Maximus caudatus. *Raii Syn.* p. 125. A. 4.? *Will.* p. 258. t. 61.

Colymbus caudatus ſtellatus. *Will.* t. 62.

Mergus minor. *Briſ.* vi. p. 108. 2. t. 10. f. 2. —*Ib.* 8vo. ii. p. 389.

Le petit Plongeon. *Buf.* viii. p. 254. t. 21.
Speckled Diver, or Loon. *Br. Zool.* ii. N° 139.
—*Ib. Sup.* 119. t. K. *Arct. Zool.* ii. N° 441.
Alcia. i. t. 12. *Latb. Syn.* vi. p. 341. 3.
Pult. Cat. Dorſet. p. 17. *Id. Syn.* i. t. 101.
Lacep. Br. Birds, vi. t. 223.

PROVINCIAL.

Sprat Loon. Greateſt Speckled Diver. Cobble.

The weight of this ſpecies is about four pounds; length twenty ſeven inches.

Bill three inches long, of a pale horn-colour, ridge of the upper mandible duſky; irides yellowiſh brown.

The head is duſky, ſpeckled with grey; the hind part of the neck plain duſky; on the ſides of the head under the eyes, the chin, and throat, white; the fore part of the neck ſpeckled with aſh-colour; back and whole upper parts duſky, marked with numerous ſmall oval white ſpots; quills and tail duſky; in ſome the latter is ſlightly tipped with white; the whole under parts of the body white, except on the ſides under the wings, which are ſtreaked with duſky; the thighs duſky brown; the vent and under-tail coverts mottled with grey; the tail conſiſts of twenty-two unequal feathers, the outer ones not being half ſo long as the middle ones; legs duſky brown, inſide greeniſh grey.

The female is not ſo large, and the ſpots on the back and ſcapulars not ſo large and diſtinct.

The

The Speckled Diver is the moſt common ſpecies found in this country. It is frequently ſeen in winter in our bays and inlets, and ſometimes in freſh-water rivers and lakes. It is obſerved to attend the ſprats in the river Thames, for which reaſon the fiſhermen called it Sprat Loon. This, like the reſt of the genus, retires northward to breed; is ſaid to be common about the Baltic and the White Sea. It lays two eggs in the graſs upon the borders of lakes, the ſize of thoſe of a Gooſe, duſky, with a few black ſpots.

DIVING-PIGEON. Vide *Guillemot-black.*

DOB-CHICK. Vide *Grebes-duſky, eared, and little.*

DOB-CHICK-BLACK and WHITE. Vide *Grebe-duſky.*

DOB-CHICK-GREATER. Vide *Grebe-tippet.*

DOTTREL.

Charadrius Morinellus. *Lin. Syſt.* i. p. 254. 5. *Gmel. Syſt.* ii. p. 686. *Ind. Orn.* ii. p. 746. 17. *Raii Syn.* p. 111. A. 4. *Will.* p. 230. t. 55. 57. *Brif.* v. p. 54. 5. t. 4. f. 2. and p. 58. 6.—*Ib.* 8vo. ii. p. 225. and p. 126.

Petit Pluvier, ou le Guignard. *Buf.* viii. p. 87.

Dottrel. *Br. Zool.* ii. N° 210. t. 73.—*Ib.* fol. 129. t. D. *Arct. Zool.* ii. p. 487. A. *Will. Angl.* p. 309. *Albin.* ii. t. 62. and t. 63. *Lath. Syn.* v. p. 208. 14. *Pult. Cat. Dorſet.* p. 16. *Walc. Syn.* ii. t. 162. *Lewin, Br. Birds,* v. t. 186. *Don. Br. Birds,* ii. t. 42.

This ſpecies of Plover weighs about four ounces, ſometimes five; length near ten inches. The bill is an inch long, duſky; irides hazel; the crown of the head black; forehead duſky and grey, mixed; a broad ſtroke of white from above the eye paſſes to the hind-head; cheeks and throat white; the neck of a cinereous olive; back and coverts of the wings olive-brown, each feather marginated with pale ferruginous; the breaſt is of a pale, dull orange, on the upper part of which is a tranſverſe line of white, bordered above with a narrow one of black; belly black; vent and thighs rufous white; the quills duſky-brown; the ſhaft and outer web of the firſt feather white; the tail conſiſts of twelve feathers of an olive-brown, barred near the end with black, tipped with white; legs duſky.

The female has the crown of the head brown, mottled with white, and the white line over the eye leſs conſpicuous; the belly is mixed black and white; the white line on the breaſt is wanting; and the colours in general more dull.

Young birds have the whole under parts of the body of a very pale ferruginous brown.

The Dottrel appears to make this country a reſting-place in its migratory flights to and from its breeding-place. It is ſeen on ſome of our downs, heaths, and moors, from April to the beginning of June, and returns again in September, and remains till November.

On

On the Wiltſhire downs it reſorts to the new-ſown corn, or fallow ground, for the ſake of worms, its principal food. They fly in families of five or ſix in the autumn, which we have obſerved to be the two old birds and their young; but ſometimes a dozen or more flock together. It is a ſtupid bird, and eaſily ſhot; when diſturbed will frequently extend one wing, and do not fly to a great diſtance. It doubtleſs goes northward to breed; but we do not find any one who mentions the neſt or eggs. It is probable ſome may breed on the mountains in Scotland. We once ſaw them in pairs in that country in the ſummer ſufficiently late to form ſuch an opinion; and we are not ſingular in this conjecture.

It is ſaid to be not unfrequent on the Lapland alps, viſits Sweden in May, and breeds in the northern parts of Ruſſia and Siberia.

DOUCKER-GREATER. Vide *Diver-Imber.*

DOUCKER-HORNED. Vide *Grebe-creſted.*

DOUCKER-NORTHERN. Vide *Diver-black-throated.*

DOUCKER-SMALL. Vide *Grebe-little.*

DOVE. A genus of birds, the characters of which are,

Bill ſlender, weak, and ſtrait at the baſe.
Noſtrils lodged in a ſoft protuberance.
Tongue entire.
Legs ſhort.
Toes divided to their origin.

bruſh-wood, and in hedges, or large hawthorn buſhes; but more frequently in the fork of a tree, or againſt the body, when ſurrounded with ivy, and particularly in fir-trees. Their food is grain and ſeeds of all kinds. In the autumn they devour acorns and beech-nuts greedily, ſwallowing them whole. In defect of theſe and grain it eats turnip-greens and young clover, or green corn and ivy-berries.

We have been at conſiderable pains to endeavour to domeſticate this bird; and though we have tamed them within doors ſo as to be exceedingly troubleſome, yet we never could produce a breed, either by themſelves or with the tame Pigeon. Two were bred up together with a male Pigeon, and were ſo tame as to eat out of the hand; but as they ſhewed no ſigns of prolificacy in the ſpring, were ſuffered to take their liberty in the month of June, by opening the window of the room in which they were confined, thinking the Pigeon might induce them to return to their uſual place of abode, either for food or to rooſt; but they inſtantly took to their natural habits, and we ſaw no more of them, although the pigeon continued to return. We ſhall here mention a curious aſſemblage we once bred up, and lived together in perfect amity: A common Pigeon, Ring Dove, White Owl, and Sparrow Hawk; of which the Ring Dove was maſter.

DOVE-ROCK.

Columba Œnas. *Lin. Syſt.* i. p. 279. 1. β. *Gmel. Syſt.* ii. p. 769. *Raii Syn.* p. 62. A. 10. *Will.* p. 136. t. 35. *Ind. Orn.* ii. p. 589. 1. *Briſ.* i. p. 86. 5.

Stock Pigeon, *or* Stock Dove. *Br. Zool.* N° 100. t. 45. — *Ib. App.* p. 605.. *Arct. Zool.* ii. p. 329. A. *Albin.* ii. p. 46. — *Ib.* iii. t. 44. *Latb. Syn.* iv. p. 604. 1. — *Ib. Sup.* p. 197. *Will. Angl.* p. 185. *Pult. Cat. Dorſet.* p. 7. *Walc. Syn.* ii. t. 86. *Lewin, Br. Birds,* iv. t. 128. *Don. Br. Birds,* v. t. 107. *White, Selb.* p. 98. 110.

PROVINCIAL.

Rockier.

Ornithologiſts ſeem to differ in opinion concerning the Rock and Stock Pigeon; though it appears moſt conceive them to be diſtinct ſpecies. In thoſe deſcribed under ſuch names there ſeems to be ſo much ſimilitude, except what may be expected from a ſpecies half reclaimed, and frequently returning to their natural wild habits again, that we cannot but conſider them as one and the ſame ſpecies.

The Rock Dove is conſidered to be the origin of our tame Pigeons, as it is ſaid to poſſeſs the white on the lower part of the back, in which part the Stock Dove is deſcribed to be aſh-coloured, and that this laſt is rather larger. But theſe va-

riations we have obſerved in Pigeons killed in their native haunts amongſt the rocks on our coaſts; and our dove-cote Pigeons frequently have no white on the back. It is therefore probable many of our common ſpecies, after having been bred in a pigeon-houſe contiguous to ſuch rocky ſituations, return to their natural habits, and there produce ſome variation in colour.

The bird now before us we killed on the cliffs in Cau'dy Iſland, in South Wales.

It weighed eleven ounces; length thirteen inches and a half; breadth twenty-two.

The bill is brown, inclining to purpliſh red, point duſky; irides light yellow.

The head dark bluiſh aſh-colour; neck and breaſt gloſſed with green and copper, as viewed in different lights, moſt conſpicuous on the ſides and back of the neck; the upper part of the back and wing coverts pale aſh-colour; acroſs the middle of the greater coverts is a broad band of black, and another of the ſame on the ends of the ſecondary quills, running into each other on thoſe feathers neareſt the body; the greater quills are duſky, daſhed with aſh-colour, the outer ones darkeſt, and all of them moſt ſo towards the tips, ſlightly edged on their exterior webs with white; the lower part of the back white; the rump and tail dark bluiſh aſh-colour, the ends of the latter black; the two exterior feathers whitiſh on the outer webs towards the baſe; the ſides under the wings, and under

wing

wing coverts, white; the belly bluiſh aſh-colour; legs red.

In another ſpecimen, killed in the ſame place, the upper part of the back, and the whole of the wing coverts, are mottled with black and light aſh-colour, without any regular bands acroſs the wings. In every other reſpect it reſembled the firſt; but this was rather larger, weighing twelve ounces; length fourteen inches. It was a female, and was ſhot as ſhe flew from her neſt. This muſt, however, be conſidered as a variety; for in the more remote parts, where the breed is pure, the bars on the wings ſeem to be a characteriſtic mark, and common to both ſexes.

Theſe birds have ſometimes appeared in prodigious flocks in winter, frequenting our beech woods for the ſake of the maſt or ſeed of that tree. Theſe flights, however, are leſs numerous and leſs frequent of late years. Sometimes they are ſeen in company with our common Pigeons at the barn-doors in ſevere winters; and are ſaid to be known by their inferior ſize and darker colour.

This bird generally breeds in caverns on our coaſts, and in ruined edifices; and we are told ſometimes nidificate in the holes of decayed trees. Perhaps theſe laſt have been called Stock Doves, ſuppoſing them a diſtinct ſpecies. Mr. White informs us, in his Natural Hiſtory of Selborne, that Stock Doves formerly abounded in the beech woods near that place; and that amongſt them were

were ſeen little parties of ſmall blue Doves, which were there called Rockiers. However this may be, we are inclined to believe it is the ſame ſpecies, with ſome little variety.

All the beautiful varieties of Pigeons under the denomination of Carrier, Pouter, Nun, Owl, &c. are ſuppoſed to be derived from this ſpecies by domeſtication.

DOVE-STOCK. This appears to be no other than the preceding ſpecies. Vide *Dove-rock.*

DOVE-TURTLE.

Columba Turtur. *Lin. Syſt.* i. p. 284. 32. *Gmel. Syſt.* ii. p. 786. *Raii Syn.* p. 61. A. 2. *Will.* p. 134. t. 35. *Briſ.* i. p. 92. 7.—*Ib.* 8vo. i. p. 21. *Ind. Orn.* ii. p. 605. 47.

Turtur auritus. *Raii Syn.* p. 184. 26.

Tourterelle. *Buf.* ii. p. 545. t. 25.

Tourterelle de Portugal. *Buf.* ii. p. 556.

Turtur luſitanicus. *Briſ.* i. p. 98. 9.—*Ib.* 8vo. i. p. 23.

Common Turtle. *Br. Zool.* N° 103. t. 45.—*Ib.* fol. 88. t. 88. *Albin.* ii. t. 47—t. 48. *Hayes, Br. Birds,* t. 14. *Will. Angl.* p. 183. t. 35. *Lath. Syn.* iv. p. 644. 40. var. A. B. C. D.—*Ib. Sup.* p. 199. *Pult. Cat. Dorſet.* p. 7. *Walc. Syn.* ii. t. 188. *Lewin, Br. Birds,* iv. t. 130.

This elegant little ſpecies weighs rather more than ſix ounces; length twelve inches. The bill

is brown; irides reddiſh yellow; a bare ſpace beneath and behind the eyes of a purpliſh red.

The top of the head and upper part of the neck behind cinereous; on each ſide of the neck is a patch of black feathers, tipped with white; the back is brown, daſhed with aſh-colour, lighteſt on the margin of each feather; ſcapulars and wing coverts black, deeply margined with ferruginous brown; quills duſky brown, with light edges; the forehead and chin dull white; breaſt pale vinaceous; belly and under tail coverts white; the ſides above the thighs aſh-colour; upper coverts of the tail duſky, edged with brown, and daſhed with aſh-colour; the tail is black, tipped with white, except the two middle feathers, which are wholly of a duſky brown; the outer feather is much ſhorter than the reſt, and white on the exterior web; legs purpliſh red.

The female, in general, is not quite ſo bright in colour, and is rather leſs than the male.

The Turtle viſits the ſouthern parts of England in the ſpring, and remigrates the beginning of September. It chiefly inhabits thick woods, where it makes a neſt in a tree, compoſed of ſticks. The eggs are two in number, and white. Theſe birds ſeem to be more plentiful in Kent than in any other county. Dr. Latham obſerves they are ſeen in that diſtrict to frequent the pea-fields in flocks of twenty or more as ſoon as the peas begin to ripen. We have alſo ſeen ſmall flocks in the

month

month of Auguſt on Romney Marſh, which were ſaid to viſit annually the ſame ſpot about that ſeaſon. It is found, though rarely, as far weſtward as Devonſhire; but we believe not far northward. This ſpecies appears to be ſubject to ſome variety. In the *General Synopſis* mention is made of four or five; if ſo, it is found in China, Manilla, Portugal, Java, and other parts. But ſome of theſe are given as diſtinct ſpecies by other authors.

In the Introduction to this work we have noticed the ſingular glands ſituated under the craw of this bird, ſecreting a lacteal fluid; probably common to all this genus.

The young of this ſpecies do not throw out the black feathers on the neck the firſt year; and the bare ſpace about the eyes is of a grey colour.

DOVE-TURTLE-SPOTTED-NECKED.

This appears to be a mere variety of the Common Turtle. The difference conſiſts in the whole ſide of the neck being black, and inſtead of thoſe feathers being tipped with white, there is a round ſpot of white on each very near the end. Dr. Latham ſays this bird was ſhot in Buckinghamſhire, and that he obſerved one of theſe amongſt ſome birds that came by the laſt expedition to the South Seas; but as it was in a parcel wherein was ſome which belonged to the Cape of Good Hope, it is poſſible that this might come from that place.

Lath. Syn. iv. p. 645. 40. A.
Ind. Orn. ii. p. 606. 47. β.

DUCK. A genus of birds whoſe character are,

Bill broad, depreſſed; the end furniſhed with a horny ſubſtance termed a nail; the edges of the mandibles pectinated.

Noſtrils ſmall, oval.

Tongue broad, fringed near the baſe.

Toes three before, webbed; one behind.

DUCK-BIMACULATED.

Anas glocitans. *Gmel. Syſt.* ii. p. 526. *Ind. Orn.* ii p. 862. 75.

Bimaculated Duck. *Br. Zool.* ii. N° 287. t. 100. f. 2. *Lath. Syn.* vi. p. 521. 75.

The length of this ſpecies is twenty inches; bill deep lead-colour; nail black; irides brown; the crown of the head brown, changeable with green, ending in a ſtreak of brown at the hind-head, with a ſmall creſt; between the bill and eye a round ferruginous ſpot; another ſpot of the ſame, but larger and oblong, behind each ear; throat of a fine deep purple; the reſt of the head bright green, continued in ſtreaks down the neck; the breaſt light ferruginous brown, waved with black; wing coverts aſh-coloured; lower coverts ſtreaked with ruſt-colour; ſcapulars cinereous; quills the ſame, inclining to brown; ſecondaries fine green, ending in a ſhade of black edged with white; twelve feathers in the tail, the two middlemoſt black, the others brown, edged with white; tail coverts deep changeable

changeable green; belly duſky, finely granulated; legs ſmall, yellow; webs duſky.

Mr. Pennant informs us a bird of this deſcription was taken in a decoy.

Said to inhabit part of Siberia and Lake Baikal.

DUCK-BLACK. Vide *Scoter.*

DUCK-BROWN-HEADED. Vide *Golden-eye.*

DUCK-BURROW. Vide *Shieldrake.*

DUCK-COMMON, or WILD.

Anas Borchas. *Lin. Syſt.* i. p. 205. 40. *Gmel. Syſt.* ii. p. 538. *Ind. Orn.* ii. p. 850. 49.

Anas fera. *Brif.* vi. p. 318. 4.—*Ib.* 8vo. ii. p. 447.

Canard ſauvage. *Buf.* ix. p. 115. t. 7. 8.

Wild Duck. *Br. Zool.* ii. N° 279. t. 97.—*Ib.* fol. 175. *Arct. Zool.* ii. N° 494. *Will. Angl.* p. 308. t. 72. 75. *Albin.* ii. t. 10.—*Ib.* i. t. 99. *Lath. Syn.* vi. p. 489. 43. *Pult. Cat. Dorſet.* p. 21. *Walc. Syn.* i. t. 77.—*Don. Br. Birds,* v. t. 124. *Lewin, Br. Birds,* vii. t. 246.

The male bird, Mallard or Drake, as it is called, weighs about two pounds and a half; length near twenty-three inches.

The bill is of a yellowiſh green; irides hazel; the head and upper part of the neck deep gloſſy green, bounded below with a white circle, which almoſt ſurrounds the neck; the lower part of the neck before and breaſt dull purpliſh cheſnut; the back

back is brown; the ſides and ſcapulars white, marked with numerous ſmall undulated lines of brown; the rump, upper and under tail coverts, black; on the wing coverts is a tranſverſe ſtreak of white, edged with another of black; beneath which is the ſpeculum, of a fine purpliſh or violet blue, on the ſecondary quills, which are ſhaded to a black near the ends, and tipped with white, and forms another narrow line of this laſt colour on the wings; the belly is pale grey, minutely ſpeckled with light brown in undulated lines; the tail conſiſts of twenty feathers, the four middle ones are of a gloſſy greeniſh black, and curve upwards in a ſingular manner, and ſo connected as to appear only as two feathers; the others are ſtrait, pointed, and of a greyiſh brown, margined with white.

The female is not ſo large, and of a ruſty brown, ſpotted with duſky black; the ſpeculum on the wing is like the male; but none of the tail feathers are curved; the legs of both ſexes are orange.

The Duck breeds on many of our rivers and lakes; ſometimes at a conſiderable diſtance from the water. It ſcrapes together a little of ſuch vegetables as are contiguous for a neſt, and lays from ten to eighteen eggs of a bluiſh white. At the time of incubation the female plucks the down from her breaſt to line the neſt, and frequently covers the eggs when ſhe leaves them.

It frequently happens that a large variety of this bird is caught in our decoys, or ſhot by the ſportſman; but theſe are only half-domeſticated Ducks, which are obliged to leave the canals or pieces of water belonging to private perſons when they become frozen. Theſe are called Rouen Ducks.

It is obſerveable in moſt kinds of birds whoſe young leave the neſt as ſoon as hatched, that they depoſit their eggs on the ground. There are, however, ſome inſtances in which this ſpecies, the Shieldrake, and perhaps others, occaſionally vary in this particular.

We have been aſſured, by a perſon of undoubted veracity, that a half-domeſticated Duck made a neſt in Rumford Tower, hatched her young, and brought them down in ſafety to a piece of water at a conſiderable diſtance. Others have been known to breed in trees; and we recollect the neſt of this bird being found in the head of an old pollard willow impending the water, from whence the young might readily drop unhurt into their natural element.

Wild Ducks pair; but when domeſticated one male will ſerve ſeveral females.

The number that breed in this country is ſmall in compariſon, ſince the extenſive marſhes and fens have been drained and cultivated; and it is probable the greater part of thoſe caught in our decoys in winter come to us from the more northern parts at that ſeaſon.

Mr.

Mr. Pennant mentions thirty-one thouſand two hundred having been taken in one ſeaſon in decoys in the neighbourhood of Wainfleet. And it appears on record that no leſs than two thouſand ſix hundred and forty-ſix Mallards, or Drakes, were taken in two days near Spalding. Theſe appear to have been young birds before they could fly, called Flappers.

DUCK-CUTHBERT. Vide *Duck-eider.*

DUCK-DUSKY and SPOTTED. Vide *Duck-Harlequin.*

DUCK-EIDER, or EDDER.

Anas molliſſima. *Lin. Syſt.* i. p. 198. 15. *Gmel. Syſt.* ii. p. 514. *Ind. Orn.* ii. p. 845. 35.

Anas S. Cuthberti. *Raii Syn.* p. 141. A. 3. F. *Will.* p. 278. t. 77. F.

Anſer lanugioſus. *Briſ.* vi. p. 294. 13. t. 29. 30. —*Ib.* 8vo. ii. p. 440.

Oie à duret, Eider. *Buf.* ix. p. 103. t. 6.

Great Black and White Duck. *Edw.* t. 98. M. & F.

Eider, *or* Cuthbert Duck. *Br. Zool.* ii. N° 271. t. 95. —*Ib.* fol. 152. t. Q. *Arct. Zool.* ii. N° 480. *Will. Angl.* p. 362. t. 76. *Lath. Syn.* vi. p. 470. 29.—*Ib. Sup.* p. 274. *Walc. Syn.* i. t. 66. *Lewin, Br. Birds,* vii. t. 244.

PROVINCIAL.

Colk.

DUC

This ſpecies is nearly double the ſize of the common Duck; length twenty-two inches.

The bill is black; the top of the head is black, taking in the eyes, and continuing in a line on each ſide, where the feathers project on the bill almoſt as far as the noſtrils; below the nape of the neck, on each ſide, the feathers are of a pale green; the reſt of the head, neck, breaſt, back, ſcapulars, and wing coverts, are white; ſome of the coverts are long, and ſomewhat curved at the ends, falling over the quills, which are black; the under parts from the breaſt are black; tail black; legs dull green. In ſome the baſe of the wings and middle of the back is black.

Theſe birds do not arrive to maturity in plumage till the third, or perhaps the fourth year.

In the firſt the back is white, and the uſual parts, except the crown, black; but the reſt of the body is variegated with black and white.

In the ſecond year the crown of the head is black, and the neck and breaſt ſpotted black and white. Other varieties have alſo been obſerved.

The female weighs about three pounds and a half; the general colour of the plumage reddiſh brown, barred with black; the hind part of the neck marked with duſky ſtreaks; on the wings are two bars of white; belly deep brown, indiſtinctly marked with black; tail duſky; legs black.

The Eider Duck is rarely if ever ſeen in the ſouth of England; it breeds in the north of Scotland,

land, particularly on the weſtern iſles, and on the Farn iſlands on the coaſt of Northumberland, in the months of June and July. It lays five or ſix eggs of a pale greeniſh olive-colour. The neſt is made on the ground, compoſed of marine plants, and lined with down of exquiſite fineneſs, which the female plucks from her body. Sometimes a ſufficient quantity is taken from one neſt to fill the crown of a hat, the weight of which is not more than three quarters of an ounce. This is a conſiderable article of trade from the more northern countries. Its exceſſive lightneſs and elaſticity admirably fits it for the purpoſe of ſtuffing quilts.

We have known the young taken from the Farn iſlands in hopes of domeſticating them; but all attempts proved ineffectual, probably for want of proper food. Is ſaid to feed principally on ſhell-fiſh.

It is found plentiful in all the northern parts; Dr. Latham obſerves even in the higheſt latitudes yet diſcovered.

DUCK-FERRUGINOUS.

Anas rufa. *Faun. Suec.* N° 134.

Anas ferruginea. *Gmel. Syſt.* ii. p. 528. *Ind. Orn.* ii. p. 866. 84.

Red Duck. *Arct. Zool.* ii. N° 576. N.

Ferruginous Duck. *Br. Zool.* ii. N° 285. t. 99. *Lath. Syn.* vi. p. 526. 17.

Red-breaſted Duck. *Lewin, Br. Birds,* vii. t. 224.

This ſpecies weighs twenty ounces.

The bill is long and flatted, a little rounded at the baſe; colour pale blue.

Head, neck, and whole upper parts of the body, reddiſh brown; throat, breaſt, and belly, the ſame, but paler; legs pale blue; webs black.

Mr. Pennant informs us this bird was killed in Lincolnſhire. It is the only ſpecimen on record in England; but is ſaid to be found in Denmark and Sweden.

No mention is made reſpecting the difference of ſexes, or whether that above deſcribed was male or female.

DUCK-FIELD. Vide *Buſtard-leſſer.*

DUCK-GREAT-BLACK. Vide *Duck-velvet.*

DUCK-GREAT-BLACK and WHITE. Vide *Duck-eider.*

DUCK-GREY-HEADED. Vide *Morillon.*

DUCK-GREY-HEADED. Vide *Duck-king.*

DUCK-HARLEQUIN.

MALE.

Anas hiſtrionica. *Lin. Syſt.* i. p. 204. 35. *Gmel. Syſt.* i. p. 534. *Phil. Tranſ.* lxii. p. 417. *Ind. Orn.* ii. p. 849. N° 45. M. F.

Anas torquata. *Briſ.* vi. p. 362. 14.—*Ib.* 8vo. ii. p. 457.

Le Canard à collier. *Buf.* ix. p. 250.

Duſky and ſpotted Duck. *Edw.* t. 99.

Harlequin Duck. *Arct. Zool.* ii. N° 490. *Lath. Syn.* vi. p. 484. 38.

FEMALE.

FEMALE.

Anas minuta. *Lin. Syſt.* i. p. 204. 36. *Gmel. Syſt.* i. p. 534.
Anas torquata. *Gmel. Syſt.* i. p. 514.
Querquedula freti Hudſonis. *Briſ.* vi. p. 469. 4.—*Ib.* 8vo. ii. p. 483.
Le Canard brun. *Buf.* ix. p. 252.
La Sarcelle brune et blanche. *Buf.* ix. p. 287.
Little brown and white Duck. *Edw.* t. 157.
Harlequin Duck (female). *Lath. Syn.* vi. p. 485. 38.

We are enabled with certainty to add this ſpecies of Duck to the liſt of Britiſh birds from the authority of Mr. Sowerby, in whoſe collection of the more rare Engliſh birds we had an opportunity of examining both ſexes, which were killed on the domain of Lord Seaforth in Scotland a few years ſince, and preſented to him by that nobleman.

The great Linnæus had conſidered the female of this ſpecies as diſtinct from his *Anas hiſtrionica*, and has given it under the name of *Anas minuta*; Gmelin has followed him, but not without expreſſing his doubts. Various other authors have conſidered the ſexes as diſtinct ſpecies; but later obſervations have clearly proved, beyond doubt, that the *Anas minuta* is no other than the female Harlequin Duck.

This ſpecies is about the ſize of the Wigeon, but ſhorter; length ſeventeen inches; weight about

twenty ounces. Bill ſmall, an inch and a half long, and black; irides hazel; from the baſe of the bill to the eye a white patch; crown of the head black, bounded by a ferruginous ſtreak; neck black, on each ſide of which is a white line pointing downwards; above that is a white ſpot; round the breaſt is a white band, marked with dots of black behind; this is bounded by a black one; between this and the wings is a tranſverſe line of white; the breaſt bluiſh aſh-colour; back duſky brown, with a purpliſh hue; rump bluiſh black; belly and thighs black; ſides dull orange yellow; wings and tail deep aſh-colour; legs black.

The Harlequin Duck, like moſt of the genus, appears to be ſubject to conſiderable variety, from the deſcription of various authors. The breaſt in ſome is marked with ſemilunar ſtripes of white; the tail is brown, and ſome of the quill-feathers tipped with white; and the ſpot between the bill and eye yellow.

The female is leſs, meaſuring about fourteen inches in length; the forehead, and between the bill and eye, is white, with a ſpot of the ſame behind the ear; the reſt of the plumage above is generally of a duſky brown; upper part of the breaſt and rump inclining to rufous; the lower part of the breaſt and belly barred with pale rufous and white; the lower belly and thighs with rufous and brown; legs duſky.

This bird is an inhabitant of the more northern

parts

parts of the world. It is found in Ruſſia, Iceland, and Greenland, and as far as Kamtſchatka; in America from Carolina to Newfoundland, and at Hudſon's Bay.

Is ſaid to frequent bays and rivers during the ſummer months, and to be fond of ſhady places, making its neſt on ſhore amongſt ſhrubs; particularly about the moſt rocky and rapid torrents. It dives admirably in ſearch of ſmall ſhell-fiſh, on which it feeds, as well as on ſpawn and the *larvæ* of gnats. Has been found frequently in the ſmall rivulets of Hudſon's Bay ninety miles inland; moſt probably frequenting ſuch places in preference to large rivers during the breeding ſeaſon. It flies ſwift, and to a great height, ſo that it is not eaſily taken. The note is a ſort of whiſtle. Lays ten or twelve eggs, like thoſe of the Pigeon. In autumn, when the young are capable of flying, they migrate ſouthward, to paſs the winter in the open ſea; and is probably not uncommon on the northern coaſts of Scotland.

DUCK-HAWK. Vide *Falcon-peregrine*, and *Moor-buzzard*.

DUCK-HOOK-BILLED.

This is a mere variety of the common domeſticated Duck. The only difference is in the bill, which is longer, and bent downwards. There is alſo another variety with divided toes, not connected by a membrane or web.

DUCK-KING.

Anas ſpectabilis. *Lin. Syſt.* i. p. 195. 5. *Gmel. Syſt.* i. p. 907. *Ind. Orn.* ii. p. 845. 36.

Le Canard à tête griſe. *Buf.* ix. p. 253.

Anas freti Hudſonis. *Briſ.* vi. p. 365. 15.—*Ib.* 8vo. ii. p. 458.

Grey-headed Duck. *Edw.* t. 154.

King Duck. *Arct. Zool.* ii. N° 481. *Lath. Syn.* vi. p. 473. *Lewin, Br. Birds,* vii. t. 245.

This ſpecies is not much inferior in ſize to the Eider Duck.

The bill is almoſt two inches long, of an orange colour; at the baſe of the upper mandible is a ridged protuberance, flat on the top, and compreſſed on the ſides, but divided into two, the elevated parts velvety black, paſſing on each ſide to the eyes; the crown of the head and nape are pale aſh-colour; at the baſe of the upper mandible the feathers are pea-green, paſſing backwards on each ſide the neck, and taking in half the eye; beneath which, and round to the chin, the feathers are of a dirty white; but here the two colours are blended, and the white is loſt by degrees in the green; under the chin is a black mark, diverging like the letter V inverted; the reſt of the neck and breaſt are whitiſh; the middle of the back, the belly, and vent, black; wings duſky, on the middle of the coverts a patch of white; quills black; the ſecondaries curve downwards over the quills; the ſhafts deep ferruginous,

ferruginous, on each ſide the outer ones a patch of white; the tail is cuneiform, ſhort, and black; legs duſky.

The female is leſs; the protuberance on the bill not ſo large, nor ſo high-coloured, but the feathery ridge on the top is broader; the whole plumage brown, the middle of each feather duſky; ſix of the leſſer quills are tipped with white, which forms a line of white on the wing; the reſt of the quills and tail brown.

The young birds do not get the gibboſity of the bill, nor the males their mature plumage, for ſome years. Theſe birds are ſubject to vary a little in their ſeveral changes, both with regard to the colour of their bill and plumage; is ſometimes found with a little white on the hind part of the head and on the back.

The King Duck reſides chiefly in the northern parts; are plentiful at Hudſon's Bay, where they breed on the ſides of pools and rivers. The neſt is made of ſticks and moſs, lined with down plucked from their own body. The eggs are five or ſix in number, rather leſs than thoſe of the Gooſe, of a whitiſh colour. It is not unfrequent in the north of Siberia and Kamtſchatka, and common in Greenland, where the down is accounted of equal value to that of the Eider Duck, the fleſh excellent, and the gibbous part of the bill a delicacy. The ſkins ſewed together are uſed for winter garments.

Theſe birds are not uncommon on the coaſts of Norway;

Norway; and we are aſſured by Mr. Pennant and others that it ſometimes frequents the Orkney iſles, which has induced us to give it a place in this work.

DUCK-LITTLE-BROWN and WHITE. Vide *Duck-harlequin.*

DUCK-LONG-TAILED.

Anas glacialis. *Lin. Syſt.* i. p. 203. 30. *Gmel. Syſt.* ii. p. 529. *Ind. Orn.* ii. p. 864. 82.

Anas hyemalis. *Lin. Syſt.* i. p. 202. 29.

Anas caudacuta Harelda. *Raii Syn.* p. 145. 14. *Will.* p. 290.

Anas longicauda iſlandica. *Briſ.* vi. p. 379. 17. —*Ib.* 8vo. ii. p. 460.

Querquedula ferroenſis. *Briſ.* vi. p. 466. 40, f. 2.—*Ib.* 8vo. ii. p. 482.

Canard à longue queue. *Buf.* ix. p. 202.

Sarcelle de Ferroe. *Buf.* ix. p. 278.

Swallow-tailed Shieldrake. *Will. Angl.* p. 364.

Long-tailed Duck. *Br. Zool.* ii. N° 283.—*Ib.* fol. 156. t. Q. 7. *Arct. Zool.* ii. N° 501.—*Ib. Sup.* p. 76. *Edw.* t. 280. & t. 156. *Lath. Syn.* vi. p. 528. 73. *Pult. Cat. Dorſet.* p. 20. *Walc. Syn.* i. t. 73. *Lewin, Br. Birds,* vii, t. 262. *Don. Br. Birds,* v. t. 111.

PROVINCIAL.

Sharp-tailed Duck.

This ſpecies is about the ſize of a Wigeon; length twenty-two inches, including the long feathers

thers of the tail. The bill is black; down the middle and acroſs the tip orange; irides red. The fore part and ſides of the head are reddiſh grey; on each ſide the neck, juſt below the head, is an oval black ſpot; the hind part of the head, the throat, and remaining part of the neck and breaſt, white; back and rump black; ſides of the upper tail coverts white, the middle black; the lower belly and vent white; the ſcapulars white, long, and pointed; the wings chiefly black, with a mixture of cheſnut; the four middle tail feathers are black, the others white; the two middle ones are narrow, and exceed the others three inches and a half; legs of a dull red; claws black.

Such is the deſcription of the male; but in ſome the black parts are more or leſs of a chocolate colour, and the ſpot on the neck occupies half of it; the length of the tail alſo varies.

The female has been deſcribed by ſome authors for a different ſpecies. The bill, however, which is the ſame in this ſex, ſeems to be an unerring guide. The ſides of the head are white, behind cinereous; the reſt of the head, the neck, breaſt, and back, duſky black; the lower part of the breaſt and ſcapulars cheſnut; belly white; upper tail coverts and wings like the male; legs duſky reddiſh brown. This ſex is alſo ſubject to ſome variation; moſt commonly the middle tail feathers are not much longer than the reſt. It is rarely met with in England, but is frequent in the north

of

of Scotland and the Orkneys in winter, where they aſſemble in large flocks; common in Sweden, Lapland, and Ruſſia. Is ſaid to breed in Greenland and at Hudſon's Bay; makes a neſt of graſs near the ſea, and lays ten or more bluiſh white eggs. The down of this bird is ſaid to be as valuable as that of the Eider Duck.

The *trachia* of this ſpecies is ſaid to be of a ſingular ſtructure, having three openings covered with a thin membrane.

DUCK-PINTAIL.

Anas acuta. *Lin. Syſt.* i. p. 202. 28. *Gmel. Syſt.* ii. p. 528. *Ind. Orn.* ii. p. 864. 81.

Anas caudacuta. *Raii Syn.* p. 147. A. 5.—*Ib.* 192. 13. *Will.* p. 289. t. 72.

Anas longicauda. *Briſ.* vi. p. 369. 16. t. 34. f. 1. 2. —*Ib.* 8vo. ii. p. 459.

Canard à longue queue. *Buf.* ix. p. 199. t. 13.

Tritzihoa. *Raii Syn.* p. 175.

Sea Pheaſant, *or* Cracker. *Will. Angl.* p. 376. t. 73. *Albin.* ii. t. 94, 95.

Pintail. *Br. Zool.* ii. N° 282.—*Ib.* fol. 156. t. Q. 8. *Arct. Zool.* ii. N° 500. *Lath. Syn.* vi. p. 526. 72. *Pult. Cat. Dorſet.* p. 21. *Walc. Syn.* i. t. 72. *Lewin, Br. Birds,* vii. t. 261.

The weight of this ſpecies is about two pounds; length twenty inches. Bill black, bluiſh on the ſides; irides dark. The head and upper part of the

the neck before rufous brown; the ſides of the head gloſſed with purple; lower part of the neck before white, running up on each ſide to the hind-head, divided by a brown line down the back of the neck; nape duſky, gloſſed with purple; breaſt and belly white; back and ſides of the breaſt marked with numerous ſmall undulated black and white lines; the ſcapulars black; the inner ones long, pointed, and margined with greyiſh white; ſmaller coverts of the wings fine aſh-colour; the greater coverts of the ſecondary quills tipped with bay; the greater quills duſky brown; leſſer quills gloſſy purpliſh-green on their outer webs, black near the ends, tipped with white; the two middle feathers of the tail are three inches longer than the reſt, narrow, pointed, and black; the others duſky, edged with white; vent black; legs duſky, inclining to lead-colour.

The female is ſomewhat leſs; the head and neck ruſty brown, ſtreaked with duſky; the back and ſcapulars duſky brown, tranſverſely marked with narrow white bars acroſs each feather; the ſpeculum in the wing ſomething like the male, but leſs conſpicuous; the under parts light ruſty-brown, mottled with a deeper ſhade; the tail is brown and cuneiform; the two middle feathers croſſed with one or two pale lines, but are not much longer than the next; the number of feathers are ſixteen.

The

The female is not diſtinguiſhable from the male in plumage.

The Scaup Duck is not uncommon in moſt parts of this kingdom in winter, and is frequently found in freſh waters. Is ſuppoſed to take its name from feeding on broken ſhells, called ſcaup.

This, like moſt of the genus, breed in the more northern parts; is common in Ruſſia, Sweden, Norway, and Lapland; is found at Hudſon's Bay in the warmer months.

DUCK-SHARP-TAILED. Vide *Duck-long-tailed.*

DUCK-SMALLER-RED-HEADED. Vide *Goldeneye.*

DUCK-TUFTED.

Anas Fuligula. *Lin. Syſt.* i. p. 207. 45. *Gmel. Syſt.* ii. p. 543. *Ind. Orn.* ii. p. 868. 90. *Will.* p. 280. t. 73.

Anas criſtata. *Raii Syn.* p. 142. A. 7.

Glaucium minus. *Briſ.* vi. p. 411. 26. t. 37. 1. —*Ib.* 8vo. ii. p. 469.

Morillon. *Buf.* ix. p. 227. 231. t. 15.

Tufted Duck. *Br. Zool.* ii. N° 274.—*Ib.* fol. 153. *Arct. Zool.* ii. p. 573. G. *Albin.* i. t. 95. *Will. Angl.* p. 365. t. 73. *Hayes, Br. Birds,* t. 26. *Lath. Syn.* vi. p. 540. 79. *Pult. Cat. Dorſet.* p. 20. *Walc. Syn.* i. t. 78. *Lewin, Br. Birds,* vii. t. 257.

This

This ſpecies weighs about twenty-five ounces; length ſeventeen inches.

The bill is near two inches long, of a bluiſh lead-colour, broadeſt towards the point; nail black; irides black yellow.

The head, neck, upper part of the breaſt, and whole upper part of the body, black; the two firſt tinged with changeable violet and green; on the back of the head a pendant creſt of very narrow feathers, ſometimes two inches long; the back and ſcapulars are very minutely ſpeckled with grey, not diſtinguiſhable at a ſmall diſtance; the three or four firſt greater quills black on their outer webs, the reſt more or leſs white towards the baſe; the ſecondaries, except four or five next the body, white, tipped with black, and ſlightly edged with the ſame; lower breaſt and belly gloſſy white; above the thighs, and behind the vent, black; the tail is ſomewhat cuneiform, and conſiſts of fourteen black feathers; the whole upper parts of the bird have a ſlight tinge of violet, when viewed in ſome particular lights; legs and feet duſky black.

This bird is ſubject to much variety in ſize and colour; will ſometimes weigh as much as thirty-one ounces, others not above twenty-two; and the black on the breaſt is intermixed with brown, as well as the plumage in general being of a ruſty brown: ſuch are probably birds of the firſt or ſecond year.

The female reſembles the other ſex in markings;

but where that is black this is of a ruſty brown; the ſides are alſo brown, which in the male are white; and the head wants the creſt. This, however, is ſometimes the caſe in young male birds.

The Tufted Duck is not uncommon with us in winter; is frequently ſeen in our freſh waters as late as the latter end of March; are difficult to ſhoot by reaſon of their inceſſant diving. It is often brought to market and ſold for Widgeon. Is found in all the northern parts of Europe as far as Norway, and throughout Ruſſia.

Scopoli has noticed a variety as large as a Mallard, with a yellowiſh bill; and another, of a leſs ſize, with a band or ſpeculum on the wing of ſhining green, above which is a ruſous one, and beneath a black band.

It is much to be doubted if theſe are the ſame ſpecies; ſuch does not ſeem to have been noticed in England; and indeed the great ſpeculum in the wing of the latter ſhould forbid it, as this in general is a pretty clear mark of diſtinction in the Duck tribe. The male of this ſpecies poſſeſſes a *labyrinth*.

DUCK-VELVET.

Anas fuſca. *Lin. Syſt.* i. p. 196. 6. *Gmel. Syſt.* ii. p. 507. *Ind. Orn.* ii. p. 848. N° 44.

Anas nigra major. *Raii Syn.* p. 141. A. 4. *Will.* p. 278. t. 70. *Briſ.* vi. p. 423. 29.—*Ib.* 8vo. ii. p. 472.

Turpan.

Turpan. *Buf.* ix. p. 291?
Grande ou double Macreuſe. *Buf.* ix. p. 242.
Great black Duck. *Will. Angl.* p. 363. t. 70.
Velvet Duck. *Br. Zool.* ii. N° 272. t. 96.—*Ib.* fol. 152. *Artt. Zool.* ii. N° 482.—*Ib. Sup.* p. 75. *Lath. Syn.* vi. p. 482. 37.—*Ib. Sup.* p. 274. *Pult. Cat. Dorſet.* p. 20. *Walc. Syn.* i. r. 58. *Lewin, Br. Birds,* vii. t. 247.

PROVINCIAL.

Black Diver.

This is rather larger than the common Drake; length about twenty inches.

The bill broad, with a black knob at the baſe; the reſt of the bill is yellow; the nail red; the edges all round black.

The plumage is black, inclining to brown on the belly; under each eye is a white mark paſſing backwards in a ſtreak; acroſs the middle of the wing is a band of white; legs red; claws black.

The female is more inclining to brown, and the protuberance on the bill is wanting.

This ſpecies much reſembles the Scoter, only that bird has no white feathers about it, and the colour of the bill is ſomewhat different. In the windpipe of this bird is a ſingular bony ſwelling the ſize of a ſmall walnut, ſituated about two-thirds of the length from the *larynx*; immediately under the *larynx* is another oblong bony cavity of

any ſpecies. The tail of the Gooſander is ſaid to be compoſed of eighteen feathers; whereas this bird has twenty feathers in that part.

From theſe circumſtances we cannot heſitate in our opinion of theſe birds being diſtinct ſpecies.

Dr. Latham obſerves that Dr. Heyſham has proved, by diſſection, that ſome of the larger Dun-divers were males, and that in Cumberland this bird is infinitely more common than the Gooſander; at leaſt ten or fifteen of the firſt to one of the laſt. We muſt, however, obſerve, that in many of the aquatic birds, in particular the young, do not arrive to maturity in plumage till the third, or perhaps fourth year, before which the males moſt reſemble the female in feathers.

In order to clear up all doubt indiſputably would be to prove, by diſſection, there are female Gooſanders, which does not at preſent ſeem to be ſatisfactorily aſcertained.

We have never had an opportunity of obſerving the tail of the Gooſander; but if it is true that it has only eighteen feathers, it will certainly be an unerring mark of diſtinction, for we can ſpeak with certainty as to the Dun-diver having twenty.

This bird is ſaid to be common in Germany; but moſt probably breeds in the lakes of the more northern parts of the world.

DUNG-BIRD. Vide *Hoopoe.*

DUNG-HUNTER. Vide *Gull-arctic.*

DUNKER,

DUNKER, or DUN-CUR. Vide *Pochard.*

DUNLIN.

Tringa alpina. *Lin. Syſt.* i. p. 249. 11. *Gmel. Syſt.* ii. p. 676. *Ind. Orn.* ii. p. 736.

Cinclus torquatus. *Brif.* v. p. 216. 11. t. 19. f. 2.—*Ib.* 8vo. ii. p. 268.

Gallinago anglicana. *Brif.* v. p. 309. 5.—*Ib.* 8vo. ii. p. 288.

Le Cincle. *Buf.* vii. p. 553.

Le Brunette. *Buf.* vii. p. 493.

Dunlin. *Br. Zool.* ii. N° 205.—*Ib.* fol. p. 126. t. E. 1. f. 2. *Arct. Zool.* ii. N° 391. *Raii Syn.* p. 109. A. 11. *Will.* p. 226.—*Ib. Angl.* p. 305. *Lath. Syn.* v. p. 185. 33.—*Ib. Sup.* p. 249. *Walc. Syn.* ii. t. 151. *Lewin, Br. Birds,* v. t. 176.

This ſpecies of Sandpiper weighs from nine to eleven drams; length of the largeſt eight inches. The bill full an inch in length, ſlender, a little bent downwards, and of a duſky colour; irides duſky.

The upper part of the head, and whole plumage above, ferruginous brown; the former and hind-neck ſtreaked with duſky, the reſt ſpotted with black; the cheeks, underſide of the neck, and breaſt, whitiſh, ſtreaked with black; thoſe on the neck are ſmall, and more inclining to duſky; the wing coverts greyiſh brown, edged with lighter; the quills black; the four firſt wholly ſo, the reſt

[illegible]

[illegible]

[illegible]

Greenland, Iceland, and Scandinavia; and on the Siberian Alps, as well as at Hudſon's Bay.

DUNNOCK. Vide *Warbler-hedge.*

E.

EAGLE. A name given to the larger ſpecies of the Falcon genus, from which it differs in nothing but ſize.

EAGLE-BLACK. Vide *Eagle-ring-tailed.*

EAGLE-CINEREOUS.

Vultur Albicilla. *Lin. Syſt.* i. p. 123. 8.

Falco Albicilla. *Gmel. Syſt.* i. p. 253. *Ind. Orn.* i. p. 9. 2.

Aquila Albicilla, ſeu Pygargus. *Briſ.* i. p. 427. 5. —*Ib.* 8vo. i. p. 123. *Raii Syn.* p. 7. 5. *Will.* p. 31.

Le grand Pygargue. *Buf.* i. p. 99.

White-tailed Eagle. *Will. Angl.* p. 61.

Cinereous Eagle. *Br. Zool.* i. N° 45. t. 18. *Arct. Zool.* ii. p. 214. B. *Lewin, Br. Birds,* i. t. 4. *Lath. Syn.* i. p. 33. 8.—*Ib. Sup.* p. 11.

Vultur Albicllia. *Walc. Syn.* i. t. 1.

This ſpecies is rather leſs than the Golden Eagle. The bill, cere, and irides, of a pale yellow; a bare ſpace between the bill and eye of a bluiſh colour; the head and neck of a pale aſh-colour; the body and

and wing coverts cinereous, clouded with brown; the quills dusky; tail dusky at the base, white at the end; the legs of a light yellow, feathered but little below the knee. The male is rather darkest in colour.

The bill of the Cinereous Eagle is rather longer, and more strait towards the base, than usual in this tribe, which probably induced Linnæus to class it with the Vulture; but as no part of the head or neck is bare of feathers (the characteristic mark of that genus), we have followed the example of later authors, and continued it amongst the Eagles or Falcon genus.

Inhabits Scotland and the Orknies. An acquaintance had two of these birds, which were taken from the highest cliffs in the last mentioned place; there were three in the nest. Dr. Latham says Dr. Heysham informed him that in a nest of one of these bids, near Keswick in Cumberland, was found a trout of about twelve pounds weight; that he obtained the bird alive, and had kept it above ten years when he communicated the account; and that it was either six or seven years before the tail became white. Those from which our description was taken were about three years old.

It is said to prey on land animals as well as fish and aquatic birds.

Inhabits the southern parts of Russia, particularly about the Wolga; found also in Sweden, Denmark, Iceland, and Lapland.

In Greenland it remains the whole year amongſt the iſlands and rocks, where it preys on the aquatic birds, darting on them as ſoon as they riſe on the ſurface of the water after diving; attempts ſometimes to ſeize a Seal, which proves its own deſtruction, being drawn under water by that animal.

Mr. Denne, of Littlebourne Court in Kent, favoured us with a live ſpecimen he ſhot in March 1797. From its inferior ſize and dark colour it is probably a male. The tail feathers are ſtill duſky at the baſe; the irides hazel.

EAGLE-FISHING. Vide *Oſprey.*

EAGLE-GOLDEN.

Falco Chryſactos. *Lin. Syſt.* i. p. 125. 5. *Gmel. Syſt.* i. p. 256. *Raii Syn.* p. 6. 1. *Will.* p. 27. t. 1. *Ind. Orn.* i. p. 12. 8. *Briſ.* i. p. 431. 7. —*Ib.* 8vo. p. 124.

Le grand Aigle. *Buf.* i. p. 76.

Golden Eagle. *Br. Zool.* i. N° 42. t. 16.—*Ib.* fol. p. 61. t. A. *Arct. Zool.* ii. p. 214. A. *Albin.* ii. t. 1. *Lewin, Br. Birds,* i. t. 2. *Lath. Syn.* i. p. 31.—*Ib. Sup.* p. 10. *Walc. Syn.* i. t. 3.

The general weight of this ſpecies is about twelve pounds; the length three feet and a half; breadth eight feet. The bill is of a bluiſh black; cere yellow; irides hazel.

The whole bird is of a dark brown; the feathers about

about the back of the head and neck long and narrow, bordered with tawny; quill-feathers chocolate brown; tail deep brown, daſhed with aſh-colour, and white at the roots; legs yellow, ſhort, ſtrong, and covered with feathers to the feet, the great characteriſtic diſtinction between this and the Sea Eagle.

The Golden Eagle is ſaid to be not unfrequent in the mountainous parts of Ireland and Scotland; but we ſuſpect it to be more rare than is generally imagined, and has undoubtedly been confounded with the Sea Eagle.

It breeds in the moſt inacceſſible rocks, and lays three or four white eggs.

Smith, in his hiſtory of Kerry, ſays a poor man in that county got a comfortable ſubſiſtence for his family during a ſummer of famine out of an Eagle's neſt.

Inſtances have been recorded of infants being carried to their neſts. And in order to extirpate theſe birds there is a law in the Orkney iſles which entitles any perſon who kills an Eagle to a hen out of every houſe in the pariſh in which it is killed.

Theſe birds are remarkable for their longevity and abſtinence from food. Mr. Pennant mentions an inſtance of one enduring hunger for twenty-one days.

EAGLE-RING-TAIL.

Falco fulvus. *Lin. Syſt.* i. p. 125. 6. *Gm. Syſt.* i. p. 256. *Ind. Orn.* i. p. 10. 4.

Chryſaë

Chryſaëtos, cauda annulo albo cincta. *Raii Syn.* p. 6. 2. *Will.* p. 28.

Aquila. *Briſ.* i. p. 419. 1.—*Ib.* 8vo. 121.

L'Aigle commun. *Buf.* i. p. 86.

Black Eagle. *Br. Zool.* i. N° 87. *Arct. Zool.* ii. N° 87.

Ring-tail Eagle. *Br. Zool.* fol. p. 62. *Will. Angl.* p. 59. *Lewin, Br. Birds,* i. t. 3. *Lath. Syn.* i. p. 32.—*Ib. Sup.* p. 10. *Walc. Syn.* i. t. 4. *Pult. Cat. Dorſet.* p. 2.

White-tailed Eagle. *Edw.* t. 1.

This ſpecies is nearly equal in ſize to the Golden Eagle. In the bill, cere, irides, and legs, it reſembles thoſe of that bird.

The plumage is rather darker; and the tail is white for two-thirds of its length; the end is black. Mr. Pennant informs us it is frequent in Scotland, where it is called the Black Eagle; and adds, that it is very deſtructive to Deer, which it will ſeize between the horns, and by inceſſantly beating it with its wings, ſoon makes a prey of the haraſſed animal; that it builds in cliffs of rocks near the deer foreſts, and makes great havoc not only amongſt them but alſo the white Hares and Ptarmigans.

Mr. Willughby gives a curious account of the neſt of this ſpecies found in the woodlands near the river Derwent in the Peak of Derbyſhire. He ſays it was made of large ſticks, lined with two

layers

layers of ruſhes, between which was one of heath; that in it was one young and an addle egg, and by them a Lamb, a Hare, and three Heath-poults. The young eagle was black, having a white ring about the tail.

From this mark in ſo young a bird it ſhould appear to be a characteriſtic of the ſpecies in all ages; for though it varies in the breadth of that band, yet it never entirely loſes it.

This bird is as rarely found in South Britain as the Golden Eagle.

Inhabits many parts of Europe and America.

EAGLE-SEA.

Falco oſſifragus. *Lin. Syſt.* i. p. 124. 3. *Gmel. Syſt.* i. p. 255. *Raii Syn.* p. 7. 3. *Will.* p. 29. t. 1. *Ind. Orn.* i. p. 12. 7.

Aquila oſſifraga. *Briſ.* i. p. 437. 9.—*Ib.* 8vo. p. 125.

L'Orfraie. *Buf.* i. p. 112. t. 3.

Sea Eagle, *or* Oſprey. *Br. Zool.* i. N° 44. t. 17.—*Ib.* fol. t. 63. *Arct. Zool.* ii. N° 86. A. *Will. Angl.* p. 59. t. 1. *Lewin, Br. Birds,* i. t. 1. *Lath. Syn.* i. p. 30.—*Ib. Sup.* p. 9. *Wale. Syn.* i. t. 2. *Pult. Cat. Dorſet.* p. 2. *Don. Br. Birds,* v. t. 105.

This ſpecies is full as large as the Golden Eagle. The bill is bluiſh black; cere, ſides of the mouth, orbits, yellow; irides light hazel. The feathers on

on the head and upper part of the neck are long and narrow, duſky brown at their ends, tawny towards the baſe, and white at the roots; the whole body dark brown, intermixed with ruſt-colour; the tail and its coverts mottled with yellowiſh white, dark and faint aſh-coloured brown; the quills are of a dark chocolate colour; the ſhafts white towards the baſe; the legs ſtrong and yellow, feathered very little below the knee, and meaſuring two inches in circumference; the claws black; the inner one, which is largeſt, is two inches long, much hooked, and nearly one inch round at the baſe.

The ſpecimen from which this deſcription is taken was killed by Sir Robert Littleton's game-keeper in Shropſhire, early in the ſpring 1792, and preſented to me by Lord Valentia, now in my muſeum.

It was accompanied by a letter from Sir Robert, the purport of which was, that his ſervant being out ſhooting ſaw two large birds feeding on the carcaſe of a ſheep, which appeared recently killed; that having nothing but ſmall ſnipe-ſhot with him he turned back, intending to go home for larger; that the eagles then followed him, and frequently came ſo near that he concluded they meant either to attack him or his dogs. Suddenly loſing ſight of one, he judged it was very near him behind, and being ſomewhat alarmed, turned round and ſhot at it in a hurry; after which the bird flew ſome hundred

dred yards, and dropped. On his approach it was vomiting blood; and he killed it after a ſtruggle of half an hour. He adds, that it was the largeſt of the two.

The other Eagle continued in the neighbourhood ſome time after, and rooſted in the high trees of a wood belonging to Sir Robert Littleton.

Another of this ſpecies was ſhot in Epping Foreſt a few weeks before. Others have been frequently killed in the New Foreſt; and we are informed ſcarce a year paſſes without one being ſeen in that part; two of which we have ſeen nailed up in the hall of the Lodge at Lyndhurſt.

John Maxwell, Eſq. of Ardbraccan in Ireland, favoured us with two young birds of this ſpecies alive, taken the preceding year on a mountainous precipice, or craggy cliff, called Slieve Donald, impending the ſea in the county of Down.

That gentleman informed us, two men, covered with ſackcloth and armed, were lowered by ropes to the area, which, with conſiderable difficulty, they robbed of two young, leaving only one addled egg behind.

The old Eagles being ſo furious as to create ſerious alarm, neither the neſt or colour of the egg were noticed. Some fragments of fleſh were in the neſt.

The Eaglets were covered with a gloſſy, dark, murry-coloured down (as it was termed). A baſket was attached to the ropes that conveyed the men down;

down; into this the young birds were put; but from the inceſſant violence and amazing ſtrength of the parent birds, were with difficulty carried off.

Theſe birds were not twelve months old when we received them. On their firſt moulting they became much darker, particularly about the breaſt and thighs, the latter almoſt wholly of a duſky black; at two years old the baſe of the bill became yellow; in the third year there was not any material change. At this time one of them killed the other, and devoured it; probably neglected to be fed, for they before lived together in perfect harmony.

The food of this bird is ſaid to be principally fiſh; but it is probable every animal of inferior ſtrength ſuffers from its rapacity.

It is not uncommon in Scotland and Ireland, and breeds generally in the neighbourhood of large lakes, or on the ſea-coaſt amongſt the moſt ſtupendous cliffs.

Between the upper and lower lakes of Killarney is a rock called the Eagle's Neſt, originating from the circumſtance of its breeding there annually.

This bird is ſaid to watch the Oſprey catching fiſh, when it purſues that bird till he quits his prey, which it ſeizes moſt dexterouſly in the air.

From the aſtoniſhing height theſe and ſome other birds fly, we are led to believe they are capable of living in a much lighter air than other animals. From the top of ſome of the higheſt mountains in Scotland we have ſeen ſeveral ſoaring together at

ſo great a diſtance as to appear ſcarce larger than a Swallow.

EAGLE-WHITE-TAILED. Vide *Eagle-cinereous*, and *Eagle-ring-tail.*

EASTERLING. Vide *Smew.*

EBB. Vide *Bunting-common.*

EGRET

Ardea Garzetta. *Lin. Syſt.* i. p. 237. 13. *Gmel. Syſt.* ii. p. 628. *Raii Syn.* p. 99. 5. *Will.* p. 206.—*Ib. Angl.* p. 280. *Ind. Orn.* ii. p. 694. 64.

Egretta. *Briſ.* v. p. 431. 16.—*Ib.* 8vo. ii. p. 322.

L'Aigrette. *Buf.* vii. p. 372. t. 20.

Little Egret. *Br. Zool. App.* t. 7. *Arct. Zool.* ii. N° 347. *Lath. Syn.* v. p. 90. 59. *Walc. Syn.* ii. t. 30. *Lewin, Br. Birds,* iv. t. 149. *Don. Br. Birds,* iv. t. 98.

This ſpecies of Heron weighs about one pound; length near a foot.

The bill is black; irides yellow; the bare ſpace about the eyes green. The feathers on the back of the head are long, forming a pendent creſt; two of the feathers are nearly five inches long, and narrow; the whole plumage is of a pure white; the feathers on the breaſt and ſcapulars are of a ſoft and looſe texture; legs greeniſh black; claws black.

If we may judge from the bill of fare of the famous

mous feaſt given by the Archbiſhop Nevil, theſe birds were plentiful in this country formerly, for no leſs than one thouſand were in that liſt. It is, however, now become a very rare bird in this kingdom; one inſtance only of its being killed in theſe realms is on record by modern authors, and that in Angleſea. In the firſt year this bird is ſaid to be ſlate-coloured; the ſecond year grey, ſpotted with white. It is not uncommon in many parts of Europe and Aſia; is found alſo in Africa, and in America about New York and Long Iſland, ſome of the Weſt India iſlands, and Cayenne.

The feathers of this bird were formerly much eſteemed as ornaments for the head-dreſs, eſpecially thoſe of the ſcapulars.

ELK. Vide *Swan-whiſtling.*

EMMET-HUNTER. Vide *Wryneck.*

ERN. Vide *Eagle-cinereous.*

F.

FALCON. A genus of rapacious birds, the characters of which are,

Bill ſtrong, hooked; the baſe covered with a bare ſkin called the cere.

Tongue bifid at the end.

Noſtrils ſmall, oval, placed in the cere.

Toes, middle toe connected to the outmoſt as far as the firſt joint.
Claws large, much hooked.
Female largeſt.

FALCON-ASH-COLOURED. *Falco cinereaceus.*
Falco hyemalis. *Gmel. ſyſt.* p. 274? *Ind. Orn.* i. p. 35. N° 73?
Northern Falcon. *Lath. Syn.* i. p. 79?
Winter Falcon. *Arct. Zool.* ii. p. 209?

Size rather inferior to the female Sparrow-hawk, but of a greater length and more ſlender make. The bill is ſmall and black; cere and irides yellow. Head, neck, breaſt, and whole upper parts, dark browniſh grey, the tips of each feather darkeſt; the belly, ſides, thighs, and under tail coverts, are white, with large ferruginous ſtreaks down their middle; greater quills black; ſome of the greater coverts impending them duſky black on the outer webs towards their middle, forming a ſmall bar on the wings; the two middle feathers of the tail above like the back; the three next, on each ſide, incline more to light grey; the outer ones almoſt white; all more or leſs barred with ferruginous, except the exterior one; the under part of the tail is of a light colour, and the bars not ſo conſpicuous; legs long, ſlender, yellow; claws ſhort, black, and not greatly hooked for a predacious bird.

This ſpecies was ſhot in Wiltſhire in the winter,

and

and we received it for the Henharrier; which indeed at firſt ſight it might readily be miſtaken for in the ſtate of changing its plumage from the Ringtail; but the want of the wreath of ſhort feathers round the head, always conſpicuous in both ſexes of that bird, at once diſtinguiſhes it; the ſize is alſo much inferior, though the tail is full as long, and the wings much longer. And, as an additional mark of diſtinction, the wings, when cloſed, reach beyond the extremity of the tail, which in the Henharrier are far ſhort of the tail. The wing is alſo more pointed, the third feather being much longer than any other; whereas in the Henharrier the point of the wing is more rounded by reaſon of the third and fourth feather being of equal length.

From the ſtate we received this bird in it was impoſſible to aſcertain the ſex. It was, however, in tolerable good plumage, and is now in good preſervation.

We cannot find this ſpecies deſcribed by any one except it is the Northern Falcon of Latham, *Synopſis*, Vol. I. N° 62, to which it bears ſome ſimilitude, but the ſize of that bird is much ſuperior. This, however, may be reconciled by ſuppoſing them to be of different ſexes.

Mr. Pennant has made his Winter Falcon of the *Arctic Zoology* to be a variety of the Northern Falcon, occaſioned by age or ſex. The deſcription, however, is materially different.

Gmelin has quoted both the above authors for

his *Falco Hyemalis*; and Dr. Latham, in his *Index Ornithologicus*, has adopted that name. It is ſpoken of as a native of North America, but does not ſeem to be much known. Viſits New York in winter, and retires in the ſpring.

The bird from which this deſcription is taken differs in the marking of the tail from thoſe of the before-quoted authors; but leaſt ſo from that of the *General Synopſis*. The bars are four in number, but theſe are moſtly confined to the inner webs, and not obſervable on the two middle feathers or the exterior one, and are mere ſpots on the ſecond outer feather.

FALCON-BLUE-BACKED. Vide *Falcon-peregrine.*

FALCON-GENTIL.

Falco gentilis. *Lin. Syſt.* p. 126. *Gmel. Syſt.* p. 270. *Raii Syn.* p. 13. 5. *Will.* p. 46. *Ind. Orn.* p. 29. 66. *Briſ.* i. p. 339.—*Ib.* 8vo. p. 98.

Falco montanus. *Raii Syn.* p. 13? *Will.* p. 45. t. 5?

Gentil Falcon. *Br. Zool.* i. N° 50. t. 21. 22. *Arct. Zool.* ii. N° 98. *Albin.* ii. t. 6. *Will. Angl.* p. 79. *Lath. Syn.* i. p. 64.—*Ib. Sup.* p. 17. *Lewin, Br. Birds*, i. t. 11. *Walc.* i. t. 11. *Pult. Cat. Dorſet.* p. 2.

This bird has been conceived by ſome to be the young of ſome other ſpecies.

Buffon

Buffon is of opinion that it is the Common Falcon in full feather; others have ſuppoſed it to be the young of the Goſhawk immatured.

It is a ſhort-winged hawk, like that of the Goshawk, the wings reaching only half the length of the tail.

The ſize is that of the laſt mentioned bird. Bill lead-colour; cere and irides yellow. Head light ferruginous, with oblong black ſpots; the whole under ſides yellowiſh white; the points of each feather marked with heart-ſhaped duſky ſpots; back brown; quills duſky, barred with black on the outer webs, and with white on the lower part of the inner webs; coverts of the wings and ſcapulars brown, edged with ruſt-colour; the tail is barred with black and cinereous alternately, the tip white. A variety, ſuppoſed to be young birds, are deſcribed to have tranſverſe bars on the breaſt. Is ſaid to make its neſt in rocks, and to inhabit the north of Scotland.

FALCON-GREY.

Falco griſeus. *Gmel. Syſt.* i. p. 275. *Ind. Orn.* i. p. 37. 86.

Grey Falcon. *Br. Zool.* i. N° 49.—*Ib.* fol. p. 65. *Lewin, Br. Birds,* i. t. 15. *Lath. Syn.* i. p. 82. 67.

Mr. Pennant firſt gave this bird to the public, from whoſe authority later ornithologiſts have con-

ıued it as a diſtinct ſpecies. It ſeems, however, ıe above author did not ſee the ſpecimen he has ecorded, but ſays it was ſhot near Halifax 1762, and the following account tranſmitted to him by Mr. Bolton.

This bird was about the ſize of a Raven; the bill was ſtrong, ſhort, much hooked, and of a bluiſh colour; the cere and edges of the eyelids yellow; the head was ſmall, flatted at the top; the fore pait of a deep brown; the hind part white; the ſides of the head and throat were cream-coloured; the belly white, marked with oblong black ſpots; the hind part of the neck and the back were of a deep grey; the wings were very long, and when cloſed reached beyond the tra'n; the firſt of the quill-feathers were black, with a white tip; the others were of a bluiſh grey, and their inner webs irregularly ſpotted with white; the tail was long, and wedge ſhaped; the two middle feathers, being the longeſt, were plain (the colour not mentioned), the reſt ſpotted; the legs were long, naked, and yellow.

We cannot help doubting whether this bird may not be a mere accidental variety of ſome other ſpecies of Falcon; and we are the more inclined to be of that opinion as only one ſingle inſtance of ſuch a bird has been mentioned by any writer; and we know this claſs of birds is ſubject to much variety.

FALCON-GYR. Vide *Falcon-jer.*

FALCON-HAGGARD. Vide *Falcon-peregrine.*

FALCON-JER.

Falco iſlandicus. *Ind. Orn.* i. p. 32. 69.
Falco candicans. *Gmel. Syſt.* ii. p. 275.
Gerfaulk. *Buf.* i. p. 241.
Gyrfalco. *Raii Syn.* p. 13. 3.
Gerfalco. *Will.* p. 44. t. 8. *Briſ.* i. p. 370. t. 30. f. 2 —*Ib.* 8vo. p. 108.
Gyrfalcon. *Br. Zool.* i. N° 47. t. 19. *Arct. Zool.* ii. p. 221. E. *Lewin, Br. Birds*, i. t. 16.
Iceland Falcon. *Lath. Syn.* i. p. 71. 50 A. B.
White Jerfalcon. *Lath. Syn.* i. p. 83. 84.—*Ib. Sup.* p. 21.
Gyrfalcon. *Walc. Syn.* i. t. 15.

This ſpecies is ſuperior in ſize to the Goſhawk. Bill bluiſh, tip black; cere blue aſh-colour; irides duſky.

The plumage is generally white, ſpotted with brown; legs bluiſh aſh colour.

This bird is ſubject to variety, either from age, ſex, or climate. In the northern countries it is found quite white. Others are ſaid to be brown above, white beneath, ſpotted with brown; tail grey, with tranſverſe brown lines.

Mr. Pennant ſays the bill is yellow; the whole plumage white, marked with duſky lines, ſpots, or bars; the head, breaſt, and belly, with narrow lines, thinly ſcattered and pointing down; the wings with large heart-ſhaped ſpots; the legs pale blue, and feathered a little below the knees.

Said

FALCON-ROUGH-LEGGED.

Falco lagopus. *Gmel. Syſt.* i. p. 260. *Ind. Orn.* i. p. 19. 33.

Rough-legged Falcon. *Br. Zool. App.* t. 1. *Arct. Zool.* ii. p. 200. 92. *Lewin, Br. Birds*, i. t. 14. *Lath. Syn.* i. p. 75.—*Ib. Sup.* p. 18. *Walc. Syn.* i. t. 14.

Greenland Falcon. *Arct. Zool.* ii. p. 220. E.

This ſpecies meaſures upwards of two feet in length; the wings are long, and reach near to the end of the tail when cloſed.

The bill is duſky; cere and irides yellow. The head, neck, and breaſt, yellowiſh white, ſtreaked with brown, thoſe on the breaſt large; the lower part of the ſides above the thighs, and belly, except a line down the middle, duſky brown; the ſcapulars and wing coverts blotched with duſky brown, the former mixed with yellowiſh white, the latter inclining to ferruginous; the quill-feathers white at the baſe, duſky black at the ends; the outer webs daſhed with cinereous, ſhafts white; the tail is brown one-third from the end, acroſs which are two faint bars of duſky black, the reſt white, with a few ſpots acroſs the upper part, reſembling a broken bar, of brown; the tip white; upper tail coverts white, ſtreaked with brown; the legs are covered with pale dull yellow feathers down to the feet, ſpotted with brown; thighs the ſame; feet yellow; claws black.

This

This bird appears to be ſubject to ſome variety. That deſcribed by Mr. Pennant, ſhot near London, had the extreme half of the tail brown, tipt with dirty white.

Another, ſhot in Suffolk, had the tail of a cream-coloured white; near the tip is a brown bar above an inch in breadth; above that another half an inch broad; and above theſe each feather had a ſpot upon it in the middle, mimicking, when ſpread, a third bar; the two outer feathers on each ſide are marked with a few irregular ſpots of brown on the outer webs, almoſt the whole of their length. This bird was leſs than the former, meaſuring only one foot ten inches.

The Rough legged Falcon is a native of the more northern parts, and is rarely met with in England. That from which the firſt deſcription is taken was picked up dead on the coaſt of Kent in the winter of 1792, preſented to us by Dr. Latham, and now in my muſeum.

Said to be a native of Denmark.

FALCON-SPOTTED.

Falco verſicolor. *Gmel. Syſt.* i. p. 272. *Ind. Orn.* i. p. 33. 73.

Spotted Falcon. *Br. Zool.* i. N° 55. t. 26. *Lewin, Br. Birds*, i. t. 13. *Lath. Syn.* i. p. 74. *Walc. Syn.* i. t. 13.

This ſpecies is about the ſize of a Buzzard.

Bill

Bill black; cere yellow; irides pale yellow. The crown of the head, and hind part of the neck, white, ſpotted with light reddiſh brown; back and ſcapulars the ſame, edged with white; quill-feathers duſky, barred with aſh-colour; under ſide of the neck, breaſt, belly, thighs, and rump, white; on the two firſt are a few ruſt-coloured ſpots; the middle feathers of the tail barred with white and deep brown; the others with light and darker brown; the legs are ſtrong.

Two of theſe birds were ſhot at Longnor in Shropſhire, and firſt deſcribed by Mr. Pennant.

FALCON-WHITE. Vide *Falcon-jer.*

FALK. Vide *Razorbill.*

FALLOW-SMICH. Vide *Wheatear.*

FLASER. Vide *Gull-arctic.*

FIELDFARE.

Turdus pilaris. *Lin. Syſt.* i. p. 291. 2. *Gmel. Syſt.* ii. p. 807. *Raii Syn.* p. 64. A. 3. *Will.* p. 138. t. 37. *Ind. Orn.* i. p. 330. 11. *Briſ.* ii. p. 214. 5.—*Ib.* 8vo. i. p. 217.

La Litorne, *ou* Tourdelle. *Buf.* iii. p. 301.

Fieldefare, *or* Feldefare. *Br. Zool.* i. N° 106. —*Ib.* fol. 90. t. P. 2. f. 1. *Arct. Zool.* ii. p. 340. A. *Will. Angl.* p. 188. t. 37. *Albin.* i. t. 36. *Hayes, Br. Birds,* t. 31. *Lewin, Br. Birds,* ii. t. 60. *Lath. Syn.* iii. p. 24. 11. *Pult. Cat. Dorſet.* p. 10. *Walc. Syn.* ii. t. 200.

PROVINCIAL.

Pigeon Fieldfare.

This ſpecies of Thruſh is in ſize between that of the Miſſel and the Throſtle; length ten inches; weight four ounces.

The bill is yellowiſh, black at the point; irides hazel. The head and hind part of the neck cinereous, the firſt ſpotted with black; the back and leſſer coverts of the wings cheſnut-brown; rump aſh colour; quill-feathers duſky brown, with paler edges; the fore part of the neck, breaſt, and ſides, yellowiſh, ſtreaked with duſky; throat white; belly and vent the ſame; tail duſky black; the middle feathers daſhed with cinereous; legs black.

The female has leſs of the rufous tinge on the breaſt; in other reſpects like the other ſex.

This is a migrative ſpecies, viſits us together with the Redwing in large flocks in October, and frequently remains till the beginning of April. With us its principal food is the fruit of the hawthorn and other berries, worms, and inſects. In very ſevere weather they migrate further ſouth; but a ſudden fall of ſnow ſometimes deprives them of the power of flying acroſs the ſeas to a warmer climate; in which caſe thouſands are ſtarved. In the winter of 1798 a very heavy ſnow fell to the northern and eaſtern parts of this country, when prodigious flocks appeared in the weſt of England; but as that part was ſoon after covered with ſnow, which laſted on the ground for a conſiderable time, they became too weak to ſhift their quarters to a more ſouthern climate, and thouſands were picked

up ſtarved to death in Devonſhire. Stares and Redwings ſuffered the ſame fate. If ſevere weather comes on more gradually, and food becomes ſcarce, few are ſeen with us after Chriſtmas; but on their return northward to breed appear again in ſmall flocks. This bird with us rooſts on the ground, and rarely perches for that purpoſe. Linnæus ſays it builds in high trees in Sweden, and frequents the places where junipers grow. It is alſo found, during the warmer months, in Ruſſia, Siberia, and Norway, where they undoubtedly breed.

FIG-EATER.

A bird ſo called by Willughby, who ſays it is found in Yorkſhire, where it is called Bean Bird. Mr. Pennant makes it the Pettychaps; but that cannot be if it makes its neſt under beams in old buildings, as it is ſaid to do. We have ſuſpected it to be no other than the Spotted Flycatcher, who frequently builds in ſuch ſituations. But we are left in great doubts about this bird. Certainly it is not the true Fig-eater found on the continent.

FINCH. A genus of birds, the characters of which are,

Bill perfectly conic, ſlender towards the point and ſharp.

Tongue truncated.

Toes, three forward, one backward.

FINCH-BEECH. Vide *Finch-chaf.*

FINCH-BRAMBLE. Vide *Brambling.*

FINCH-BUL.

Loxia pyrrhula. *Lin. Syſt.* i. p. 300. 4. *Gmel. Syſt.* ii. p. 846. *Raii Syn.* p. 86. A. *Will.* p. 130. t. 43. *Ind. Orn.* i. p. 387. 56. *Briſ.* iii. p. 308. 1.—*Ib.* 8vo. i. p. 395.

Bouvreuil. *Briſ.* iv. p. 372. t. 17.

Bulfinch. *Br. Zool.* i. N° 116 —*Ib.* fol. 106. t. U. f. 3. 4. *Arct. Zool.* ii. p. 353. A. *Albin.* i. t. 59. 60.—*Ib. Song Birds*, t. 15. *Will. Angl.* p. 247. t. 17. *Hayes, Br. Birds*, t. 37. *Lewin, Br. Birds*, ii. t. 70. *Lath. Syn.* ii. p. 143. 51. —*Ib. Sup.* p. 152. *Walc.* t. 209. *Pult. Cat. Dorſet.* p. 111.

PROVINCIAL.

Red-hoop. Tony-hoop. Alp. Nope.

This ſpecies of Groſbeak is ſo well known as to make it unneceſſary to be very particular in deſcription.

The bill is black, ſhort, and thick; irides duſky. The crown of the head is black; upper part of the neck and back fine cinereous grey; cheeks, breaſt, and belly, bright crimſon; vent white; coverts of the wings croſſed with a white line juſt above the quill-feathers, which laſt are duſky; rump white; tail black.

The female is very unlike the male in plumage,

xcept in the crown of the head, which is black; he whole bird beſides is of a dirty brown; rump white.

The Bullfinch is not gregarious; ſeldom more than a brood is ſeen together; is moſt commonly ſeen in pairs. It is not migratory, but frequents our woods and thickets all the year. In the ſpring it affects gardens and orchards, where it is very deſtructive among the buds of fruit-trees.

It makes a neſt the latter end of April or beginning of May, preferring the thickeſt places for that purpoſe, moſt frequently in a black or white thorn-buſh, either in woods or hedges. The neſt is compoſed of ſmall dry twigs, lined with fibrous roots.

The eggs are four or five in number, of a bluiſh white, ſpeckled, and ſtreaked with purple: rather larger than thoſe of a Linnet.

The young birds at firſt reſemble the female, except that it has no black on the head. This, as well as the crimſon on the breaſt of the male, does not appear till two months after they leave the neſt.

In confinement this bird not unuſually turns wholly black; ſuppoſed to be effected by being fed too much on hempſeed. Its native notes are few, but remarkably ſoft, and delivered ſo low as to eſcape a common obſerver; the call notes are ſimple, but more audible. When caged it becomes very docile, and may be taught a variety of tunes; for which reaſon, as well as for its beauty, it is frequently deprived of liberty.

FINCH-BUL-GREATEST. Vide *Groſbeak-pine.*

FINCH-CHAF.

Fringilla cælebs. *Lin. Syſt.* i. p. 318. 3. *Gmel. Syſt.* ii. p. 901. *Raii Syn.* p. 88. 16. A. *Will.* p. 186. t. 45. f. 4. *Ind. Orn.* i. p. 437. N°12. *Briſ.* iii. p. 148. 36.—*Ib.* 8vo. i. p. 437.

Le Pençon. *Buf.* iv. p. 109. f. 4.

Chaffinch. *Br. Zool.* i. N° 125.—*Ib.* fol. 108. t. 5. f. 2. 3. *Arct. Zool.* ii. p. 381. F. *Will. Angl.* p. 253. 45. *Albin.* i. t. 63.—*Ib. Song Birds*, t. p. 25. *Lewin, Br. Birds*, ii. t. 79. *Lath. Syn.* iii. p. 257. 10.—*Ib. Sup.* p. 165. *Walc.* ii. t. 217. *Pult. Cat. Dorſet.* p. 12.

PROVINCIAL.

Beechfinch. Horſefinch. Pink. Twink.

This bird is rather leſs than the Sparrow.

The bill is bluiſh; irides hazel; the forehead black; crown of the head, back part, and ſides of the neck, bluiſh aſh-colour; the cheeks, under ſide of the neck, and breaſt, dull pink; back cheſnut-brown; rump greeniſh; belly white, tinged with pink; the baſtard wing and coverts of the primary quills are black; thoſe of the ſecondary tipped with white; the ſmaller coverts black and greyiſh, on which is a ſpot of white; the quill-feathers duſky, ſlightly edged with greeniſh yellow on the outer webs, marked with white on both webs at the

aſe; tail duſky; the exterior feather is obliquely marked with white, taking in the whole of the outer web, the next is tipped with white; legs duſky.

The female is of a dull green above; the breaſt and belly of a brown or dirty white; the wings have the ſame markings as the male, but leſs brilliant.

This bird makes a moſt elegant neſt of green moſs, curiouſly ſtudded with lichen interwoven with wool, and lined with feathers and hair. It builds againſt the ſide of a tree, particularly in ivy, or in ſome forked branch of a buſh; but particularly in apple-trees overgrown with moſs and lichen, and, like many other birds, adapts the materials of its neſt to the ſurrounding colour; an inſtinct of no ſmall importance.

The eggs are four or five in number, larger than thoſe of the Goldfinch, of a dirty white, tinged with purple, marked with ſtreaks and ſpots of dark purple. Its notes are few, and ſcarce deſerve the name of ſong. Both ſexes have a monotonous call note, which ſeems to expreſs the word *twink*; from whence it is provincially called by that name.

In the ſummer theſe birds live chiefly upon inſects, with which they feed their young; in the winter become gregarious, and feed on ſeeds and grain. They remain with us the whole year, and flock with other hard-billed birds in the colder months; but the ſexes do not ſeparate, as they are known to do in Holland and other countries

Mr. White makes mention of flocks of females being ſeen in Hampſhire; theſe probably came from ſome more northern county.

FINCH-CHERRY. Vide *Groſbeak-haw.*

FINCH-COLD. Vide *Flycatcher-pied.*

FINCH-FALLOW. Vide *Wheatear.*

FINCH-GREAT-PIED-MOUNTAIN. Vide *Bunting-tawny.*

FINCH-GREEN. Vide *Groſbeak-green.*

FINCH-GOLD.

Fringilla Carduelis. *Lin. Syſt.* i. p. 318. 7. *Gmel. Syſt.* ii. p. 903. *Raii Syn.* p. 89. A. 1. *Will.* p. 189. t. 46. *Ind. Orn.* i. p. 449. 58. *Briſ.* iii. p. 53. 1—*Ib.* 8vo. i. p. 320.

Le Chardonneret. *Buf.* iv. p. 187. t. 10.

Goldfinch, *or* Thiſtlefinch. *Br. Zool.* ii. N° 124. —*Ib.* fol. 108. t. v. f. 1. *Arct. Zool.* ii. p. 283. H. *Will. Angl.* p. 246. t. 46. *Albin*, i. t. 64.—*Ib. Song Birds*, t. p. 18. *Hayes, Br. Birds*, t. 34. *Lewin, Br. Birds*, t. 81. *Lath. Syn.* iii. p. 281. *Walc. Syn.* ii. t. 219. *Pult. Cat. Dorſet.* p. 12. *Don. Br. Birds*, v. t. 103.

This beautiful bird is rather leſs than the Chaffinch. The bill is white, with a black point; irides duſky. The forehead and chin rich ſcarlet; top of the head black; cheeks white, bounded with black; hind part of the head white; breaſt pale tawny brown; the coverts of the wings black; quill-feathers duſky

black, barred acroſs with bright yellow; tips white; belly white; the tail feathers black; moſt of them marked with a white ſpot near their ends; legs whitiſh.

The female differs very little in plumage from the male; in general the ſmaller coverts of the wings are not ſo black.

Young birds are brown about the head for ſome time after they leave the neſt, and are by ſome called Grey Pates.

The Goldfinch is ſubject to variety on confinement; ſometimes wholly black; others black and white, or quite white. A variety is ſometimes taken by the birdcatchers with white ſpots under the throat: ſuch is termed a Cheverel.

It makes a very elegant neſt, formed externally of bents, moſs, and liverwort, wove together with wool; lined ſometimes with wool, or hair covered with thiſtle down, or willow cotton.

The eggs are four or five in number, of a bluiſh white, with a few ſmall ſpots, chiefly at the larger end.

It ſometimes builds in hedges, but moſt commonly in trees, eſpecially thoſe which are evergreen. The native ſong of this bird is not inelegant, but which is ſeldom pure in confinement. It readily breeds with the Canarybird; the production are mules, and generally termed Canary-goldfinch.

Their principal food is ſeed of various kinds, particularly the thiſtle and teaſel; is fond of plantain, chickweed, and hempſeed.

FINCH-HAW. Vide *Groſbeak-haw.*

FINCH-HORSE. Vide *Finch-chaf.*

FINCH-LESSER-MOUNTAIN. Vide *Bunting-mountain.*

FINCH-MOUNTAIN. Vide *Brambling.*

FINCH STORM. Vide *Petrel-ſtormy.*

FINCH-THISTLE. Vide *Finch-gold.*

FINCH-TREE. Vide *Sparrow-tree.*

FLUSHER. Vide *Butcherbird-red-backed.*

FLYCATCHER. A genus of birds, the characters of which are,

Bill beſet with briſtles, and flatted at the baſe; upper mandible notched at the end.

Toes divided to their origin.

FLYCATCHER-PIED.

Muſcicapa atricapilla. *Lyn. Syſt.* i. p. 326. 9. *Gmel. Syſt.* ii. p. 935. *Ind. Orn.* ii. p. 467. 1.

Œnanthe noſtra quarta. *Raii Syn.* p. 77. A. 5. *Will.* p. 170.

Rubetra anglicana. *Briſ.* iii. p. 436. 27.—*Ib.* 8vo. i. p. 271.

Traquet d'Angleterre. *Buf.* v. p. 222.

Cold Finch. *Will. Angl.* p. 236. *Edw.* t. 30.

Pied Flycatcher. *Br. Zool.* N°135.—*Ib.* fol. 103. t. S. f. 1. (mas.) *Arct. Zool.* ii. p. 391. B. —*Ib. Sup.* p. 64. *Lewin, Br. Birds,* iii. t. 88. *Lath. Syn.* iii. p. 324. 2. *Walc.* ii. t. 225. *Don. Br. Birds,* i. t. 22.

FLY

This ſpecies is about the ſize of a Linnet.

The bill is black; irides hazel. Forehead white; crown of the head, cheeks, taking in the eyes, and whole upper parts, black; the leſſer wing coverts and the greater coverts of the primores are duſky; the ſix firſt quill-feathers wholly duſky, the reſt white at the baſe, increaſing till the three laſt feathers are wholly white, except a ſpot of black near the point of the exterior web; the under parts of the bird, from chin to tail, are white; the tail is duſky black; the two exterior feathers white on their outer webs, and near the ſhaft on the inner webs, except at the point; the legs are black.

The female has no white on the forehead; the whole upper parts are duſky brown; the under parts dirty white; in the wings the white is leſs conſpicuous than in the male.

This bird is ſubject to ſome variety: ſome are ſaid to be mixed with grey on the upper parts, or ſpotted with white on the rump or upper tail coverts; others have white on the three exterior feathers of the tail; ſome have only the outer feather marked with white.

Young birds at firſt reſemble the female, and in their change of plumage the males have all the intermediate ſhades from brown to black. This has probably given riſe to an opinion that the male changes his plumage in the winter, and becomes like the female.

It is ſaid to be indigenous to England. It is, however,

however, very local, and by no means plentiful; but is ſpoken of as an inhabitant of Yorkſhire, Lancaſhire, and Derbyſhire.

We do not find the neſt has been diſcovered in this country. Dr. Latham ſays it builds in ſome hole of a tree, not very near the ground; and adds, that the neſt is only a few fibres, mixed with moſs, covering the bottom of the hole; and that the eggs are ſix in number.

With us it ſeems to affect wild and uncultivated tracts of furze, where there are no trees. In ſuch ſituations, therefore, it is probable it builds on the ground amongſt the high graſs and furze like the Whinchat and Stonechat, as it has all the actions and manners of thoſe birds.

FLYCATCHER-SPOTTED.

Muſcicapa Griſola. *Lin. Syſt.* i. p. 328. 20. *Gmel. Syſt.* ii. p. 949. *Raii Syn.* p. 81. 7. *Will.* p. 153. 171. 7.—*Ib. Angl.* p. 211. *Brif.* ii. p. 357. 1. t. 35. f. 3.—*Ib.* 8vo. i. p. 257. *Ind. Orn.* ii. p. 467. 1.

Stoporola Aldr? *Raii Syn.* p. 77. A. 1. *Will.* p. 159.—*Ib. Angl.* p. 217.

Le Gobemouche. *Buf.* iv. p. 517. t. 25. f. 2.

Cobweb. *Mort. North.* p. 426.

Spotted Flycatcher. *Br. Zool.* N° 134.—*Ib.* fol. 99. t. P. 2. f. 4. *Lewin, Br. Birds,* iii. t. 87. *Lath. Syn.* iii. p. 323. 1. *Walc.* ii. t. 224. *Pult. Cat. Dorſet.* p. 11.

PROVINCIAL.

FLY

PROVINCIAL.

Rafter. Bee-bird. Cherry-ſucker. Chanchider.

This ſpecies is about the ſize of a Titlark; length not quite ſix inches. The bill is duſky, and broad at the baſe; inſide of the mouth yellow; irides hazel.

The whole upper parts of the bird are brown; the head obſcurely ſpotted with duſky; breaſt and belly dull white; the ſhafts of the feathers on the breaſt duſky; ſides under the wings tinged with dull orange; legs ſhort and black.

This bird comes to us late in the ſpring, and departs in September. It frequents orchards and groves, will often make its neſt on the limb of ſome fruit-tree nailed againſt the wall, or in a hole; ſometimes in outbuildings, upon the end of a rafter or beam, and at other times againſt the body of a large tree upon the ſtump of a decayed branch. The neſt is formed of bents, moſs, and ſuch like materials, interwoven with ſpiders webs, and lined with feathers.

It lays four or five eggs, not much unlike thoſe of the Redbreaſt, but rather leſs, and the ruſt-coloured ſpots more diſtinct, and not ſo much confined to the larger end.

Its food ſeems to be entirely winged inſects; though we are told it is particularly fond of cherries. Perhaps they may frequent that and other

fruit-trees for the ſake of flies, which are attracted by the fruit.

As ſoon as the young birds leave the neſt they are led by the old ones to ſome neighbouring wood or grove where inſects abound, and where they may be ſeen darting in every direction in purſuit of flies, and frequently returning to the ſame ſtand.

Its note is a ſimple weak chirp, not frequently uſed till after the young are fled, ſo that the bird is not eaſily diſcovered, though it is by no means uncommon, and ſeems to be more generally diffuſed in England than the Redſtart, and many other of the ſummer migrants. We have obſerved it in ſeveral parts of Cornwall, where the laſt mentioned bird is rarely if ever ſeen; and it is more common than that bird in the weſt of Devonſhire.

FRENCH-PIE. Vide *Shrike-cinereous.*

FULMAR, or FULMER.

Procellaria glacialis. *Lin. Syſt.* i. p. 213. 3. *Gmel. Syſt.* ii. p. 562. *Ind. Orn.* ii. p. 823. 9.

Procellaria cinerea. *Brif.* vi. p. 143. 2. t. 12. f. 2. —*Ib.* 8vo. ii. p. 399.

Fulmar, *ou* Petrel-puffin gris-blanc. *Buf.* ix. p. 325. t. 22.

Wagellus Cornubienſium Mallemucke. *Raii Syn.* p. 130. A. 13.

Haffhert, *ſeu* Equus marinus. *Will.* p. 306. —*Ib. Angl.* p. 395.

Fulmar

Fulmar Petrel. *Br. Zool.* ii. N° 257.—*Ib.* fol. 145. t. M. 2. *Arct. Zool.* ii. N° 461. *Lath. Syn.* vi. p. 403. 9. *Lewin, Br. Birds*, vi. t. 217. *Walc. Syn.* ii. t. 89.

This ſpecies of Petrel is about the ſize of the Common Gull. The bill is yellow, very ſtrong, and much hooked at the end; the noſtrils are placed in a convex ſheath.

The head, neck, tail, and whole under ſide of the body, white; the back and coverts of the wings aſh-colour; quill-feathers duſky; the legs yellowiſh; back-toe wanting, inſtead of which is a ſort of ſpur.

The Fulmar is not frequently ſeen on our ſouthern coaſts; we never remember but one inſtance, and that was in South Wales.

It is not uncommon in ſome of the iſlands off the north of Scotland. At St. Kilda it breeds, and ſupplies the inhabitants with a vaſt quantity of oil, which is uſed for culinary as well as medical purpoſes.

Its food is entirely fiſh, particularly that of the moſt oily nature, by which means it is always charged with oil, and has the power of ejecting it with force from its bill; a mode of defence peculiar to the Petrel genus. Is ſaid to lay one large white egg the latter end of May.

FURZE-CHAT. Vide *Whinchat.*

G.

GADWALL.

Anas ſtrepera. *Lin. Syſt.* i. p. 200. 20. *Gmel. Syſt.* ii. p. 520. *Ind. Orn.* ii. p. 859. 69. *Briſ.* vi. p. 339. 8. t. 33. f. 1.—*Ib.* 8vo. ii. p. 452.

Anas platyrhynchos. *Raii Syn.* p. 145. A. 2. *Will.* p. 287.

Chipeau. *Buf.* ix. p. 187. t. 12.

Gadwall, *or* Grey. *Br. Zool.* ii. N° 288.—*Ib.* fol. 157. t. Q. 3. *Arct. Zool.* ii. p. 575. I. *Will. Angl.* p. 374. t. 72. *Lewin, Br. Birds,* vii. t. 258. *Lath. Syn.* vi. p. 515. 16. *Walc. Syn.* ii. t. 68. *Pult. Cat. Dorſet.* p. 21.

This ſpecies of Duck is about the ſize of a Widgeon; length nineteen inches.

The bill is two inches long, flat and black. Head, and moſt part of the neck, reddiſh brown, ſpotted with black; ſides of the head, throat, and fore part of the neck, rufous white, ſpotted with brown; the lower part of the neck, upper part of the back, breaſt, and ſcapulars, elegantly marked with black and white curved lines; lower part of the back duſky brown; rump, upper and under tail coverts, black; belly dirty white; ſides and thighs

thighs marked with light brown lines; the coverts on the ridge of the wing are pale reddiſh brown; beneath that of a purpliſh red; the loweſt order of a deep black; greater quills duſky; on the leſſer quills is a white patch; the tail aſh-coloured, edged with white; legs orange.

The female has the ſame markings in the wings, but not near ſo bright in colour; the breaſt reddiſh brown, ſpotted with black; the feathers on the neck, back, and rump, edged with pale reddiſh brown; the curved lines on the neck and breaſt of the male are wanting in this ſex.

This bird viſits us in winter, but does not appear to be a plentiful ſpecies. Suppoſed to bred in Sweden, and probably in Ruſſia and Siberia, being found in thoſe parts.

It is ſaid to be a great diver; feeds chiefly by night, concealing itſelf amongſt the reeds and ruſhes during the day. It makes a noiſe not unlike that of the Mallard, but louder.

GALLINULE. A genus of birds, the characters of which are,

Bill thick at the baſe, ſloping towards the point, and ſomewhat compreſſed ſideways.
Forehead more or leſs bare.
Body compreſſed.
Wings ſhort and concave.
Tail ſhort.
Toes divided to their origin.

GALLINULE-COMMON.

Fulica chloropus. *Lin. Syſt.* i. p. 258. 4. *Gmel. Syſt.* ii. p. 698.

Gallinula chloropus. *Ind. Orn.* ii. p. 770. 13.

Gallinula chloropus major. *Raii Syn.* p. 113. A. 1.—*Ib.* p. 190. 15. *Will.* p. 223. t. 58. *Briſ.* vi. p. 3. 1. t. 1.—*Ib.* 8vo. ii. p. 358.

Poule d'eau. *Buf.* viii. p. 171. t. 15.

Common Water-Hen, *or* More-Hen. *Albin.* ii. t. 71.—iii. t. 91. *Will. Angl.* p. 312. t. 58.

Common Gallinule. *Br. Zool.* ii. N° 217. t. 77. —*Ib.* fol. 131. t. L. 1. *Arct. Zool.* ii. N° 411. —*Sup.* p. 69. *Hayes, Br. Birds,* t. 30. *Lath. Syn.* v. p. 258. 12. *Lewin, Br. Birds,* v. t. 191. *Pult. Cat. Dorſet.* p. 16. *Walc. Syn.* ii. t. 169. *Don. Br. Birds,* v. t. 110.

PROVINCIAL.

Moor-hen. Marſh-hen. Cuddy. Moor-coot.

This birds weighs from fourteen to ſixteen ounces; length fourteen inches.

The bill is red towards the baſe, greeniſh at the point; irides reddiſh hazel.

The crown of the head and whole upper parts are of a deep olive-brown; the under parts cinereous; ridge of the wing and under tail coverts white; in ſome there are two or three white feathers at the baſe of the thigh; legs and toes duſky green; above the knee is a garter of red.

The

The female is leſs than the male, and wants the red on the bill. This mark, however, is moſt conſpicuous in the ſpring.

This is a very common ſpecies, found in moſt ſedgy and ſlow rivers, or ſtreams of water, and frequently in ponds abounding with weeds, where it can lay concealed.

Its food is principally inſects, ſeeds, and vegetable productions of various ſorts, for which it frequently quits the water. It runs faſt, and is equally expert in ſwimming and diving, although the feet do not ſeem calculated for the latter.

It is continually flirting up the tail as it runs or ſwims, at which time the white underneath is very conſpicuous.

The neſt is made of flags or ruſhes, which is placed near the ſurface of the water on ſome branch of a tree or buſh; ſometimes on the ſtump of an old willow. The eggs are nine or ten in number, but moſt commonly five or ſix, of a light yellowiſh brown, marked with ruſt-coloured ſpots. The young are hatched in about three weeks, and inſtantly take the water: they are covered with a black down.

When diſturbed will frequently perch. It flies badly, with its legs hanging down. In the act of diving the wings are uſed as fins, like moſt others of the aquatic tribe.

The breed of this bird is very much deſtroyed by ſummer floods in our rivers; by reaſon of their neſts

neſts being placed ſo near the water the eggs are continually ſwept away.

The young are frequently deſtroyed by fiſh of prey; and we have taken an old bird out of the ſtomach of a pike.

Is found in many parts of the European continent; as alſo in America, and in the Weſt Indies.

GALLINULE-CRAKE.

Rallus Crex. *Lin. Syſt.* i. p. 261. 1. *Gmel. Syſt.* ii. p. 711.

Gallinula Crex. *Ind. Orn.* ii. p. 766. 1.

Ortygometra. *Raii Syn.* p. 58. A. 8. *Will.* p. 122. t. 29. *Brif.* v. p. 159. 3.—*Ib.* 8vo. ii. p. 253.

Porphyrio rufeſcens. *Brif.* v. p. 533. 5.—*Ib.* 8vo. ii. p. 353. *Will.* p. 236.

Rale de Genet, *ou* Roi des Cailles. *Buf.* viii. p. 146. t. 12.

Land-Hen, Daker-Hen, *or* Rail. *Will. Angl.* p. 170. t. 29.—*Ib.* p. 316. *Albin.* i. t. 32.

Crake Gallinule. *Br. Zool.* ii. N° 216. t. 75.—*Ib.* fol. 131. *Arct. Zool.* ii. N° 412. *Lath. Syn.* v. p. 250. 1. *Walc. Syn.* ii. t. 170. *Lewin, Br. Birds,* v. t. 190. *Pult. Cat. Dorſet.* p. 15. *Don. Br. Birds,* v. t. 116.

PROVINCIAL.

Land-Rail. Corn-Crake, Crek, or Cracker. Bean-Crake. Corn-Drake.

The weight of this ſpecies is ſeven or eight ounces; length nine inches and a half.

The bill is of a light brown colour; irides hazel. The whole upper parts of the bird are of a rufous brown; the top of the head, back, and ſcapulars, marked with duſky black; the coverts of the wings and tail are of a fine bay; under parts pale yellowiſh brown, approaching to white on the belly; the ſides barred with dark and light rufous brown; the ſides of the head, over the eyes, inclining to aſh-colour; legs cinereous brown.

This bird has been ſuppoſed by ſome to be the Water-Rail in its ſummer plumage; but this is too obvious an error to dwell on; their characters and mode of life are entirely different. It never takes the water, but reſides in high graſs or corn, where it breeds, making a neſt of a few dry plants, and lays twelve or fourteen eggs, ſome ſay as many as twenty. Theſe are not much leſs than thoſe of a Partridge, but of a longer ſhape; of a dull white, marked with ruſt-coloured ſpots.

It is a migrative ſpecies, appearing with us about the latter end of April, and departing in October. On its firſt appearance, and till the female begins to ſit, the male is frequently heard to make a ſingular noiſe, much reſembling that of a comb when the finger is drawn along the teeth; and which has been uſed as a decoy.

It is much ſought after for the delicacy of its fleſh, but is a difficult bird to ſpring, depending on its

its legs for ſafety more than its wings. In the autumn, when the corn and graſs are cut, it takes to the thickets and brakes, and is with difficulty rouſed to take flight. It flies ſlow, and with its legs hanging down.

It is moſt plentiful in the northern parts of the kingdom, and in Ireland. On their firſt coming they are very lean, but before their departure become exceſſively fat. It is found in the Hebrides and Orknies; inhabits Germany, France, and Italy, and as far north as Norway.

Their food is chiefly worms, ſnails, and inſects, in defect of which ſeeds and various vegetables.

GALLINULE-SPOTTED.

Rallus Porzana. *Lin. Syſt.* i. p. 262. 3. *Gmel. Syſt.* ii. p. 712. *Ind. Orn.* ii. p. 772. 19.

Rallus aquaticus minor, ſive Maruetta. *Briſ.* v. p. 155. 2. t. 13. f. 1.—*Ib.* 8vo. ii. p. 252.

Gallinula ochra Geſneri. *Raii Syn.* p. 115. 7. *Will.* p. 236.—*Ib. Angl.* p. 316.

Petit Rale d'eau, *ou* le Marouette. *Buf.* viii. p. 157.

Spotted Gallinule. *Br. Zool.* ii. N° 215.—*Ib.* fol. 130. t. L.* 1. *Arct. Zool. Sup.* p. 69. *Lath. Syn.* v. p. 264. 18. *Walc. Syn.* ii. t. 172. *Lewin, Br. Birds,* v. t. 192. *Don. Br. Birds,* v. t. 122.

PROVINCIAL.

Spotted Water-Hen.

The weight of this elegant ſpecies is about four ounces; length nine inches.

The bill is yellowiſh green; the baſe red-orange; irides reddiſh hazel.

The crown of the head, back, and rump, olive-brown, ſtreaked with duſky; the two laſt ſpotted with white; the ſides of the head and neck daſhed with cinereous, and minutely ſpotted with white; the breaſt and wing coverts olive-brown, ſpotted with white; the larger coverts barred and ſtreaked with white; belly and vent dirty yellowiſh white; the ſides, above the thighs, barred with white; legs pale green.

This is rather a ſcarce bird with us, and is certainly a migrative ſpecies. We have met with it as early as the firſt of April, and as late as the middle of October, about which time it probably retires ſouth to paſs the winter. We have received it from Suſſex, Hampſhire, and Caermarthenſhire. It does not appear to be found far inland, but chiefly reſorts to the marſhes and borders of ſmall ſtreams well clothed with reeds and ruſhes, amongſt which it is concealed, and difficult to be rouſed.

It has all the manners and habits of the Common Gallinule; or Water-hen; is ſaid to make a neſt compoſed of ruſhes, placed amongſt reeds on the ſurface of the water, and to lay ſeven or eight white eggs. The young take the water as ſoon as hatched, and are ſaid to be quite black.

It is found in France and Italy in the ſpring, and

is

is there ſuppoſed to be migratory. Is found alſo in the ſouth of Ruſſia and weſtern parts of Iberia. In England it has not been obſerved farther north than Cumberland.

GAMBET.

Tringa gambetta. *Lin. Syſt.* i. p. 248. 3. *Gmel. Syſt.* ii. p. 671. *Ind. Orn.* ii. p 728, 9.

Gambetta. *Raii Syn.* p. 107. 2. *Will.* p. 222. —*Ib. Angl.* p. 300.

Totanus ruber. *Brif.* v. p. 192. 4.—*Ib.* 8vo. ii. p. 262.

Red-legged Horſeman. *Alb'n.* ii. t 68.

Gambet Sandpiper. *Br. Zool.* ii. N° 198. t. 70. *Arct. Zool.* ii. N° 394. *Lath. Syn.* v. p. 167. 9. *Lewin, B. Birds,* v. t. 168.

The ſpecies of Sandpiper, we are informed by Mr. Pennant, has been ſhot on the coaſt of Lincolnſhire. He deſcribes it to be the ſize of the Greenſhank; the head, back, and breaſt, cinereous brown, ſpotted with dull yellow; the coverts of the wings and ſcapulars cinereous, edged with yellow; the primores duſky, bordered with yellow; legs yellow.

Dr. Latham adds, that the bill is of a reddiſh colour, with a black tip; the irides yellowiſh green.

This bird ſeems to have been rarely met with in England. Said to be known in France, but is eſteemed there a rare bird. Is ſpoken of as inhabiting Scandinavia and Iceland.

GANNET.

Pelicanus baſſanus. *Lin. Syſt.* i. p. 217. 5. *Gmel. Syſt.* ii. p. 577. *Ind. Orn.* ii. p. 891. 26.

Anſer baſſanus. *Raii Syn.* p. 122. A. 2. *Will.* p. 247. t. 63.

Sula Hoieri. *Raii Syn.* p. 123. 5. *Will.* p. 249.

Sula baſſana. *Briſ.* vi. p. 503. 5. t. 44.—*Ib.* 8vo. ii. t. 492.

Fou de Baſſan. *Buf.* viii. p. 376.

Soland Gooſe. *Will. Angl.* p. 328. t. 63. *Albin.* i. t. 86.

Gannet. *Br. Zool.* ii. N° 293. t. 103.—*Ib.* fol. 160. t. L. *Arct. Zool.* ii. N° 510. *Lath. Syn.* vi. p. 608. 25. *Lewin, Br. Birds,* vii. t. 265. *Walc. Syn.* i. t. 94. *Pult. Cat. Dorſet.* p. 22.

This ſpecies of Pelican weighs about ſeven pounds; length full three feet; breadth above ſix. The bill is about ſix inches long, nearly ſtrait, except towards the point, where it declines a little; the edges of the mandibles are ſharp, and a little jagged irregularly; the colour dirty yellowiſh white; near the baſe of the upper mandible is a ſharp proceſs and ſuture, which enables the bird to move it a little in the act of ſwallowing large fiſh; along the ſides are two longitudinal grooves running the whole length, but no noſtrils. The irides are pale yellow; round the eyes the ſkin is bare, and of a blue colour; the chin is deſtitute of feathers, and of a

duſky colour, capable of great diſtention, forming a kind of pouch. The whole plumage is white, except the crown of the head, which is buff-colour, and the baſtard wing and greater quill-feathers, which are black; the tail feathers are twelve, ſtrong and pointed.

A variety is mentioned by Mr. Pennant to have been killed in Caernarvonſhire. *Vide* Sula.

The young birds, during the firſt year, and perhaps longer, are duſky, ſpeckled with white.

It lays one egg nearly the ſize of that of the Gooſe; the neſt is compoſed chiefly of ſea-weeds, and generally placed upon the moſt inacceſſible parts of the higheſt rocks.

The Gannet is found plentiful on ſome of the northern coaſts, particularly the Orknies and the Baſs Iſland in the Firth of Edinburgh; but we believe breeds no where further ſouth. In the autumn it leaves thoſe places, and purſues the herrings and pilchards round our coaſt, returning to their uſual haunts in ſpring. During their winter migration they are frequently found off the coaſt of Cornwall, and are ſeen in every part of the Britiſh and Iriſh channel, but generally keep far out at ſea.

It takes its prey by darting down upon it with great velocity from a conſiderable height.

The pouch with which it is furniſhed is of peculiar ſervice to carry food to its young.

GANNET-CORNISH. Vide *Gull-ſkua.*

GARGANEY.

Anas Querquedula. *Lin. Syſt.* i. p. 203. 32. *Gmel. Syſt.* ii. p. 531. *Raii Syn.* p. 148. 8. *Will.* p. 291. t. 74. *Ind. Orn.* ii. p. 872. 99. *Briſ.* vi. p. 427. 31. t. 39. f. 1. 2.—*Ib.* 8vo. ii. p. 473.

Phaſcas forte Geſnero D. Johnſon. *Raii Syn.* p. 147. A. 4. (fem.) *Will.* p. 289.

Sarcelle. *Buf.* ix. p. 260.

Garganey. *Br. Zool.* ii. N° 289. t. 101.—*Ib.* fol. 158. t. Q. 9. *Arct. Zool.* ii. p. 576. O. *Will. Angl.* p. 377. 7. t. 74. *Lath. Syn.* vi. p. 550. 87. *Lewin, Br. Birds,* t. 259. *Pult. Cat. Dorſet.* p. 21. *Walc. Syn.* i. t. 75. *Don. Br. Birds,* i. t. 21.

PROVINCIAL.

Pied Widgeon. Summer Teal.

This ſpecies of Duck is in ſize between the Widgeon and Teal; length about ſeventeen inches. The bill is dark lead-colour; irides light hazel.

The upper part of the head duſky brown, ſtreaked with duſky; over the eye is a broad white line, paſſing down the ſide of the neck; the cheeks and upper part of the neck purpliſh, marked with minute lines of white pointing downwards; the breaſt marked with ſemicircular lines of brown and black; chin black; belly dirty white, ſtreaked with duſky towards the vent; the ſides croſſed with numerous

ſmall

ſmall black undulated lines; coverts of the wings cinereous grey, the larger ones tipped with white; ſcapulars long and narrow, the upper ones ſtriped with black, white, and aſh-colour, the reſt cinereous grey; the ſpeculum on the ſecondary quill-feathers is green; the tail is duſky; legs lead-colour.

The female is brown above, ſtreaked with duſky; the white ſtreak behind the eye is very faint, and the green on the wing wanting.

This laſt diſtinguiſhes it from the female Teal, which in other reſpects it reſembles.

Whether this bird ever breeds with us is not aſcertained; we have received it from the decoys in Somerſetſhire in the month of April by the name of Summer Teal, and were informed it made its appearance on thoſe pools always about that time.

It is found with us in winter, as alſo in France at that ſeaſon. Has been obſerved in Sweden, Ruſſia, and Siberia, and even as far as Kamtſchatka, where it is ſaid to be plentiful, and moſt probably breed.

GID. Vide *Snipe-jack.*

GILLIHOWTER. Vide *Owl-white.*

GLEAD. Vide *Kite.*

GOATSUCKER. A genus of birds, the characters of which are,

Bill ſhort, hooked at the end; the baſe of the upper mandible beſet with a row of ſtiff briſtles; mouth wide.

Tongue

Tongue ſmall, entire.

Toes connected by a membrane as far as the firſt joint.

Legs ſhort; middle claw ſerrated.

Tail conſiſts of ten feathers.

GOATSUCKER-EUROPEAN.

Caprimulgus Europæus. *Lin. Syſt.* i. p. 346. 1. *Gmel. Syſt.* ii. p. 1027. *Raii Syn.* p. 26. A. 1. *Will.* p. 70. t. 14. *Ind. Orn.* ii. p. 584. 5. *Briſ.* ii. p. 470. 1. t. 44.—*Ib.* 8vo. i. p. 289.

L'Engoulerent. *Buf.* vi. p. 512.

Nocturnal Goatſucker. *Br. Zool.* ii. N° 173. t. 59.—*Ib.* fol. 97. t. R. 1.

European Goatſucker. *Arct. Zool.* ii. p. 437. A. *Will. Angl.* p. 107. *Albin.* i. t. 10. *Borlaſ. Cornw.* t. 24. f. 13. *White, Selb.* p. 62. 94. *Lath. Syn.* iv. p. 593. 5.—*Ib. Sup.* p. 194. *Lewin, Br. Birds,* iii. t. 127. *Pult. Cat. Dorſet.* p. 13. *Walc. Syn.* ii. t. 255. *Don. Br. Birds,* iii. t. 67.

PROVINCIAL.

Night-hawk. Dorr-hawk. Churn-owl. Goat-owl, Wheel-bird. Night-jarr.

This ſpecies weighs between two and three ounces; length full ten inches.

The bill is duſky and weak; mouth exceſſive wide; eyes very large; irides duſky.

The

The plumage is beautifully diverſified with black, brown, ferruginous, and white, ſprinkled and daſhed with cinereous; the under parts are ferruginous brown, with numerous undulated tranſverſe lines; the legs are very ſhort, ſcaly, and feathered below the knee; the claw of the middle toe ſerrated on the inner edge.

The male has a large oval ſpot of white on the inner web of the three firſt quill-feathers, and at the end of the two outmoſt tail-feathers.

The female wants the white ſpots on the wings and tail; in other reſpects is like the male.

With us this bird is only a ſummer viſitant, appearing about the middle of May, and departing again the latter end of September or beginning of October. It is the only ſpecies found in Europe out of nineteen or twenty enumerated by different authors.

It makes no neſt, but lays two eggs on the bare ground amongſt fern, heath, or long graſs, ſometimes in woods or furze; but at all times contiguous to woods, where it chiefly conceals itſelf by day. The eggs are larger than thoſe of a Blackbird, of an oblong oval, whitiſh, elegantly marbled with light brown and aſh-colour. It generally ſits on the ground, but if diſturbed frequently perches on the limb of a tree, moſt commonly lengthwiſe, not acroſs, as is uſual with moſt birds.

In the duſk of the evening it begins its flight in purſuit of the larger inſects, particularly the Scarabæus Melolontha and Solſtitialis, which riſe from their

their earthy abode about that time. Is alſo fond of the large-bodied moths; but few winged inſects eſcape its wide-extended mouth.

It makes a ſingular noiſe, like the ſound of a large ſpinning wheel, and which it is obſerved to utter perched, with the head lowermoſt; beſides which it emits a ſharp ſqueak, repeated as it flies.

The Goatſucker is moſt plentiful in the wild tracts of uncultivated land, interſperſed with rocks and wood. We have ſeen in Scotland eight or ten on wing together in the duſk of the evening, ſkimming over the ſurface of the ground in all directions, like the Swallow, in purſuit of inſects.

GODWIT-CAMBRIDGE.

Scolopax Cantabrigienſis. *Gmel. Syſt.* i. p. 668. *Ind. Orn.* ii. p. 721. 23.

Cambridge Godwit. *Br. Zool.* ii. N° 185. *Lath. Syn.* v. p. 146. 16. *Lewin, Br. Birds,* iv. p. 36.

We are informed this ſpecies of Snipe was ſhot near Cambridge. Mr. Pennant, to whom we are obliged for the deſcription of this bird, ſays it is larger than the Common Redſhank. The head, upper part of the neck, and the back, are of a cinereous brown; the leſſer coverts of the wings brown, edged with dull white, and barred with black; the primores duſky, whitiſh on their inner ſides; ſecondaries barred with duſky and white; under

under ſide of the neck and breaſt of a dirty white; belly and vent white; tail barred with cinereous and black; legs of an orange-red.

GODWIT-CINEREOUS.

Scolopax caneſcens. *Gmel. Syſt.* ii. p. 668. *Ind. Orn.* ii. p. 721. 22.

Cinereous Godwit. *Br. Zool.* ii. N° 180. t. 66. *Lath. Syn.* v. p. 145. 15.

This is another ſpecies of Snipe for which the public are indebted to Mr. Pennant, who ſays, the bill was two inches and a half long. The head, neck, and back, variegated with aſh-colour and white; the tail ſlightly barred with cinereous; the throat and breaſt white, the laſt marked with a few aſh-coloured ſpots; the legs long, ſlender, and aſh-coloured.

This was about the ſize of the Greenſhanks; approaches it nearly in colour; but the bill was ſo much thicker as to form a ſpecific diſtinction.

It was killed near Spalding.

GODWIT-COMMON.

Scolopax œgocephala. *Lin. Syſt.* i. p. 246. 16. *Gmel. Syſt.* ii. p. 667. *Ind. Orn.* ii. p. 719. 16.

Œgocephalus Bellonii. *Raii Syn.* p. 105. A. 4. *Will.* p. 215.—*Ib. Angl.* p. 294.

Limoſa rufa major. *Briſ.* v. p. 284. 6.—*Ib.* 8vo. ii. p. 282.

Le grande Barge rouſſe. *Buf.* vii. p. 505.

Godwit. *Lath. Syn.* v. p. 145. 14. A. *Albin.* ii. t. 70. *Br. Zool.* fol. t. B. B. *Wale. Syn.* ii. t. 141.

Scolopax leucophæa. *Ind. Orn.* ii. p. 719. 17?

Limoſa griſea major. *Briſ.* v. p. 272. 3. t. 24. f. 2.—*Ib.* 8vo. ii. p. 279.

Common Godwit. *Br. Zool.* ii. N° 179.—*Ib.* fol. 120. t. B. *Artt. Zool.* ii. N° 373. *Lath. Syn.* v. p. 144. 15.—*Ib. Sup.* p. 245.

Grey Godwit. *Lewin, Br. Birds,* iv. t. 161. *Don. Br. Birds,* iv. t. 75.

PROVINCIAL.

Yarwhelp, or Yarwip. Sea Woodcock.

This ſpecies of Snipe is ſubject to very conſiderable variety, both in ſize and plumage; and we conceive authors have erred in making more than one ſpecies out of theſe varieties, which is only a difference occaſioned by age or ſex. Dr. Latham had, in his *Synopſis*, given a variety which, afterward, in his *Index Ornithologicus*, he gave as a diſtinct ſpecies, under the title of Leucophæa; but we do not find any ſufficient character to make it diſtinct from the Œgocephala of Linnæus. In the many we have examined, the markings ſeem to run ſo much into one another that we cannot find a permanent diſtinction.

The weight of this bird is from ſeven to twelve ounces;

ounces; length fifteen or ſixteen inches. Bill from three inches and a quarter to upwards of four inches, a very little reflected, of a pale brown, duſky at the point; irides hazel. The head, neck, and upper parts, are of a light ruſty brown, in ſome inclining to aſh colour; the middle of each feather duſky; breaſt cinereous-brown; belly and under tail coverts white; in ſome the throat and rump are white; from the bill to the eye a whitiſh ſtroke; the prime quill-feathers are black; the ſhaft of the firſt white; tail white, barred more or leſs with duſky brown; legs long and duſky, ſometimes bluiſh grey.

In ſome the upper and under tail coverts are barred or ſpotted with brown or duſky; and other trifling varieties.

One now before us has the greater coverts of the wings ſo deeply margined with light grey-brown as to appear almoſt all whitiſh at a little diſtance; and the ſides of the body have a few long ſtreaks of brown.

Old male birds are ſaid to have ſome black lines on the chin, breaſt, and throat.

Theſe birds continue with us the whole year; in the ſpring reſort to the fens, where they breed; and are taken in nets with the Ruffs, by means of a ſtale or ſtuffed bird.

In the winter it is found on our ſhores, particularly at the mouths of large rivers and inlets, where

where the mud and ſand become bare at low water, where it feeds on inſects.

It is met with in various parts of the continent of Europe and Aſia, as well as in America.

GODWIT-LESSER. Vide *Snipe-jadreca.*

GODWIT-RED.

Scolopax Lapponica. *Lin. Syſt.* i. p. 246. 15. *Gmel. Syſt.* ii. p. 667. *Ind. Orn.* ii. p. 718. 15.

Limoſa ruſa. *Briſ.* v. p. 281. 5. t. 25. f. 1.—*Ib.* 8vo. ii. p. 281.

La Barge rouſſe. *Buf.* vii. p. 504.

Red Godwit. *Br. Zool.* ii. N° 181. t. 67.—*Ib.* fol. tab. add. *Arct. Zool.* ii. N° 372. *Edw.* t. 138. *Lath. Syn.* v. p. 142. 13. *Lewin, Br. Birds,* iv. t. 160. *Pult. Cat. Dorſet.* p. 14. *Walc. Syn.* ii. t. 140.

Small Curlew, *or* Red-breaſted Godwit. *Lin. Tranſ.* i. p. 128.

This ſpecies of Snipe is generally rather larger than the Common Godwit; weight about twelve ounces; length eighteen inches. The bill is full three inches and a half long, a trifle reflected, ſlender, duſky towards the point; the baſe yelloviſh fleſh-colour; irides hazel. The head, neck, breaſt, back, and ſcapulars, ferruginous; the top of the head is ſtreaked with duſky; the back and ſcapulars marked with large black ſpots or bars; from the

the bill to the eye a light-coloured ſtreak; the belly and under tail coverts white; the ſides under the wings barred with duſky; the ſmaller coverts of the wings on the ridge duſky, the next inclining to ferruginous; the larger ones cinereous-brown, light at the tips; greater quill-feathers black, ſhafts white; the outer webs ſlightly edged with white half way down; inner webs white at the baſe; the ſecondary quills duſky from their points half way; baſe white; thoſe next the body ferruginous, like the ſcapulars, barred with black; the rump and upper tail coverts white; the middle feathers of the tail duſky black, the reſt white half way from the baſe, ends black; legs near four inches long, and black; the thighs bare of feathers full an inch above the knee.

In ſome the breaſt is ſtreaked with black, others mottled rufous and white; and the upper tail coverts barred with rufous and brown.

The ſpecimen from which the above deſcription is taken was ſent to us by ſome unknown friend; it was killed early in the autumn.

It is a rare ſpecies in England, and not frequently met with; and we believe never in ſummer. One we ſaw in Cornwall had the breaſt and neck of a bright ferruginous.

It is ſaid to be plentiful in ſome parts of America; is found in numbers in the fens about Hudſon's Bay, where they breed and retire ſouthward.

GOLDEN-EYE.

Anas Clangula. *Lin. Syſt.* i. p. 201. 23. *Gmel. Syſt.* ii. p. 523. *Raii Syn.* p. 142. A. 8. *Will.* p. 282. 13. t. 73. *Ind. Orn.* ii. p. 867. 87. *Briſ.* vi. p. 416. 27. t. 37. f. 2.—*Ib.* 8vo. ii. p. 470.

Le Garrot. *Buf.* ix. p. 222.

Golden-eye. *Br. Zool.* ii. N° 276.—*Ib.* fol. 154. t. Addend. *Arct. Zool.* ii. N° 486. *Albin.* i. t. 96. *Will. Angl.* p. 368. t. 73. *Lath. Syn.* vi. p. 535. 76. *Lewin, Br. Birds,* vii. t. 255. *Walc. Syn.* i. t. 69.

Brown-headed Duck. *Lewin, Br. Birds,* vii. t. 256.

Morillon, *or* Grey-headed Duck. *Br. Zool.* ii. N° 277? *Arct. Zool.* ii. p. 573. F? *Lath. Syn.* vi. p. 537. 77? *Will. Angl.* p. 367.

Anas Glaucion. *Ind. Orn.* ii. p. 868. 88.

Willughby and Ray ſeem to have given the young, or female, of this ſpecies under the title of Smaller red-headed Duck.

Lewin conſiders his Brown-headed Duck as a diſtinct ſpecies, and ſtates the bill to be eſſentially different.

Nothing has tended ſo much to the diſcovery and aſcertaining the ſpecies of Ducks as the ſingular conformation in the *trachia*, or windpipe, of the males of ſome of this tribe of birds. That of the Golden-

Golden-eye is ſo very different from moſt others that it is an unerring mark of diſtinction.

We have examined many with the ruſty-brown head, as deſcribed by authors for a different ſpecies, ſome of which were males, and had the enlargement of the *trachia.*

A matured male Golden-eye weighs near two pounds; length between eighteen and nineteen inches. Bill black; irides fine bright yellow. Head and upper half of the neck black, gloſſed with green and violet, changeable as viewed in different lights; at the corner of the mouth is a large white ſpot; the lower part of the neck, the breaſt, and all beneath, are white; the back, rump, and upper tail coverts, black; ſcapulars black and white; the coverts of the wings are black, with a white patch on the leſſer, and another on the larger coverts; quill-feathers black, except ſeven of the ſecondaries, which are moſtly white; tail black; legs orange.

The female is conſiderably leſs. The bill is yellowiſh towards the point; the head in this, as well as in the male, is full of feathers, which makes it appear large, but inſtead of being black, it is, as well as the upper part of the neck, of a ruſty-brown; round the middle of the neck is a greyiſh ring; the lower part of the neck and breaſt mottled duſky and cinereous; back duſky, daſhed with cinereous; the coverts of the wings and quills like the male, but the black parts of a duſky colour; the fore part

of the legs and toes yellowiſh, the hind part and webs black.

The young male birds are like the female in plumage, but are larger.

We have taken no ſmall pains in diſſecting a great many of this ſpecies which appeared to be females, but proved males; and all ſuch poſſeſſed the ſwelling in the *trachia*, and the *labyrinth* of the Golden-eye.

It is probable the male of this ſpecies is many years arriving at full maturity, for it is rarely found with the full black head, and the white ſpot at the corner of the mouth.

The ſexes are readily diſcovered in their firſt feathers by paſſing the finger and thumb down the windpipe; the enlargement of the *trachia* is eaſily felt.

This bird, in its immature ſtate, is probably the Glaucion, or Morillon, of various authors. It viſits us in winter in ſmall flocks, and feeds chiefly in rivers contiguous to the ſea; ſometimes killed in freſh waters more remote; is a quick diver, and not eaſily ſhot, except on wing.

Is ſeldom caught in the decoys; but we have frequently bought them in Briſtol market, where they are indiſcriminately ſold for Widgeon.

It retires northward to breed; is found at that ſeaſon in Norway and Sweden; is known alſo in America.

GOOSANDER.

GOOSANDER.

Mergus Merganſer. *Lin. Syſt.* i. p. 208. 2. *Gmel. Syſt.* ii. p. 544. *Raii Syn.* p. 134. A. 1. *Will.* p. 253. t. 64. *Ind. Orn.* ii. p. 828. 1. *Brif.* vi. p. 231. t. 32.—*Ib.* 8vo. ii. p. 423.

Le Harle. *Buf.* viii. p. 267. t. 23.

Gooſander, *or* Merganſer. *Br. Zool.* ii. N° 260. t. 92. f. 1.—*Ib.* fol. 147. *Artt. Zool.* ii. N° 465.—*Ib. Sup.* p. 73. *Will. Angl.* p. 335. t. 64. *Lath. Syn.* vi. p. 418. 1. *Lewin, Br. Birds,* vi. t. 231. *Pult. Cat. Dorſet.* p. 19. *Walc. Syn.* i. t. 79. *Don. Br. Birds,* iii. t. 49.

This is the largeſt ſpecies of Merganſer; weight about four pounds; length two feet four inches. The bill three inches long, narrow, ſerrated, or toothed, on the edges of both mandibles; the tip of the upper hooked; colour red; irides the ſame.

The head and upper part of the neck gloſſy greeniſh black; the feathers on the crown and back of the head are long and looſe; the reſt of the neck, breaſt, and under parts, white; the ſides, above the thighs, undulated with duſky lines; the upper part of the back black; lower part of the back, rump, and tail coverts, browniſh aſh-colour; the leſſer wing coverts white; the reſt aſh-colour, with ſome white; the greater quill-feathers are black, with aſh-colour on the interior webs of ſome of the inner ones; the ſecondaries white, margined with greeniſh

black on the outer webs; the ſcapulars neareſt the body black, the others white; the tail conſiſts of eighteen aſh-coloured feathers, with duſky ſhafts; legs orange.

In ſome ſpecimens the breaſt is of a roſy buff-colour.

Mr. Pennant has given the Dundiver as the female of this ſpecies; but in the deſcription of that bird we have given our reaſons for continuing it as a diſtinct ſpecies.

The Gooſander ſometimes viſits our rivers and lakes in ſevere winters, but retires to the more northern latitudes to breed. Is ſaid to be found in the Hebrides in ſummer, and to continue in the Orknies the whole year. It is not uncommon on the continent of Europe and Aſia, but moſt plentiful towards the north; breeds in Greenland and Iceland; is alſo found in ſome parts of America.

Its food is fiſh, for which it dives with great celerity, and holds its ſlippery prey with great ſecurity by means of its toothed bill, ſo admirably adapted to the purpoſe.

GOOSANDER-RED-BREASTED. Vide *Merganſer red-breaſted.*

GOOSE. Several of the larger ſpecies of the Duck genus, commonly ſo called. The Common Gooſe is given under the title of Gooſe grey-leg.

GOOSE-BEAN.

Anas ſegetum. *Gmel. Syſt.* ii. p. 512. *Ind. Orn.* ii. p. 843. 28.

Bean Gooſe. *Br. Zool.* ii. N° 267. t. 94. f. 2. *Arct. Zool.* ii. N° 472. *Lath. Syn.* vi. p. 464. 23. *Lewin, Br. Birds*, vii. t. 239. *Pult. Cat. Dorſet.* p. 20. *Wale. Syn.* i. t. 65.

PROVINCIAL.

Small Grey Gooſe.

This ſpecies is leſs than the Common Gooſe; weight from five pounds to upwards of ſeven; length from two feet and a half to three feet. The bill is ſmall, compreſſed near the end, of a whitiſh fleſh-colour; the tip, or what is termed the nail, is black (the great characteriſtic mark of diſtinction); irides hazel. The head and neck cinereous brown; the whole under part, as far as the legs, the ſame, but lighter, darkeſt on the thighs; the forehead ſpeckled with white, behind which the feathers are duſky brown; the back is aſh-colour; the lower part of the belly, upper and under tail coverts, white; ſcapulars brown aſh-colour, edged with white; the greater quill-feathers are black; exterior webs grey; ſecondaries cinereous grey, margined with black on the outer webs; coverts grey, the larger ones tipped with white; legs dull orange. There appears to be ſome little variation in the plumage of theſe birds; in ſome the bill is of a dull browniſh red; the upper

part of the back, ſcapulars, and wing coverts, brown, daſhed with cinereous, and tipped with white; greater quills plain duſky black; ſecondaries grey, tipped and margined with white.

Theſe birds come to us early in the winter, ſometimes in large flocks, and are as often ſeen in the uplands as in the fens, reſorting to fields of green wheat, which at that ſeaſon they ſeem to prefer to any other food.

In the ſpring they retire northward to breed; many are ſaid to inhabit Lewis, one of the Hebrides, all the ſummer, and breed there.

It is frequently killed and ſold at market for the Common Wild Gooſe; and indeed is more frequent, but has long been confounded for that bird. The ſpecimen from which our deſcription is taken was killed in Wales; but we have ſeen it as far weſt as Devonſhire.

GOOSE-BERNACLE.

Anas erythropus. *Lin. Syſt.* i. p. 197. 11. *Gmel. Syſt.* ii. p. 512. *Ind. Orn.* ii. p. 842. 31.

Bernicla. *Raii Syn.* p. 137. A. 5. *Will.* p. 274. *Briſ.* vi. p. 300. 14.—*Ib.* 8vo. ii. p. 411.

La Bernache. *Buf.* ix. p. 93. t. 5.

Canada Gooſe. *Albin.* i. t. 92.

Bernacle, *or* Clakis. *Br. Zool.* ii. N° 269.—*Ib.* fol. 150. *Arct. Zool.* ii. N° 479. *Will. Angl.* p. 359. *Hayes, Br. Birds*, t. 24. *Lath. Syn.* vi. p. 466. 26. *Lewin, Br. Birds*, vii. t. 242. *Pult. Cat. Dorſet.* p. 20. *Walc. Syn.* i. t. 62.

The

The weight of this ſpecies is between four and five pounds; the length about two feet.

The bill is black and ſhort.

The forehead, chin, and cheeks, are white; from the bill to the eye a black line; the reſt of the head, neck, and upper part of the back, black; the breaſt and under parts white; thighs mottled duſky and white; black about the knee; the back, ſcapulars, and coverts of the wings, barred with black, white, and grey; upper tail coverts white; rump black; tail the ſame; legs duſky black.

Theſe birds frequent the north-weſt coaſts of England in winter in large flocks, and are plentiful in ſome parts of Ireland; but are rarely met with in the ſouthern parts of this country, except in very ſevere weather.

Is ſaid to retire north as early as February to breed, and are then found in Ruſſia, Lapland, Norway, and Iceland. Has been met with at Hudſon's Bay.

In the darker ages ſtrange accounts were given of the hiſtory of this bird, which was ſuppoſed to be produced from the ſhell called by Linnæus *Lepas anatifera*, which are frequently found in vaſt abundance adhering by a pedicle to logs of wood that have laid long in the ſea; from which it was given the name of Tree Gooſe. Strange as this may appear in this enlightened age, it was credited and handed down by various authors. Gerard, who

was one of theſe credulous perſons, has given a curious account of it, as may be ſeen in his Herbal.

GOOSE-BERNACLE-SMALLER. Vide *Gooſe-brent.*

GOOSE-BRAND. Vide *Gooſe-brent.*

GOOSE-BRENT.

Anas Bernicla. *Lin. Syſt.* i. p. 198. 13. *Gmel. Syſt.* ii. p. 513. *Ind. Orn.* ii. p. 844. 32.

Brenta. *Raii Syn.* p. 130. 8. *Will.* p. 275. t. 69. *Briſ.* vi. p. 304. 16. t. 31.—*Ib.* 8vo. ii. p. 442.

Le Cravant. *Buf.* ix. p. 87.

Brent, *or* Brand Gooſe. *Br. Zool* ii. N° 270.—*Ib.* fol. 151.—*Ib. Addend.* t. Q. *Arct. Zool.* ii. N° 478.—*Ib. Sup.* p. 75. *Albin.* i. t. 93. *Will. Angl.* p. 360. *Hayes, Br. Birds,* t. 25. *Lath. Syn.* vi. p. 467. 27. *Lewin, Br. Birds,* vii. t. 243. *Pult. Cat. Dorſet.* p. 20. *Walc. Syn.* i. t. 63.

PROVINCIAL.

Rat, or Road Gooſe. Clatter-gooſe.

This ſpecies is conſiderably leſs than the Bernacle. The ſpecimen before us weighed two pounds twelve ounces; length twenty-nine inches. Bill ſhort and black; irides duſky. The head, neck, and upper part of the breaſt, black; on each ſide the ſmall part of the neck is a patch of white, mixed

mixed with black; the back, ſcapulars, and wing coverts, are of a browniſh ſlate-colour, daſhed with cinereous; quills black; the under parts of the body ſlate-colour; the ſides a little barred with white; behind the thighs, the vent, and under tail coverts, white; the rump is black; upper tail coverts white; tail black; legs black.

The female is rather leſs; the plumage not ſo dark, and the feathers of the body above and beneath margined with grey, greatly ſo on the wing coverts; the lower part of the rump, as well as the tail coverts, are white.

Young birds are ſaid to want the white mark on the neck.

Theſe birds appear on our coaſts in winter, particularly in the weſt of England, but are moſt plentiful in Ireland, where they are taken in nets placed acroſs the rivers, and are eſteemed good eating; are in greateſt abundance in thoſe rivers that empty themſelves into the northern part of the Iriſh channel. Sometimes appear in vaſt flocks on the coaſts of Picardy, even to a peſt, deſtroying all the corn near the ſea. In the year 1740 theſe birds, we are told, appeared in ſuch prodigious numbers on that coaſt that the inhabitants knocked them down with clubs.

It is ſaid to be eaſily tamed.

They breed far north, and return ſouthward in autumn. In Shetland are called Horra Geeſe. Are common at Hudſon's Bay, and probably breed there.

there. We are informed ſome of theſe birds breed in Ireland: however that may be, the greater part retire more northward for that purpoſe.

GOOSE-CANADA. Vide *Gooſe-bernacle.*

But this is not the true Canada Gooſe. That bird has not been found in this country in a wild ſtate, though it is frequently domeſticated with us, and will breed with the common ſpecies.

GOOSE-EMBER. Vide *Diver-imber.*

GOOSE-FEN. Vide *Gooſe grey-lag.*

GOOSE-GREY-LAG.

Anas Anſer (ferus). *Lin. Syſt.* i. p. 197. 9. *Gmel. Syſt.* ii. p. 510. *Raii Syn.* p. 136. A. 4.—138. A. 3. *Will.* p. 274. 2. t. 69. *Ind. Orn.* ii. p. 841. 26.

Anſer ſylveſtris. *Briſ.* vi. p. 265. 2.—*Ib.* 8vo. ii. p. 432.

Oie ſauvage. *Buf.* ix. p. 30. t. 2.

Wild Gooſe. *Will. Angl.* p. 358. *Albin.* i. t. 90.

Grey-lag Gooſe. *Br. Zool.* ii. N° 266.—*Ib.* fol. 150. *Arct. Zool.* ii. N° 473. *Lath. Syn.* vi. p. 459. 31. *Lewin, Br. Birds,* vii. t. 238. *Pult. Cat. Dorſet.* p. 20. *Walc. Syn.* i. t. 61.

PROVINCIAL.

Fen Gooſe.

This is the origin of our common domeſtic Gooſe. It weighs eight or nine pounds; length about thirty-three inches.

The

The bill is large and elevated, of a fleſh-colour; the nail white; irides greyiſh.

The head and neck aſh-colour, mixed with light ruſt-colour; back duſky grey; the breaſt and belly whitiſh, clouded with aſh-colour; ſcapulars grey, edged with white; the wing-coverts are compoſed of different ſhades of cinereous, margins of the feathers lighteſt, the leſſer coverts darkeſt; quill-feathers more or leſs black at their ends; the upper and under tail coverts pure white; the tail feathers are duſky, tipped with white, the outer ones almoſt entirely white; legs fleſh-coloured; claws black.

Theſe birds are ſaid to reſide in the Lincolnſhire fens the whole year, where they breed, and the young frequently taken and become tame. It lays eight or nine eggs of a dirty white. The greater part of the large flocks ſeen with us in the winter doubtleſs retire northward to breed. Is frequently killed upon the downs in the ſouth of England, feeding on green wheat. We remember one being ſhot in the wing by a farmer in the neighbourhood of the Wiltſhire downs, was kept alive many years, but would never aſſociate with the tame ones.

This and moſt of the other ſpecies are indiſcriminately called Wild Geeſe.

GOOSE-LAUGHING. Vide *Gooſe white-fronted.*

GOOSE-RAT, or ROAD. Vide *Gooſe-brent.*

GOOSE-RED-BREASTED.

Anas ruficollis. *Gmel. Syſt.* ii. p. 511. *Ind. Orn.* ii. p. 841. 23.

Red-breaſted Gooſe. *Arct. Zool.* ii. p. 571. C. *Lath. Syn.* vi. p. 455. 17. *Lewin, Br. Birds,* vii. t. 241.

This ſpecies is leſs than the Common Gooſe; weight about three pounds; length twenty-one inches. Bill brown; nail black; irides yellow-brown. The fore part of the head and crown black, paſſing backwards in a narrow ſtripe to the back; forehead and cheeks ſprinkled with white; between the bill and eye an oval white ſpot, above which is a black line; chin and throat black; behind the eye is white, paſſing down the neck on each ſide; the middle of this white is rufous; the reſt of the neck deep rufous; on the breaſt is a band of black, and another of white; belly white; ſides ſtriped with black; back and wings black; greater wing coverts tipped with grey; upper and under tail coverts white; legs black.

This is a rare ſpecies in England; two or three inſtances only on record. One, Dr. Latham informs us, was ſhot near London in the ſevere froſt of 1766; and another was taken alive in Yorkſhire about the ſame time, ſoon became tame, and was kept amongſt other Ducks in a pond.

This beautiful ſpecies is ſaid to breed in the northern parts of Ruſſia.

GOOSE-

GOOSE-SMALL-GREY. Vide *Gooſe-bean.*

GOOSE-SOLAND. Vide *Gannet.*

GOOSE-WHITE-FRONTED.

Anas albifrons. *Gmel. Syſt.* ii. p. 509. *Ind. Orn.* ii. p. 842. 27.

Anas erythropus. *Faun. Suec.* N° 116. (fem.)

Anſer ſeptentrionalis ſylveſter. *Briſ.* vi. p. 269. 3. —*Ib.* 8vo. ii. p. 433.

L'Oie rieuſe. *Buf.* ix. p. 81.

Laughing Gooſe. *Edw.* t. 153.

White-fronted Gooſe. *Br. Zool.* ii. N° 268. t. 94. 1.—*Ib.* fol. 150. *Arct. Zool.* ii. N° 476. *Lath. Syn.* vi. p. 463. 22. *Lewin, Br. Birds,* vii. t. 240. *Pult. Cat. Dorſet.* p. 20. *Walc. Syn.* i. t. 64. *Don. Br. Birds,* v. t. 102.

This is rather leſs than the Common Gooſe; is ſaid to weigh ſometimes as much as ſeven pounds. The bird now before us is of the general ſize, and was only four pounds four ounces. (It ſhould, however, be remembered, that great allowance is to be made in the weight of birds, as it depends on their condition, and whether the ſtomach is full or empty.) The length twenty-nine inches. The bill is fleſh-coloured, with an orange ſpot on the top at the baſe between the noſtrils; the nail white; irides duſky. The feathers round the baſe of the bill are white, running ſome way on the forehead, bounded by duſky brown; the head, neck, and upper parts, dark brown; the upper part of the back, ſcapulars, and

Greater creſted and horned Ducker. *Will. Angl.* p. 340. 5. t. 61. f. 1. *Albin.* i. t. 81. *Plot's Hiſt. Staff.* p. 229. t. 22.

Creſted Grebe. *Br. Zool.* ii. N° 223.—*Ib.* fol. 132. t. K. *Artt. Zool.* ii. p. 498. A. *Lath. Syn.* v. p. 281. 1. *Lewin, Br. Birds,* v. t. 106. *Walc. Syn.* i. t. 102. *Don. Br. Birds,* iii. t. 68.

Colymbus cinereus-major. *Raii Syn.* p. 124. A. 1. *Will.* p. 357. *Albin.* ii. t. 75.

Colymbus criſtatus. *Briſ.* vi. p. 38. 2. t. 4.—*Ib.* 8vo. ii. p. 368.

Le Grebe huppé. *Briſ.* viii. p. 233.

Grey or aſh-coloured Loon. *Will. Angl.* p. 340. 4. t. 61. f. 4.

Colymbus urinator. *Lin. Syſt.* i. p. 223. 9. *Gmel. Syſt.* ii. p. 593.

Colymbus. *Briſ.* vi. p. 34. 1. t. 3. f. 1.—*Ib.* 8vo. ii. p. 368.

Colymbus major Aldror. *Raii Syn.* p. 125. 6. *Will.* p. 256. t. 51.

Le Grebe. *Buf.* viii. p. 227.

Greater Loon, *or* Arſefoot. *Will. Angl.* p. 339. t. 51. *Edw.* t. 360. f. 2.

Tippet Grebe. *Br. Zool.* ii. N° 222. t. 78.—*Ib.* fol. 133. *Lath. Syn.* v. p. 283. 2.

PROVINCIAL.

Cargooſe, Gaunt.

The variety of ſpecies authors appear to have made from this bird has obliged us to be more than uſually diffuſe in the ſynonyms; all of which we conſider to be mere variety occaſioned by age, ſex, and ſeaſon.

A full-grown male Creſted Grebe weighs between two and three pounds; length about two feet. The bill is two inches and three quarters long, duſky brown along the ridge of the upper mandible and at the point; the reſt reddiſh fleſh-colour; irides and lore crimſon. The head is much enlarged by a creſt of a duſky colour, ſtanding up on each ſide; the cheeks and throat are ſurrounded by long feathers of a ferruginous colour; from the bill to the eye is a black line, above which is a white one; the chin is white; the hind part of the neck, and the upper part of the body and wings, duſky brown; the under part of the neck, breaſt, and all beneath, beautiful gloſſy white; the primary quill-feathers duſky, ſome of the inner ones tipped with white, the reſt are nearly all white, which, when the wing is cloſed, makes an oblique bar of that colour acroſs it; legs duſky on the outſide; ſome wholly duſky green.

In a male bird now before us the long feathers on the ſides of the head and throat are moſtly duſky, with a mixture of ferruginous; the ſides under the wings, and the thighs, duſky brown; two or three of the outer ſcapulars, and the ridge of the wing, white; the lore is duſky green.

This bird is probably of the ſecond year, and not arrived at maturity.

The female now before us did not weigh two pounds; the length twenty inches to the end of the rump feathers; bill nearly the ſame, but lighter fleſh-coloured; irides rufous brown; the feathers about the head ſcarcely elongated; the upper part of the head, back of the neck, and whole upper parts, duſky, daſhed with aſh-colour; on the crown of the head the feathers are a little lengthened; the lore is brown, beneath which a ſtroke of ſmall brown feathers reach from the mouth to the eye; the cheeks are white, with a few black ſpots near the ſides of the throat; the under parts are wholly of a fine ſatin white from chin to vent; the ſhoulders, and ſmalleſt coverts of the wings, white; the eleven firſt quill-feathers are duſky, the four laſt of them tipped with white; the reſt are white, but a few of them have a duſky ſtreak down the ſhaft; the legs are duſky without, inſide pale fleſh-colour; inner edge yellow; feet duſky beneath, pale fleſh-colour above; edges of the fins yellow; nails bluiſh.

Theſe and other varieties frequently occur; and it is probable that even in maturity two are ſcarcely alike in plumage, and that the full feathers of the head are not perfected till the third year. It muſt alſo be obſerved, that this, as well as many other birds, vary in the colour of thoſe parts deſtitute of feathers, ſuch as the bill, lore, and legs; theſe change with the ſeaſon, and are brighteſt in the ſpring.

There

There can be no doubt that the Tippet Grebe is the female or young of this ſpecies.

Dr. Latham ſeems to have been inclined to this opinion in his *Synopſis*; and in his *Index Ornithologicus* has judiciouſly brought them together.

That author informs us that a large flock of the Creſted Grebes appeared on the ſhores of the river Thames, many of which came under his inſpection; amongſt which he found the greateſt variety about the head, from being perfectly without a creſt, to the moſt complete one with all the intermediate ſtages.

This bird is indigenous to England; breeds in the meres of Shropſhire and Cheſhire, and in the fens of Lincolnſhire.

The neſt is large, compoſed of a variety of aquatic plants; it is not attached to any thing, but floats amongſt the reeds and flags, penetrated by the water.

The female lays four white eggs, about the ſize of that of a Pigeon.

Their principal food is fiſh, in purſuit of which they dive admirably. On the leaſt appearance of danger they plunge under water, depending very little on their wings for ſafety. Is very rarely ſeen on land, and at this ſeaſon ſeldom fly farther than from one ſide of the pool to the other. Mr. Pennant ſays it will carry its young upon its back, or under its wing, when they are tired, and feeds them with ſmall eels.

In the winter theſe birds viſit our coaſts and large rivers, eſpecially in hard weather, when the ſtanding waters are frozen.

GREBE-DUSKY.

Colymbus obſcurus. *Gmel. Syſt.* ii. p. 592.
Colymbus minor. *Brit.* vi. p. 567.—*Ib.* 8vo. ii. p. 373.
Podiceps obſcurus. *Ind. Orn.* ii. p. 782. 4.
Le petit Grebe. *Buf.* viii. p. 232.
Black and white Dobchick. *Edw.* t. 96. f. 1.
Duſky Grebe. *Br. Zool.* ii. N° 225. t. 78. 1.—*Ib.* fol. 133. t. K. 1. *Arct. Zool.* ii. N° 420. *Lath. Syn.* v. p. 286. 5. *Lewin, Br. Birds,* v. t. 193. *Walc. Syn.* i. t. 106. *Don. Br. Birds,* ii. t. 44.

This ſpecies is rather larger than the Little Grebe; length eleven inches.

Bill rather more than an inch long, and black; edges of the mandible in ſome red, others fleſh-coloured; irides and lore red; ſometimes the latter is fleſh-colour or brown. The upper part of the head, neck, and body, duſky; in ſome the cheeks are ferruginous, and forehead white; the under parts, from chin to vent, white; the breaſt and belly very gloſſy; in ſome the white on the throat almoſt encompaſſes the neck; the ridge of the wing is white; primary quills duſky; ſecondaries tipped with white; thighs mixed with duſky. In

ſome

ſome the under parts of the neck and ſides of the body are duſky grey, or aſh-coloured; others ſpotted duſky between the legs; legs duſky green on the outſide, fleſh-coloured within; toes fleſh-coloured; fins duſky.

Inhabits the fens of Lincolnſhire, where it breeds. Makes a neſt in the ſame manner as the Creſted Grebe; the eggs are alſo white.

Is found in the winter in our inlets on the coaſt, particularly in Devonſhire, where it is by no means uncommon.

GREBE-EARED.

Colymbus auritus. *Lin. Syſt.* i. p. 222. 8. *Gmel. Syſt.* ii. p. 590.

Podiceps auritus. *Ind. Orn.* ii. p. 781. 3.

Colymbus auritus. *Brif.* vi. p. 54. 6.—*Ib.* 8vo. ii. p. 372.

Le petit Grebe huppé. *Buf.* viii. p. 235.

Eared Dobchick. *Edw.* t. 96. f. 2.

Eared Grebe. *Br. Zool.* ii. N° 224. t. 79.—*Ib.* fol. 133. *Arct. Zool.* ii. p. 499. B. *Lath. Syn.* v. p. 285. 4. *Walc. Syn.* i. t. 104. *Lewin, Br. Birds,* v. t. 107. *Don. Br. Birds,* ii. t. 29.

Colymbus criſtatus minor. *Brif.* vi. p. 42. 3. t. 3. f. 2.—*Ib.* 8vo. ii. p. 369.

This ſpecies is larger than the Duſky Grebe; length about one foot. Bill about an inch long,

duſky; a little turned up at the point; reddiſh at the baſe; irides and lore crimſon. The head is well clothed with feathers of a duſky black; the neck and upper parts of the body duſky; behind the eyes, on each ſide, is a tuft of feathers falling backwards of a ferruginous colour; the breaſt and under parts ſilvery white; the feathers on the ſides ferruginous; legs black.

This, like moſt of the genus, is ſubject to ſome variety. In ſome the head and tufts behind the eyes are brown; ſides of the head white; neck white on the fore part, marked with ferruginous ſpots; ſides of the body marked the ſame; the back of the neck and upper parts of the body brown; ſome of the ſecondary quill-feathers and wing coverts next the body are white.

The female, in general, is not ſo full of feathers on the head.

Inhabits the fens of Lincolnſhire, where it breeds; lays four or five white eggs on a floating neſt. Feeds on fiſh and water plants. Is ſometimes found in winter in the inlets and rivers on the coaſt.

Found in the north of Europe, in Iceland, and Siberia.

GREBE-HORNED. Vide *Grebe-ſclavonian.*

GREBE-LITTLE.

Colymbus criſtatus. *Lin. Syſt.* i. p. 223. *γ.*

Colymbus minor. *Gmel. Syſt.* ii. p. 591.

——— Fluviatilis. *Briſ.* vi. p. 59. 9.—*Ib.* 8vo. ii. p. 374.

Podiceps minor. *Ind. Orn.* ii. p. 784. 9. *Raii Syn.* p. 125. A. 3. *Will.* p. 258. t. 61.

Yacapitzahoac. *Raii Syn.* p. 177.

Le Grebe de la rivier ou le Caſtagneux. *Buf.* viii. p. 244. t. 20.

Little Grebe. *Br. Zool.* ii. N° 226.—*Ib.* fol. 134. t. F. *Will. Angl.* p. 340. t. 61. *Lath. Syn.* v. p. 289. 10. *Lewin, Br. Birds,* v. t. 200.

Didapper. *Walc. Syn.* i. t. 105. *Don. Br. Birds,* iii. t. 56.

PROVINCIAL.

Dipper. Dobchick, or Dabchick. Small Doucker. Loon. Arſefoot.

The weight of this ſpecies is between ſix and ſeven ounces; length ten inches.

Bill ſcarce an inch long, brown; irides reddiſh hazel. The whole upper parts are of a ruſty brown; pale on the rump; cheeks light ferruginous; the fore part of the neck grey; in ſome the cheeks are of the ſame colour; breaſt and belly gloſſy white, mottled with aſh-colour and light ferruginous; in ſome the ferruginous is wanting, eſpecially in the females; but both ſexes are ſubject to vary in plumage according to age; legs dark dull green.

This is the leaſt and moſt plentiful ſpecies; is common in moſt lakes, ſlow rivers, ſmall ſtreams, and even fiſh-ponds.

It flies with ease, but dives at the least alarm, and will remain under water amongst the roots of other aquatic plants with only its bill above for respiration.

Its nest is formed, like the rest of the Grebes, of a prodigious quantity of flags or other water plants, but is generally fastened to the reeds or flags, in order to prevent it being carried away by the current.

The eggs are five or six in number, of a dirty white; the shape oblong, less than those of a Pigeon. These are generally covered with weeds, so that when the bird quits her nest suddenly they are not exposed to view. Notwithstanding this precaution they are frequently destroyed by the water rat.

Mr. Pennant supposes that the quantity of vegetables collected to form the nest ferments and gives warmth to the eggs; but we never could discover the least warmth in the nest.

In large rivers these birds are frequently devoured by pike and trout while they are diving in pursuit of small fish. We once took from the stomach of the last a water rat weighing fifteen ounces; the trout weighed only four pounds. In the spring the males are very active in pursuit of the other sex, and then frequently fly along the surface of the water to a small distance: at this season it also emits a shrill chattering noise. After the breeding season it is frequently seen in some of our

inlets

inlets of the ſea, where it is ſaid to feed on ſhrimps.

This bird is not uncommon in moſt parts of the old continent, and ſome parts of America, particularly at Hudſon's Bay.

GREBE RED-NECKED.

Colymbus rubricolis. *Gmel. Syſt.* ii. p. 529.
——— ſubcriſtatus. *Gmel. Syſt.* ii. p. 590.
Podiceps rubricolis. *Ind. Orn.* ii. p. 783. 6.
Le Grebe à joues griſes, Jougriſe. *Buf.* viii. p. 241.
Red-necked Grebe. *Arct. Zool.* ii. p. 499. C. —*Ib. Sup.* p. 69. *Lath. Syn.* v. p. 288. 7.—*Ib. Sup.* p. 260. t. 118. *Lewin, Br. Birds,* v. t. 199. *Walc. Syn.* i. t. 103. *Don. Br. Birds,* i. t. 6.

Length eighteen inches; bill nearly two inches long; ſides of the baſe of both mandibles, for three quarters of an inch, of a fine orange-yellow, the reſt black; lore brown or blackiſh; irides fine orange red; the crown, and ſides of the head above the eyes, nearly black, and the feathers a little elongated; the hind part of the neck, the back, and wings, dark brown; ſix of the middle ſecondaries white, a little mottled with duſky at the tips; the two or three next outward ones more or leſs white near the tips and inner webs; the chin, ſides under the eyes, and fore part of the neck for above

an

an inch, pale aſh-colour; the reſt of the neck ferruginous-cheſnut, mottled with duſky; from thence to the vent white, like ſatin, mottled on the ſides with duſky irregular ſpots; legs black.

The deſcription of this rare ſpecies we have borrowed from Dr. Latham's *Supplement* to his *Synopſis*, who ſays two of theſe birds were taken alive in Eaſt Kent in April 1786. He alſo mentions two other ſpecimens, one killed at Teignmouth in January, the other at Sandwich in October. The firſt of theſe had the colours of the head and neck much blended, and the ferruginous of the neck only juſt breaking forth. The other weighed nineteen ounces and a half; length twenty-one inches and a half; the bill yellow at the baſe; duſky olive towards the tip; lore duſky; irides pale brown; head quite ſmooth; the ferruginous colour of the neck much blended with duſky; the white on the under parts greatly mottled with the ſame; legs without duſky, within greeniſh yellow. Another variety is mentioned to have a band of white acroſs the lower part of the neck.

This bird is ſuppoſed to inhabit Denmark or Norway.

GREBE-SCLAVONIAN.

Podiceps cornutus. *Ind. Orn.* ii. p. 783. 5. β.

Colymbus, ſive Podiceps minor. *Raii Syn.* p. 190. 14.

Colymbus

Colymbus cornutus minor. *Briſ.* vi. p. 50. 5.
—*Ib.* 8vo. ii. p. 371.
Le petit Grebe cornu. *Buf.* viii. p. 237.
Horned Grebe. *Lath. Syn.* v. p. 288. 6. var. A.

This bird is larger than the Duſky Grebe; length thirteen inches and a half to the end of the rump-feathers; breadth twenty-two inches.

The bill is near an inch long, duſky, lighter at the baſe of the under mandible, and inclining to pink, the very tip light horn-colour; lore crimſon; irides the ſame, but round the pupil a circle of white, and the exterior edge of the iris is ſhaded to nearly white. The head is greatly enlarged by the feathers; thoſe on the top are black, tinged with dark green; the cheeks and throat the ſame; the feathers very long, forming a ſort of ruff; from the baſe of the upper mandible originates a broad bar of dull orange yellow that paſſes through the eye to the hind-head, growing gradually broader; theſe form a tuft on each ſide, and are ſomewhat erectable appearing like ears; the forehead duſky ferruginous; the back of the neck and upper part of the back dark brown, daſhed with ferruginous; the back, ſcapulars, and rump, duſky, faintly edged with cinereous; the wing coverts and twelve firſt quill-feathers brown; the thirteenth white on the inner web; the eleven next all white, except the laſt, which is brown on the outer web; the chin is black, a little mottled with white; the under part

of

of the neck and upper breaſt running far behind and down under the wings, bright ferruginous cheſnut; the reſt of the under parts gloſſy ſatin white; the back part of the thighs ferruginous brown; legs duſky on the outſide, pale on the inſide; toes pale down their middle, duſky at the edges.

This bird, which was reſcued from the hands of a fiſherman as he was juſt going to pick it, was killed near Truro in Cornwall on the fourth of May 1796, and preſented to me by a friend. It was a male bird, and is now in my muſeum. It had no *labyrinth*, nor any thing uncommon in the *trachia*.

We are very much inclined to believe, with Dr. Latham, that it is a variety of the

Horned Grebe. *Synopſis* v. p. 287. 6. t. 91.
The eared or horned Dobchick. *Edw.* t. 145.
The Colymbus cornutus. *Lin. Syſt.* i. p. 591.

Dr. Latham ſays it is found in Sclavonia. Mr. Pennant ſays, in his *Arct. Zool.* ii. N° 417, that the Horned Grebe is found at Hudſon's Bay in June, and breeds in freſh water; appears at New York in the ſpring, where it is called the Water Witch, from its vaſt quickneſs in diving.

We cannot, however, help expreſſing our doubts concerning theſe birds. If we conſider that the Grebes are all ſubject to great variety in plumage, occaſioned by age; and if we compare the various deſcriptions given by authors of the Horned, or

Sclavonian Grebe; we ſhall not find any very eſſential difference from the Eared Grebe; at leaſt not more than there is in individual ſubjects of that ſpecies. The ſize is nearly the ſame; the moſt material diſtinction is in the colour of the neck and breaſt. We muſt, however, leave this matter to thoſe who have more frequent opportunities of examining the Eared Grebe in thoſe parts where it is known to breed.

GREBE-TIPPET. Vide *Grebe creſted.*

GREEN-LEGGED-HORSEMAN. Vide *Greenſhank.*

GREENSHANK.

Scolopax glottis. *Lin. Syſt.* i. p. 245. 10. *Gmel. Syſt.* ii. p. 664. *Ind. Orn.* ii. p. 720. 21.

Limoſa griſea. *Brif.* v. p. 267. 2. t. 23. f. 1.—*Ib.* 8vo. ii. p. 278.

La Barge variée. *Buf.* vii. p. 503?

Pluvialis major. *Raii Syn.* p. 106. A. 8.—p. 190. 6. *Will.* p. 220. t. 55.—*Ib. Angl.* p. 298.

Green-legged Horſeman. *Albin.* ii. t. 69.

Greenſhank. *Br. Zool.* ii. N° 183.—*Ib.* fol. 121. t. C. 1. *Arct. Zool.* ii. N° 379. *Lath. Syn.* v. p. 147. 18.—*Ib. Sup.* p. 245. *Lewin, Br. Birds,* iv. t. 163. *Walc. Syn.* ii. t. 142.

PROVINCIAL.

Greater Plover.

The length of this ſpecies of Snipe is fourteen inches; weight about ſix ounces.

Bill about two inches and a half long, duſky, ſlender; irides hazel.

The head, back and ſides of the neck, and back, cinereous, ſtreaked with duſky; from the upper mandible to the eye a white ſtreak; the under parts, from chin to tail, white, but the white part is narrow down to the fore part of the neck; lower part of the back and rump white; greater wing coverts, ſcapulars, and three or four of the quill-feathers next the body, brown, gloſſed with green; the edges of each feather ſcallopped with duſky and aſh-colour, ſome of them barred on both webs; ſmaller coverts duſky; quills duſky, the inner webs of ſome ſpotted with white; tail white, croſſed with duſky bars; legs very long and ſlender, of a duſky green colour; the outer toe united to the middle one as far as the firſt joint.

In ſome the coverts of the wings, ſcapulars, and upper part of the back, are aſh-colour. Such are probably adults, for we have obſerved the ſpots and bars frequently found on a variety of Snipes and Sandpipers, as well as others, in the autumn, are rarely met with in the ſpring. Some variation is alſo obſerved in the length of the bill and legs; and the bare part of the thigh is in ſome nearly two inches above the knee, in others not above an inch.

Theſe birds are ſometimes ſeen in ſmall flocks

on our coaſts in winter; as alſo in the marſhes and fens contiguous to the ſea. Some few are ſuppoſed to remain with us all the ſummer, and to breed in our fens, from whence we received an egg ſaid to belong to this bird. It is rather leſs than that of the Lapwing, and not very unlike in ſhape and colour, being of an olive-brown, covered with duſky ſpots all over, but ſmaller than thoſe of the Lapwing. The greater part, however, retire northward to breed, and are found in Sweden, Ruſſia, and Siberia. Has been alſo obſerved in America in the province of New York.

GREY, or GRAY. Vide *Gadwall.*

GREY-COOTFOOTED-TRINGA. Vide *Phalarope-grey.*

GREY-PATE. The young of the Goldfinch, ſo called by the birdcatchers before the crimſon on the head appears. *Vide* Finch-gold.

GREY-SKIT. Vide *Rail-water.*

GROSBEAK. A genus of birds, the characters of which are,

Bill ſtrong, both mandibles convex and thick.
Noſtrils ſmall, round.
Tongue truncated.
Toes, three before, one behind.

GROSBEAK-GREEN.

Loxia Chloris. *Lin. Syſt.* i. p. 304. 27. *Gmel. Syſt.* ii. p. 854. *Raii Syn.* p. 85. A. 4. *Will.* p. 129.

p. 129. t. 44. *Ind. Orn.* i. p. 382. 39. *Brif.* iii. p. 190. 54.—*Ib.* 8vo. i. p. 359.

Le Verdier. *Buf.* iv. p. 172. t. 15.

Greenfinch, *or* Green Groſbeak. *Br. Zool.* N° 117.—*Ib.* fol. 107. *Arct. Zool.* ii. p. 253. B. *Will. Angl.* p. 246. t. 44. *Albin.* i. t. 58. *Lath. Syn.* iii. p. 134. 36.—*Ib. Sup.* p. 152. *Lewin, Br. Birds,* ii. t. 69. *Pult. Cat. Dorſet.* p. 11. *Walc. Syn.* ii. t. 208.

PROVINCIAL.

Green Linnet.

This ſpecies of Groſbeak is rather larger than the Houſe Sparrow; weight near eight drams; length ſix inches and a half.

The bill is thick and whitiſh; irides dark hazel. The head and back yellowiſh green; the edges of the feathers greyiſh, inclining to aſh-colour about the ſides of the head and neck; the rump and breaſt more yellow; the greater quills are yellow on the outer webs; thoſe next the body greyiſh; the tail is ſomewhat forked; the middle feathers duſky; the four outer feathers on each ſide are yellow on their exterior webs; legs fleſh-colour.

The plumage of the female is much leſs vivid, inclining to brown.

This is a very common bird in moſt parts of England in ſummer; becomes gregarious in winter, flocking with Chaffinches and Yellowhammers; but

but in ſevere weather entirely quits ſome diſtricts. It is rather a late breeder. The neſt is compoſed of ſmall dry twigs, bents, and moſs interwoven with wool, and lined with hair and feathers; is commonly placed amongſt ivy ſurrounding a tree, or in ſome thick buſh. The eggs are four or five in number, white, ſpeckled with ruſty red at the larger end, much like thoſe of the Linnet, but larger; their weight thirty-ſeven grains. Its food is principally ſeed and grain.

The native ſong of this bird is triffling, but in confinement it becomes very tame and docile, and will catch the notes of other birds.

GROSBEAK-HAW.

Loxia Coccothrauſtes. *Lin. Syſt.* i. p. 299. *Gmel. Syſt.* ii. p. 844. *Raii Syn.* p. 85. A. 1. *Will.* p. 178. t. 44. *Ind. Orn.* i. p. 371. 4. *Briſ.* iii. p. 219. 1.—*Ib.* 8vo. i. p. 370.

Le Gros-bec. *Buf.* iii. p. 444. t. 27. f. 1.

Groſbeak, *or* Hawfinch. *Br. Zool.* N° 113.—*Ib.* fol. 105. t. U. f. 1. *Arct. Zool.* ii. p. 354. C. *Will. Angl.* p. 244. t. 44. *Albin.* i. t. 56. *Edw.* t. 188. *Lewin, Br. Birds,* ii. t. 67. *Lath. Syn.* iii. p. 109. 4.—*Ib. Sup.* p. 148. *Pult. Cat. Dorſet.* p. 11. *Walc. Syn.* ii. t. 206. *Don. Br. Birds,* ii. t. 43.

PROVINCIAL.

Cherry Finch.

The length of this ſpecies is ſix inches; weight about two ounces.

The bill is three quarters of an inch long, half an inch thick at the baſe, and remarkably ſtrong, of a pinkiſh hue when alive, but ſoon fades to a light-brown horn-colour; irides grey. The crown of the head and cheeks are bay; between the bill and eye, and round the noſtrils, is black; the chin and throat the ſame; hind part of the neck aſh-coloured; the back and ſcapulars cheſnut-brown; rump and upper tail coverts light brown, inclining to aſh-colour; the breaſt light brown, tinged with bloſſom-colour, whitiſh toward the vent; the greater quills are duſky black; the points, from the fourth, including the ſecondary quills, as far as the fifteenth or ſixteenth, are gloſſy bluiſh-black, truncated at their ends, and four or five of them bent in form of a battle-axe; on the middle of their inner webs a white ſpot; three or four of the leſſer quills cloſe to the body are the colour of the back; the ſmalleſt coverts duſky; beneath a few are white, making a ſmall bar of that colour on the wing; the greateſt coverts of the ſecondary quills are cinereous grey on their outer webs; the tail is black; the inner webs half white from the points, except the two middle feathers; the coverts of the tail are very long, and reach nearly to the end; the legs are pale brown.

We are informed the female wants the black ſpot on the chin. It is probable, however, ſuch may

may be a variety, as we have examined a great many of both ſexes, all of which had more or leſs black on that part; but the females in general have leſs of the bay-colour on the head. However, this is not to be depended on, as they vary conſiderably in that part from age.

Dr. Latham ſays this ſpecies is ſubject to great variety of plumage; that the top of the head in ſome is whitiſh, ſurrounded with rufous, in others wholly black; that the band in the wings in ſome is almoſt white, in others grey, and again wholly wanting; and that ſpecimens have been ſeen wholly black.

Theſe birds uſually viſit England in the autumn, and continue with us till the month of April. It appears in ſmall flocks, ſeldom more than four or five, but are in no parts common. We once ſaw as many as a dozen together, feeding on the haw-thorn berries in Badmington Park in Glouceſterſhire. The facility with which they break the hard ſtones of that fruit to get at the kernel is aſtoniſhing. It is done apparently with as much eaſe as other ſmall birds break hempſeed.

No inſtance has been recorded of its breeding with us; but Dr. Latham aſſures us he had one ſent to him in the ſummer months. What the ſong of this bird may be in the ſeaſon of love, authors are ſilent about; but we have heard it ſing pleaſantly, in low plaintive notes, even in winter, when the weather has been unuſually warm.

It is more plentiful in France; appears about Burgundy in April, where it breeds.

The neſt is compoſed of dried fibres intermixed with liverwort, and lined with finer materials.

The eggs are of a bluiſh green, ſpotted with olive-brown, with a few irregular black markings. Is alſo common in Italy, Germany, Sweden, and part of Ruſſia.

GROSBEAK-PINE.

Loxia Enucleator. *Lin. Syſt.* i. p. 299. 3. *Gmel. Syſt.* ii. p. 845. *Ind. Orn.* i. p. 372. 5.

Coccothrauſtes canadenſis. *Briſ.* iii. p. 250. 15. t. 12. f. 3.—*Ib.* 8vo. i. p. 378.

Le Dur-bec, *ou* Gros-bec de Canada. *Buf.* iii. p. 457.

Greateſt Bulfinch. *Edw.* t. 123. 124. M. and F.

Pine Groſbeak. *Br. Zool.* i. N° 114. t. 49. f. 2. *Arct. Zool.* ii. N° 209.—*Ib. Sup.* p. 64. *Lath. Syn.* iii. p. 111. 5.—*Ib. Sup.* p. 148. *Lewin, Br. Birds*, ii. t. 68. *Wale. Syn.* t. 207. *Don. Br. Birds*, i. t. 17.

This is larger than the laſt ſpecies; weight rather more than two ounces; length above ſeven inches. Bill thick at the baſe, and hooked at the point, colour duſky; irides hazel.

The head, neck, breaſt, and rump, crimſon; the back and leſſer coverts of the wings black, edged with reddiſh; the greater wing coverts tipped with

with white, forming two bars on the wing; quill-feathers duſky, edged on their external webs with dirty white; lower part of the belly and vent aſh-colour; the tail is a little forked, duſky, margined whitiſh; legs brown.

The female is brown, tinged with green; in ſome yellowiſh.

This bird is met with in this kingdom only in the moſt nothern parts, inhabiting the pine forests of Scotland, feeding on the ſeeds of that tree. It is ſuppoſed they breed in thoſe parts, as Mr. Pennant ſaw them flying about the pines in the foreſt of Invercauld in Aberdeenſhire on the 5th of Auguſt.

Dr. Latham obſerves they are found in North America and Hudſon's Bay; that they make their neſt in trees at a ſmall height from the ground, compoſed of ſticks, lined with feathers, laying four white eggs.

Is found alſo in the pine foreſts of Siberia, Lapland, and the northern part of Ruſſia.

GROUS. A genus of birds, the characters of which are,

Bill convex, ſhort, ſtrong.
Skin above the eyes naked, ſcarlet.
Noſtrils hid in the feathers.
Tongue pointed.
Legs feathered.

GROUS-BLACK.

Tetrao Tetrix. *Lin. Syſt.* i. p. 274. 2. *Gmel. Syſt.* ii. p. 784. *Ind. Orn.* ii. p. 635. 3.

Urogallus minor. *Raii Syn.* p. 53. A. 2. *Will.* p. 124. t. 31. *Briſ.* p. 186. 2.—*Ib.* 8vo. i. p. 52.

Le Coq de bruyeres a queue fourchue. *Briſ.* ii. p. 210. t. 6.

Black Cock, Black Game *or* Grous, Heath Cock. *Br. Zool.* i. N° 93. t. 42.—*Ib.* fol. 85. t. M. 1. 2. *Arct. Zool.* ii. p. 314. C. *Will. Angl.* p. 173. t. 31. *Albin.* i. t. 22. *Lath. Syn.* iv. p. 733. 3.—*Ib. Sup.* p. 213. *Lewin, Br. Birds,* iv. t. 133. *Pult. Cat. Dorſet.* p. 7. *Walc. Syn.* ii. t. 181. *Don. Br. Birds,* iv. t. 97.

PROVINCIAL.

Heath Fowl, or Heath Poult.

This ſpecies ſometimes weighs as much as four pounds; length about twenty-three inches. Bill duſky; irides hazel. The head, neck, and whole body, is of a gloſſy blue-black, particularly about the neck, breaſt, and rump; over the eye the bare ſcarlet ſkin is granulated; the coverts of the wings duſky brown; the four firſt quill-feathers black; the next white at the bottom; the lower half and tips of the ſecondaries white; under wing coverts white; the thighs are dark brown, ſometimes marked with a few white ſpots; the tail conſiſts of

ſixteen

ſixteen black feathers; the exterior ones bend outwards, and are much longer than thoſe in the middle, which makes the tail very forked; the under tail coverts pure white; legs covered with hair-like feathers of a dark brown, ſpeckled with grey; toes pectinated.

The female weighs about two pounds; the plumage is very different from that of the male. The general colour is ferruginous, barred and mottled with black above; the under parts paler, with duſky and brown bars; the tail-feathers are ſtrait and even at the end, variegated with ferruginous and black.

The Black Grous is at preſent chiefly confined to the more northern parts of this kingdom, population and culture having driven them from the ſouth, except in a few of the more wild, uncultivated parts; in the New Foreſt in Hampſhire, Dartmoor and Sedgemoor in Devonſhire, and the heathy hills in Somerſetſhire, contiguous to the latter. It is alſo found in Staffordſhire, and in North Wales, and again in the north of England; but no where ſo plentiful as in ſome parts of the highlands of Scotland. The males are polygamous, and fight deſperately for the females.

In the month of April the male places himſelf on an eminence as ſoon as it is light in the morning, crows and claps his wings, to which the females reſort.

After the courting ſeaſon the males aſſociate peaceably

Notwithſtanding the above accounts we cannot help entertaining ſome doubt of its being a mixed breed; a circumſtance ſo unnatural in a ſtate of nature, and of which we find no other inſtance.

If this ſpecies is met with ſo common in Sweden, how is it that none but males are found?

The Wood Grous, as well as the Black Grous, are found plentiful in Norway, and numbers are brought into this country every year; and yet we cannot learn that any ſuch bird as the Hybridus is found there. And yet Dr. Latham ſays he was informed by Dr. Tunſtall, that he was told by ſome old Scotch gentlemen that both the Wood, as well as the Spurious Grous, were extant in Scotland within their memory.

We muſt, however, conſider this matter as in a ſtate of obſcurity, and that nothing has been advanced on the ſubject to induce us to believe the bird in queſtion is any other than a mere variety of the male Black Grous. Some authors have, indeed, deſcribed the female to be of a grey colour, ſpotted with black.

A ſingular bird is mentioned in the *Naturaliſt's Calendar*, ſuppoſed to be a mixed breed between the Common Fowl and Pheaſant, ſhot in the woods. But then, in this caſe, the male or female was a domeſticated bird, and moſt probably the former. For further particulars, ſee the article Pheaſant.

GROUS-RED.

Tetrao Lagopus. *Gmel. Syſt.* ii. p. 750. 4. γ.

Tetrao Scoticus. *Ind. Orn.* ii. p. 641. 15.

Bonaſa Scotica. *Briſ.* i. p. 199. 5. t. 22. f. 1.—*Ib.* 8vo. i. p. 55. *Buf.* ii. p. 252.

Attagen. *Briſ.* i. p. 209. 9.—*Ib.* 8vo. i. p. 58. *Buf.* ii. p. 252.

Red Game, Moorcock, Gorcock. *Raii Syn.* p. 54. A. 3. *Will. Angl.* p. 177. *Albin.* i. t. 23. 24.

Red Grous. *Br. Zool.* i. N° 94. t. 43.—*Ib.* fol. 85. t. M. 3. *Lath. Syn.* iv. p. 746. 13.—*Ib. Sup.* p. 216. *Lewin, Br. Birds,* t. 135. *Walc. Syn.* ii. t. 183.

This ſpecies weighs about twenty ounces; length ſixteen inches. Bill black; irides hazel; above the eye is a ſcarlet fringed membrane bare of feathers. The noſtrils are covered with black and ferruginous feathers; the head and neck pale tawny, ſpotted black; breaſt and belly dull purpliſh-brown, croſſed with numerous narrow duſky lines; quills duſky; the tail conſiſts of ſixteen feathers; the four middle ones are barred with tawny-red, the reſt black; legs covered with ſoft whitiſh feathers down to the claws, which are of a light horn-colour, broad and concave underneath.

The female weighs about ſixteen ounces; the colour not ſo dark as in the male.

This bird is only to be met with in the extenſive uncultivated

uncultivated waſtes covered with heath, particularly the moſt mountainous ſituations; are driven from the ſouth by cultivation. The mountains of Wales are now the moſt ſouthern parts theſe birds are found in; are not uncommon in Yorkſhire, and from thence northward upon the moor lands; but no where ſo plentiful as in the highlands of Scotland, where the moors are unbounded.

It is alſo found on the weſtern iſlands, and in the mountains and bogs of Ireland; but it is remarkable that theſe birds ſhould ſeem to be confined to theſe kingdoms. Linnæus did not ſeem to be acquainted with it, and Gmelin has given it as a variety of the Ptarmigan. Buffon ſpeaks of a white variety which he names L'altagas blanc, and ſays it is found about the mountains of Switzerland, and thoſe of Vicenza. But there is little doubt this is the Ptarmigan.

The Red Grous never reſort to woods, but confine themſelves wholly to the open moors, feeding on the mountain and bog berries, and, in defect of theſe, on the tops of the heath.

It lays from eight to fourteen eggs, much like thoſe of the Black Grous, but ſmaller. The young keep with the parent birds till towards winter, and are called a pack or brood; in November they flock together in greater numbers, ſometimes thirty or forty, where they are plentiful, at which time they are extremely ſhy, and difficult to be ſhot.

We never remember but one inſtance of its

being found at a diſtance from the moors. This was a female, taken alive near Wedhampton in Wiltſhire, in the winter of the year 1794, and communicated by the late Edward Poore, Eſq. who ſhewed us a part of the bird. By what unaccountable accident it ſhould have been driven to ſo great a diſtance from its native moors is difficult to be aſſigned, as the neareſt place they are known to inhabit is the South of Wales; a diſtance, in a ſtrait line, not leſs than ſixty miles.

GROUS-SPURIOUS. Vide *Grous-hybridus.*

GROUS-WOOD.

Tetrao urogallus. *Lin. Syſt.* i. p. 273. 1. *Gmel. Syſt.* i. p. 746. *Raii Syn.* p. 53. A. 1. *Will.* p. 123. t. 30. *Ind. Orn.* ii. p. 634. 1.

Urogallus major. *Briſ.* i. p. 182.—*Ib.* 8vo. i. p. 51.

Coq de Bruyere, *ou* Tetras. *Buf.* ii. p. 191. t. 5.

Capricalca. *Sib. Scot.* xvi. t. 14. 18.

Cock of the Wood or Mountain. *Raii Syn.* p. 53. A. 1. *Will. Angl.* p. 172. t. 30. *Albin.* ii. t. 29, 30.

Wood, *or* Great Grous. *Br. Zool.* i. N° 92. t. 40, 41.—*Ib.* fol. M. M. *Arct. Zool.* ii. p. 312. A.—*Ib. Sup.* p. 62. *Lath. Syn.* iv. p. 729. 1. *Lewin, Br. Birds,* iv. t. 132. *Walc. Syn.* ii. t. 180. *Don. Br. Birds,* iv. t. 89.

In

In Scotland commonly known by the name of Capercalze, or Caperkally.

This noble ſpecies is not much inferior in ſize to a Turkey; weight ſometimes twelve or thirteen pounds, but more frequently ſeven or eight; length two feet eight or nine inches. The bill is above two inches long, very ſtrong, the upper mandible much convex and hooked, the point hanging over the under mandible very conſiderably when cloſed, as in birds of prey, and projecting over the ſides, not meeting at the edges, as in moſt birds, by which means it can cut its food like a pair of ſciſſars; the colour yellowiſh; irides hazel. The noſtrils are covered with duſky feathers; over the eye is a bare red ſkin, under the eye a ſmall ſpot of white feathers; the head is duſky, a little daſhed with aſh-colour; the feathers on the chin and throat are duſky black, and long; the neck dark aſh-coloured, finely ſpeckled with duſky; the breaſt is of a fine dark gloſſy green; the reſt of the under parts black, with ſpots of white, moſt about the thighs and vent; the wing coverts and ſcapulars cheſnut-brown, finely ſpeckled with duſky, at the junction of the wing to the body is a little white; greater quills duſky; ſecondaries like the coverts, ſlightly tipped with white; the lower part of the back, rump, and upper tail coverts, aſh-colour, marked with innumerable ſmall undulated lines and ſpecks of black; the two outer rows of feathers

thers covering the tail are greatly longer than thoſe in the middle, and gradually lengthening, the under ones reaching nearly to the end of the tail; theſe are ſeven or eight in number, laying immediately over each other; their ends are white, making as many white bars on each ſide; the tail is conſiderably rounded, conſiſting of eighteen black feathers, marked with a few ſpots of white on the ſides; the legs are covered with brown hair-like feathers; the edges of the toes ſtrongly pectinated; claws duſky and blunt.

The female differs exceedingly both in ſize and colour; weight about four pounds. Bill duſky; head, neck, and back, are barred with tawny-red and black; the throat tawny-red; breaſt pale tawny, with ſome white ſpots on the upper part; the belly barred with pale tawny and black, the feathers tipped with white; ſcapulars tipped the ſame; quill-feathers duſky, mottled on the exterior webs with light brown; the tail is of a dark ruſt-colour, barred with black, and tipped with white.

This ſex is deſcribed by ſome authors to have but ſixteen feathers in the tail. This variation muſt have been occaſioned by the loſs of ſome not noticed, for we have never ſeen an inſtance where the ſexes differ in this particular. It is a ſtrong character, that ſeems to divide birds of apparent ſimilitude, as in the Corvorant and Shag, and many of the Duck tribe, the females of which ſo frequently reſemble each other.

The male of this ſpecies is polygamous, and lives ſeparate from the females, except in the amorous ſeaſon. Their manner and habits are very like thoſe of the Black Grous, except that this ſeems to be confined wholly to foreſts of pine, on the tender ſhoots of which it feeds. It was formerly met with in Scotland and Ireland, but are are now extinct.

The female is ſaid to lay from eight to ſixteen eggs, of a white colour, ſpotted with yellow, larger than thoſe of our domeſtic fowl. Dr. Latham ſays he is well informed the neſt of one found in Scotland was placed on a Scotch pine; if ſo, it differs from all the genus, who are known to lay their eggs on the bare ground.

It is not uncommon in the pine foreſts of Norway, from whence we have received it. Is alſo found plentiful in Ruſſia and Siberia, in Italy, and ſeveral parts of the Alps.

GUILLEM. Vide *Guillemot-fooliſh*.

GUILLEMOT. A genus of birds, the characters of which are,

Bill ſlender, pointed; the upper mandible ſlightly bending towards the end; baſe covered with ſhort feathers.

Noſtrils lodged in a hollow near the baſe.

Tongue ſlender, almoſt the length of the bill.

Legs furniſhed with three toes, all placed forward, and webbed.

GUILLEMOT-BLACK.

Colymbus Grylle. *Lin. Syſt.* i. p. 220. 1. *Gmel. Syſt.* ii. p. 584.

Uria Grylle. *Ind. Orn.* ii. p. 797. 2.

Uria minor nigra; Columba groenlandica. *Briſ.* vi. p. 76. 3.—*Ib.* 8vo. ii. p. 379. *Raii Syn.* p. 121. 6. *Will.* p. 245.

Le petit Guillemot. *Buf.* ix. p. 354.

Greenland Dove, *or* Sea Turtle. *Albin.* ii. t. 80. *Will. Angl.* p. 326. t. 78.

Black Guillemot. *Br. Zool.* ii. N° 236.—*Ib.* fol. 138. t. H. 4. *Arct. Zool.* ii. N° 437. *Lath. Syn.* vi. p. 332. 3. *Lewin, Br. Birds,* vi. t. 221. *Walc. Syn.* i. t. 95. *Pult. Cat. Dorſet.* p. 17.

PROVINCIAL.

Scraber.

This ſpecies weighs fourteen ounces; length near fourteen inches. The bill is black, ſtrait, ſlender; inſide of the mouth yellow-red.

The whole plumage is black, except ſome of the wing coverts and ſecondary quills, which are tipped with white; legs red.

Some are ſaid to be found wholly black; others ſpotted black and white; and that the plumage is changed to white in winter.

Theſe birds are not frequent on the ſouthern coaſts, but are not uncommon in the north, on the Farn iſlands, and in Scotland in the Hebrides. We

have ſeen it rarely on the coaſt of Wales near Tenbeigh, where a few breed annually; but no where elſe that we could find from thence to St. David's.

It lays one egg of a dirty white, blotched with pale ruſt-colour, which is depoſited under ground, or in ſome hole in the rocks.

Frequent in Greenland, Hudſon's Bay, and other northern parts.

GUILLEMOT-FOOLISH.

Colymbus Troile. *Lin. Syſt.* i. p. 220. 2. *Gmel. Syſt.* ii. p. 585.

Uria. *Briſ.* vi. p. 70. 1. t. 6. f. 1.—*Ib.* 8vo. ii. p. 377.

Uria Troile. *Ind. Orn.* ii. p. 796. 1.

Lomwia Hoieri. *Raii Syn.* p. 120. A. 4. *Will.* p. 244. t. 65.

Le Guillemot. *Buf.* ix. p. 350. t. 25.

Fooliſh Guillemot. *Br. Zool.* ii. N° 234.—*Ib.* fol. 138. t. H. 3. *Artt. Zool.* ii. N° 436. *Will. Angl.* p. 324. *Albin.* i. t. 84. *Edw.* t. 359. f. 1. *Lath. Syn.* vi. p. 329. 1.—*Ib. Sup.* p. 265. *Lewin, Br. Birds,* vi. t. 220. *Pult. Cat. Dorſet.* p. 17. *Walc. Syn.* i. t. 96. *Don. Br. Birds,* ii. t. 28.

PROVINCIAL.

Sea-hen. Scout. Kiddaw. Murre. Lavy. Willock. Tinkerſhire.

This

This ſpecies weighs about twenty-four ounces; length ſeventeen inches.

The bill is black, three inches long, ſtrait, ſharp-pointed; inſide of the mouth yellow; irides duſky. The baſe of the bill is covered with ſoft feathers, which, with the head, neck, back, wings, and tail, are of a deep mouſe-coloured brown; the tips of the leſſer quill-feathers white; the whole under ſide of the body pure white; on the ſides under the wings a few duſky lines; from the eye to the hind-head is a ſingular line occaſioned by a diviſion of the feathers; legs duſky.

Theſe birds are found in great abundance in various parts of our high rocky coaſts from north to ſouth, and in ſome places perfectly ſwarm. It is not uncommon to ſee hundreds ſitting upon their eggs on the ledge of a rock in a line, nearly touching each other.

The female lays but one egg, of a greeniſh colour, blotched and marbled with duſky, ſo variable that ſcarce two are ſeen alike. They ſeldom quit their eggs unleſs diſturbed, but are fed with ſprats and other ſmall fiſh by the male. In places where they are ſeldom diſturbed it is with difficulty they are put to flight, and may ſometimes be taken with the hand; others flutter into the water, appearing not to have much uſe of their wings.

They begin to ſettle on their breeding places early in May, and wholly leave the ſouthern parts of the kingdom the latter end of Auguſt.

The Razorbill is frequently found in the ſame ſituation, but ſeldom breed on the ſame cliff, at leaſt not cloſe to or intermixed with the groups of Guillemots.

Both theſe birds are indiſcriminately called Willocks and Murres in ſome parts.

The young ſeldom leave the rocks till they can fly, and are of the ſame colour with the parent birds.

GUILLEMOT-LESSER.

Colymbus minor. *Gmel. Syſt.* i. p. 585.
Uria Troile. *Ind. Orn.* ii. p. 797. 1. β.
Leſſer Guillemot. *Br. Zool.* ii. N° 235. t. 83.
—*Ib.* fol. 138. *Arct. Zool. Sup.* p. 69. *Lath. Syn.* vi. p. 332. 2. *Wale. Syn.* t. 97.

PROVINCIAL.

Winter Guillemot. Morrot.

This bird is leſs than the preceding ſpecies; its weight is eighteen or nineteen ounces; length about ſixteen inches. Bill and irides the ſame as the laſt. The upper part of the head, hind part of the neck, back, wings, and tail, are black; from the eye a duſky ſtreak pointing backwards; the tips of the ſecondary quill-feathers are ſlightly marked with white; the cheeks, throat, and all beneath, white; along the ſides and on the thighs a few duſky ſtreaks; legs black.

Theſe

Theſe are found ſparingly on the ſouthern coaſts in winter only, but more frequently in the north.

Many doubts have been entertained with regard to this bird. Dr. Latham, in his *Synopſis*, was inclined to believe it no other than the Fooliſh Guillemot, and in his *Index Ornithologicus* has put them together. There are ſome circumſtances, however, that forbid us from coinciding with that excellent ornithologiſt. It is to be obſerved, that beſides the difference in ſize and plumage, this bird is rarely met with in the ſouth till the month of November; whereas the other ſpecies always leave us before September, and do not appear again till the enſuing ſpring; and that the young are at the time they depart exactly like the old ones. In the ſame ſtate of doubt and uncertainty has the Razorbill and the Black-billed Auk been held, under the heads of which we ſhall have occaſion to ſay more on the ſubject.

Mr. Pennant obſerves, theſe and the Black-billed Auks haunt the Firth of Forth, during winter, in large flocks, where they are called Morrots; and that they retire before ſpring. Where ſuch opportunity offers it is much to be wiſhed ſome naturaliſt would pay attention to the doubts in queſtion.

GUILLEMOT-SPOTTED.

Uria Grylle. *Ind. Orn.* ii. p. 797. 2. var.
Black Guillemot. *Lath. Syn.* vi. p. 333. var.
Spotted Guillemot. *Br. Zool.* ii. t. 83. f. 2.
Spotted Greenland Dove. *Edw.* t. 50.

There is no doubt but this bird is a mere variety of the Black Guillemot, more or leſs ſpotted.

GULDEN-HEAD. Vide *Puffin.*

GULL. A genus of birds, the characters of which are,

Bill ſtrong, ſtrait, but bending down at the point; on the lower mandible an angular prominence.

Noſtrils oblong, narrow, placed in the middle of the bill.

Tongue a little cloven.

Body light; very full of feathers; wings long.

Legs ſmall, naked above the knees.

Feet ſmall; toes before webbed; back toe ſmall.

GULL-ARCTIC.

Larus Paraſiticus. *Lin. Syſt.* i. p. 226. 10. *Gmel. Syſt.* ii. p. 601. *Ind. Orn.* ii. p. 819. 15.

Stercorarius longicaudus. *Briſ.* vi. p. 155. 3.—p. 150. 1.—*Ib.* 8vo. ii. p. 402.—p. 401.

L'Abbe à longue queue. *Buf.* viii. p. 445.

Strunt-jager. *Raii Syn.* p. 127. 2.

Arctic Birds. *Edw.* t. 148. 149. M. & F.

Arctic Gull. *Br. Zool.* ii. N°245. t. 87. M. & F. *Arct. Zool.* ii. N°459. *Lath. Syn.* vi. p. 389. 16. t. 99. *Lewin, Br. Birds,* vi. t. 207. *Walc. Syn.* i. t. 116.

GUL

PROVINCIAL.

Feaſer. Dung-hunter.

The length of this ſpecies is about twenty-one inches. The bill is an inch and a half long, pretty much hooked, and of a duſky colour; noſtrils placed in a kind of cere.

The top of the head is black; on the ſides, forehead, neck, and under parts of the body, white; acroſs the breaſt a pale duſky bar; the upper parts of the body, wings, and tail, black; the baſe of the quills white on the inner webs; the two middle feathers of the tail are four inches longer than the reſt; legs black, ſcaly.

This, like all the genus, is liable to vary in plumage according to age. In ſome the white parts are mixed with brown, and the black parts duſky brown; the middle feathers of the tail alſo very little longer than the reſt.

The female is ſaid by ſome to be entirely brown, paleſt beneath; others aſſert this ſex to be like the male, which is moſt probable, and that the brown variety is only young birds in their firſt feathers.

This bird is not uncommon in the Hebrides and in the Orknies, where they breed amongſt the heath, appearing in May, and retire in Auguſt. Has been met with as far ſouth as Yorkſhire.

It is ſaid to make a neſt of graſs and moſs in ſome marſhy place, and lay two eggs, the ſize of thoſe of a Hen, aſh-coloured, ſpotted with black.

This

This and other ſpecies of Gulls purſue the leſſer ones, not for the ſake of their dung, as ſome have aſſerted, but to make them diſgorge, which they catch with great dexterity before it reaches the water.

It is to be remarked all this tribe are voracious, and if purſued by a Hawk or other bird that creates alarm, it readily diſgorges in order to lighten itſelf and eſcape by flight. It is no uncommon thing to ſee them bring up a large quantity of half-digeſted food when ſlightly wounded by ſhot: tamed Gulls will do the ſame if driven by a dog. Gulls float lightly on the ſurface of the water by reaſon of the quantity of feathers in proportion to their weight, and ſeem incapable of diving. If wounded ever ſo ſlightly and fall in the water, they never attempt to dive like other aquatic birds.

The Arctic Gull is found in Norway, Sweden, Denmark, and Ruſſia, as far as Kamtſchatka; is common in Greenland.

GULL-BLACK-HEADED.

Larus ridibundus. *Lin. Syſt.* i. p. 225. 9. *Gmel. Syſt.* ii. p. 601. *Ind. Orn.* ii. p. 811. 2.

Larus cinereus. *Raii Syn.* p. 128. A. 5. *Will.* p. 264.

Gavia ridibundus phœnicopos. *Briſ.* vi. p. 197. 14. —*Ib.* 8vo. ii. p. 413.

La Mouette rieuſe. *Buf.* viii. p. 433.

Brown-headed Gull. *Albin.* ii. t. 86.

Black-

Black-headed Gull. *Br. Zool.* ii. N° 252. — *Ib.* fol. 143. t. L. 5. *Arct. Zool.* ii. N° 455. *Lath. Syn.* vi. p. 380. 9. — *Ib. Sup.* p. 268. *Lewin, Br. Birds,* vi. t. 212. *Walc. Syn.* i. t. 115. *Plott. Hift. Staff.* p. 231. 7.

PROVINCIAL.

Puit, or Pewit Gull. Black-cap. Sea-Crow. Mire-Crow. Crocker.

This ſpecies is fifteen inches in length; weight nine or ten ounces. Bill ſlender, deep red; orbits the ſame; irides hazel. The head and throat black; on each eyelid a ſmall white ſpot; back and wings aſh-colour; the prime quills white, more or leſs margined and tipped with black, the firſt and ſecond ſlightly ſo on the outer web, the three next wholly white on that part; from the ſixth the exterior webs are aſh-coloured; the neck, whole under parts, and tail, are white; legs and feet red; claws black.

Theſe birds appear to be ſubject to great variety from age, or at leaſt in their change in the firſt year, and in thoſe changes have been deſcribed as different ſpecies. The Red-legged Gull of authors is only this bird before it is arrived to maturity; and there ſeems no doubt but the old birds loſe the black on the head in the winter, and do not aſſume it again till the breeding ſeaſon; but there is generally

nerally a little black about the ears; the bill and legs alſo loſe their bright colour.

We have ſeen hundreds of theſe birds together in the winter, but have never ſeen one with a black head at that ſeaſon.

They appear in great abundance in the autumn on the coaſt of Caermarthen and Glamorganſhire, particularly about the mouths of rivers. At that time the head is white, in ſome mottled with brown, with a duſky ſpot behind the ear; the back and wing coverts in young birds are mottled with brown and white; the tail croſſed with a duſky bar at the end; the bill and legs ſcarcely tinged with red.

Towards ſpring the back begins to aſſume the aſh-colour; then the wing coverts, and the bill and legs, obtain their proper colour; the black behind the ears ſpread and meet behind, and on lifting up the feathers of the crown about the month of March the ſtubs of the black feathers are to be obſerved. At this time alſo ſome few black feathers appear on the throat; but the perfect black head is not aſſumed during their ſtay in thoſe parts. In Devonſhire we have ſeen them complete in feather later in the ſpring, but have obſerved the ſame appearance in winter.

The Black-headed Gull is ſaid to breed in Lincolnſhire on the fens, and in other parts of England upon the borders of rivers.

Dr. Plott aſſures us, in his *Hiſtory of Staffordſhire*, that

that in his time theſe birds annually viſited a pool in Staffordſhire, in the pariſh of Norbury, which pool was called Pewit Pool, on account of theſe birds breeding on the iſlands. He alſo aſſures us that they would not breed on any other land than that of the proprietor of the before-mentioned place; and that, on the death of the owner, they deſerted the pool for three years, but only retired to another eſtate belonging to the next heir.

In theſe more enlightened times more ſubſtantial reaſons might be aſſigned for their leaving their uſual breeding-place. The Doctor further ſtates, that they appeared about the latter end of March or beginning of April, and retired again before winter. The young birds were accounted good eating, and were taken by driving them into nets before they could fly; that fifty dozen were taken at a driving; and that five ſhillings per dozen was the uſual price.

The young were kept alive and fattened on offal. It is alſo added, that three drivings were generally made in a ſeaſon; and that anciently as many were taken, as the profit amounted to fifty or ſixty pounds.

No author mentions their being ſeen in winter, having at that time been made a diſtinct ſpecies under various denominations, particularly that of Red-legged Gull. See that bird, under which the ſynonyms will be found.

It makes a neſt on the ground with ruſhes, dead graſs,

graſs, and ſuch like materials, and lays three eggs of an olivacious brown, marked with ruſty brown blotches.

Found in Ruſſia and Siberia. Inhabits North America.

GULL-BLACK-TOED.

Larus crepidatus. *Gmel. Syſt.* ii. p. 602. *Ind. Orn.* ii. p. 819. 14.

Catarracta Cepphus. *Raii Syn.* p. 129. 11. *Will.* p. 267.—*Ib. Angl.* p. 351. t. 67.

Stercorarius ſtriatus. *Briſ.* vi. p. 152. 2. t. 13. f. 2. —*Ib.* 8vo. ii. p. 401.

L'Abbé, *ou* Stercoraire. *Buf.* viii. p. 441. t. 34.

Black-toed Gull. *Br. Zool.* ii. N° 244. t. 86. *Arct. Zool.* ii. N°460. *Lath. Syn.* vi. p. 387. 15. —*Ib. Sup.* p. 268. *Lewin, Br. Birds,* vi. t. 216. *Walc. Syn.* i. t. 118.

The weight of this ſpecies is about eleven ounces; length fifteen or ſixteen inches.

The bill is an inch and a half long; the noſtrils are placed in a kind of cere.

The head and neck dirty white; the ſides of the laſt marked with duſky; breaſt and belly white, croſſed with numerous duſky and yellowiſh lines; ſides and vent barred tranſverſely with black and white; the back, ſcapulars, wing coverts, and tail, black, edged with white or pale ruſt-colour; the ſhafts and tips of the quills white, the exterior webs

and

and upper half of the interior black; the lower part of the latter white; tail black, tipped with white; the two middle feathers in ſome near an inch longer than the reſt; ſhafts white; the exterior webs of the outmoſt ſpotted with ruſt-colour; the legs of a bluiſh lead-colour; lower part of the toes and webs black.

We have obſerved one in the muſeum of Dr. Latham darker in colour, and mixed with ferruginous; breaſt mottled with white; the tail a little rounded at the end. This was killed near Horſham in Suſſex. Two others only are recorded to have been killed in England; one ſhot near Oxford, the other in Lancaſhire.

Said to be more frequently met with on the continent, particularly in Denmark.

Suppoſed to breed at Hudſon's Bay, arriving there in April, and makes a ſlight neſt of graſs; lays two ferruginous eggs, ſpotted with black.

GULL-BROWN. Vide *Gull-ſkua.*

GULL-BROWN-HEADED.

Larus erythropus. *Gmel. Syſt.* ii. p. 597.
Larus ridibundus. *Ind. Orn.* ii. p. 812. var. γ.
Red-legged Gull. *Arct. Zool.* ii. p. 533. E.
Brown-headed Gull. *Lath. Syn.* vi. p. 383. 11.

This bird is undoubtedly a variety of the Black-headed Gull in its approach to maturity. We have ſeen many with the head and throat mottled brown and

and white; though it is not uncommon that when they are ſo far advanced towards perfection on that part, to find them, with the coverts of the wings brown, edged with white, and the tail tipped with black, as this bird is deſcribed to be. *See* Black-headed and Red-legged Gulls.

GULL-BROWN and WHITE. Vide *Gull-wagel.*

GULL-CLOVEN-FOOTED. Vide *Tern-black.*

GULL-COMMON.

Larus canus. *Lin. Syſt.* i. p. 224. 3. *Gmel. Syſt.* ii. p. 596. *Ind. Orn.* ii. p. 815. 9.

Larus cinereus minor. *Raii Syn.* p. 127. A. 3. *Will.* p. 262. t. 76.

Gavia cinerea. *Briſ.* vi. p. 175. 8. t. 16. f. 1. —*Ib.* 8vo. ii. p. 408.

Le Grand Mouette cendrée. *Buf.* vii. p. 384.

White web-footed Gull. *Albin.* ii. t. 84.

Common Gull. *Br. Zool.* ii N° 249. t. 89. f. 2. —*Ib.* fol. 142. *Arct. Zool.* ii. N° 458.—*Ib. Sup.* p. 70. *Will. Angl.* p. 345, t. 76. *Lath. Syn.* vi. p. 378. 8. *Lewin, Br. Birds,* vi. t. 215. *Walc. Syn.* i t. 110. *Pult. Cat. Dorſet.* p. 18. *Don. Br. Birds,* ii. t. 46.

PROVINCIAL.

Sea Mall, or Sea Mew.

This ſpecies weighs fifteen or ſixteen ounces; length about ſeventeen inches.

Bill yellow; irides hazel. The head, neck, tail, and under parts of the body, white; the back, ſcapulars, and wing coverts, aſh-colour, the former tipped with white; the primary quills black; the two or three firſt have a ſpot of white acroſs the ends, but the tips are black, the reſt are tipped with white; the ſecondaries like the back, tipped with white; legs dull white, or tinged with green.

This is the deſcription of the bird maturely feathered. In the firſt year it is more or leſs mottled all over with brown and white, with a duſky bar acroſs the end of the tail. In the ſecond year the head, neck, under parts, and tail, are white, the two firſt ſtreaked with duſky; the laſt retains the bar at the end; the back and ſcapulars are aſh-coloured, but the coverts of the wings ſtill retain brown and white mottled feathers. In this ſtate it has long been conſidered as a diſtinct ſpecies, and called the Winter Mew, or Gull (ſee that bird). We ſhall here remark, that too much caution cannot be obſerved with regard to this tribe of birds; none perhaps have puzzled the naturaliſt more from the variation in plumage at different ages.

All the Gulls found in this country are more or leſs mottled for the firſt year, and ſome probably do not arrive to maturity till the third or fourth year, which we can ſpeak of as a fact from having kept them on purpoſe to aſcertain this point, and which we ſhall enlarge more upon under the article Herring Gull.

The Common Gull is one of the moſt plentiful ſpecies; is found on moſt parts of our coaſt. They breed upon the ledges of the rocks cloſe to the ſea-ſhore; ſometimes not far above the water. We ſaw ſome hundreds ſitting on their neſts in an iſland off St. David's; the neſts were made of ſea-weed, and were placed near together about fourteen feet from the beach. The eggs were two or three in number, of a dull olive-brown, blotched with duſky, the ſize of a ſmall Hen's egg. When diſturbed were exceedingly clamorous, and not much alarmed by repeatedly ſhooting at.

This bird is frequently ſeen in winter at a conſiderable diſtance from the coaſt; will flock with Rooks in ſevere weather, and follow the plough for the ſake of the *larvæ* of the Chaffer (*Scarabæus Melolonthe*) and worms.

GULL-GREAT-BLACK-BACKED.

Larus marinus. *Lin. Syſt.* i. p. 225. 6. *Gmel. Syſt.* ii. p. 598. *Ind. Orn.* ii. p. 813. 6.

Larus niger. *Briſ.* vi. p. 158. 1.—*Ib.* 8vo. ii. p. 403.

Le Goéland noir. *Buf.* viii. p. 405. t. 31.

Larus maximus ex albo et nigro varius. *Raii Syn.* p. 127. A. 1. *Will.* p. 261.

Great Black and White Gull. *Will. Angl.* p. 344. t. 67. *Albin.* iii. t. 94.

Black-backed Gull. *Br. Zool.* ii. N° 242.—*Ib.* fol. 140. t. L. *Arct. Zool.* ii. N° 451. *Lath. Syn.*

GUL

Syn. vi. p. 371. 2. *Lewin, Br. Birds,* vi. t. 208. *Pult. Cat. Dorſet.* p. 18. *Walc. Syn.* t. 112.

PROVINCIAL.

Cobb.

The weight of this ſpecies is four pounds three quarters; length near thirty inches; breadth five feet nine or ten inches.

Bill bright yellow, very thick and ſtrong, three inches and a half in length; on the lower mandible is a blood-red ſpot, duſky in the middle; irides fine yellow; eyelids red-orange. The head, neck, rump, tail, and whole under parts, white; back and wing coverts duſky black; prime quills black, the firſt tipped with white for two inches or more, the ſecond the ſame, with a black bar acroſs the white, the reſt more ſlightly tipped with white; the ſecondaries are alſo tipped more or leſs with white, as are their coverts and two or three of the ſcapulars; legs fleſh-colour.

This bird and the Leſs Black-backed Gull, hereafter deſcribed, have hitherto been confounded for the ſame ſpecies; but from frequent opportunities of obſerving their manners, as well as by diſſection, proving both ſexes to be alike in plumage, we are able without doubt to pronounce them diſtinct ſpecies.

It is not a very plentiful ſpecies, but is ſometimes ſeen on moſt of our coaſts, but no where ſo plen-

tiful as on the extenſive ſandy flats of the coaſt of Caermarthenſhire, between Laugharne and Tenbeigh, where we have ſeen it at all times of the year.

They generally keep in ſmall flocks of eight or ten, ſometimes in pairs, but never herd with the other Gulls.

It was natural to believe, as they were ſeen on that coaſt all the ſummer, they muſt breed ſomewhere near, but in our reſearches from that part as far as St. David's we could not diſcover where they bred, but were informed by the fiſhermen (who call them Cobbs) they breed on the Steep Holmes and on Lundy iſlands in the Briſtol channel. The Leſs Black-backed Gull, as well as the Herring Gull, we found the neſts of in great abundance: but none of theſe birds were to be ſeen in the ſame places.

The young, for the firſt two or three years, are mottled all over with brown and white; the bill is light horn-colour, tip black; quill-feathers duſky; tail mottled, near the end a duſky bar; tips white; irides and orbits duſky. We ſhot ſeveral birds of this deſcription in company with them, of both ſexes; their weight and ſize little inferior; and as theſe are always found to aſſociate with them, and as there is no other ſpecies of Gull half ſo large, there can be no doubt of its being the young of this bird.

In this imperfect ſtate this has been deſcribed by

by ſome authors for a diſtinct ſpecies, under the title of Wagel; others, who have not conſidered the ſize as eſſential, may have made the young of the Leſs Black-backed and Herring Gulls the Wagel; all of which are mottled nearly in the ſame manner during the firſt two or three years.

We have ſeen the young of this ſpecies with the dark colour appearing on the back. It is a great enemy to the fiſhermen; will tear and devour the largeſt fiſh from the hooks when left dry by the ebbing tide.

GULL-GREAT-BLACK AND WHITE. Vide *Gull-great-black-backed.*

GULL-GREAT-GREY.

The bird given by Ray under this title is evidently the young of one of the larger ſpecies of Gulls, probably that of the Herring Gull or Great Black-backed Gull.

GULL-HERRING.

Larus fuſcus. *Lin. Syſt.* i. p. 225. 7. *Gmel. Syſt.* ii. p. 599. *Ind. Orn.* ii. p. 815. 8.

Larus cinereus maximus. *Raii Syn.* p. 127. A. 2. *Will.* p. 262.

Larus griſeus. *Briſ.* vi. p. 162. 3.—*Ib.* 8vo. ii. p. 405.

Gavia griſea. *Briſ.* vi. p. 171. 6.—*Ib.* 8vo. ii. p. 407.

Le Goéland a manteau gris brun. *Buf.* viii. p. 379.

Herring Gull. *Br. Zool* ii. N° 246. t. 88.—*Ib.* fol. 141. *Arct. Zool.* ii. N° 452. *Lath. Syn.* vi. p. 372. 3. *Will. Angl.* p. 345. *Lewin, Br. Birds*, vi. p. 7. *Walc. Syn.* i. t. 113. *Pult. Cat. Dorſet.* p. 18.

This ſpecies weighs about thirty-three ounces; length twenty-three inches.

Bill yellow; on the lower mandible a reddiſh-orange ſpot; irides light yellow; orbits red. Head, neck, tail, and under parts, white; back, ſcapulars, and wing coverts, aſh-colour; quill-feathers duſky, the five firſt black towards their ends, with a white ſpot near the tip; legs pale fleſh-colour.

The young are at firſt mottled all over with brown and dirty white; the quills duſky, without any white at the tips; tail with a duſky bar at the end; bill horn-colour; irides duſky; legs dark. In the ſecond year the colours continue the ſame, but rather lighter; in the autumnal moulting the back becomes aſh-colour; the irides get lighter, inclining to yellow; the bill the ſame; the reſt of the plumage as before. In this ſtate we have one now living of two years old laſt breeding ſeaſon; and now, in October, ſhews no inclination to make any further change this year; from which we may fairly conclude theſe birds do not arrive at maturity till the ſpring of the third year, and perhaps not till the autumn, the principal moulting ſeaſon. Nor can we be yet certain it is then perfect; for, upon examining

examining ſpecimens we have of the next change, we find the wing coverts ſtill mottled with brown; the head and neck ſtreaked with duſky brown; the bar in the tail broken by numerous white undulated ſtreaks running down the webs; the quills, back, ſcapulars, and under parts of the body, perfect; bill and irides yellowiſh; legs pale fleſh-colour. This ſhould ſeem the laſt ſtage towards perfection; and as this was killed in the ſummer, it is moſt probable they are not perfected till the autumn of the third year.

In the two firſt years the young of this and the Leſs Black-backed Gull are ſo much alike that they cannot be aſcertained till the matured feathers begin to ſhew on the back.

Whether theſe immatured birds breed we cannot be certain; but we are inclined to believe they do, as we ſaw a great many of them intermixed with the perfect ones in a Gullery on an iſland off St. David's, where the neſts were innumerable, and were equally clamorous with the others when diſturbed. The neſts were on the top of the iſland, amongſt the graſs and looſe ſtones; they were compoſed of a ſmall quantity of long dry graſs; the eggs, two in number, of a dark olive-brown, with duſky blotches.

This bird, like others of the genus, feeds indiſcriminately on fiſh, and various other productions of the ſea, particularly the *Aſterias*, or Star-fiſh. It is ſometimes obſerved to trample the ſoft ſand by

moving the feet alternately in the ſame place. For what purpoſe this ſingular action is intended we cannot ſay, unleſs it is to force up the ſand eels, or ſome hidden prey.

The Herring Gull is found plentiful on all our ſhores, and not leſs frequent in all the northern parts of Europe, and elſewhere.

GULL-LAUGHING.

Larus atricilla. *Lin. Syſt.* i. p. 225. 8. *Gmel. Syſt.* ii. p. 600. *Ind. Orn.* ii. p. 813. 4.

Larus major cinereus Baltneri. *Raii Syn.* p. 129. 8, *Will.* p. 263. t. 67.

Gavia ridibunda. *Briſ.* vi. p. 192. B. t. 18. f. 1. —*Ib.* 8vo. ii. p. 413.

Baltner's great aſh-coloured Sea-mew. *Will. Angl.* p. 346. t. 67.

Laughing Gull. *Arct. Zool.* ii. N° 454. *Lath. Syn.* vi. p. 383. 12.

This ſpecies is larger than the Black-headed Gull; length eighteen inches. It differs from that bird only in the legs, which are black; the bill is, however, ſtronger, and the head larger. We do not recollect this bird having been noticed in England, or at leaſt recorded in the liſt of Britiſh birds.

In the month of Auguſt 1774 we ſaw five of them together feeding in a pool upon the Shingly Flats near Winchelſea; two only were black on the head, the others were mottled all over with brown.

One

One of them was ſhot; but although the remaining four continued to reſort to the ſame place for ſome time, the old ones were too ſhy to be procured. We alſo ſaw two others near Haſtings in Suſſex. They may eaſily be known from the Black-headed Gull even flying; the flight is different; the bird appears much larger, and the tail ſhorter in proportion.

It is found in Ruſſia, and in ſome parts of America; and is ſaid to be plentiful in the Bahama iſlands. Is ſuppoſed to breed at Hudſon's Bay; at leaſt a bird ſimilar to it is ſaid to make its neſt in the pine-trees, and lay four lead-coloured eggs.

GULL-LESS-BLACK-BACKED.

Larus argentatus. *Gmel. Syſt.* ii. p. 600?
Silvery Gull. *Arct. Zool.* ii. p. 533. C.—*Ib. Sup.* p. 70. *Lath. Syn.* vi. p. 375. 5.
Larus marinus. *Ind. Orn.* ii. p. 814. 6. var. β.

This ſpecies is greatly inferior in ſize to the Great Black-backed Gull, and rather ſuperior to the Herring Gull. It weighs about thirty-ſix ounces, and rarely more; length twenty-four inches. Bill yellow, with an orange ſpot on the lower mandible; ſize and ſhape like that of the Herring Gull; irides pale yellow; orbits red-orange.

Head, neck, tail, and whole under parts, pure white; back, ſcapulars, and wings, duſky black; prime quills duſky, towards their ends black; the point

point of the firſt is white, with a black tip; the ſecond the ſame, with only a white ſpot in the black; the others very ſlightly tipped with white; two or three of the ſcapulars are alſo tipped with white; legs yellow.

This bird has been confounded by moſt authors with the *Marinus*, or Great Black-backed Gull, and both, by the later writers, conſidered as the Silvery Gull; and in their firſt plumage, which is mottled all over with brown and white, have been made into another ſpecies under the title of Wagel. Both Dr. Latham and Mr. Pennant, in their laſt works on Ornithology, have, however, made the Wagel no other than the young of one or both theſe ſpecies, as well as the *Argentatas*, or Silvery Gull; but have ſtill blended the two Black-backed Gulls together. The former author has alſo made the *Larus Nævius* of Gmelin to be a variety of one of theſe birds, of which there is no doubt. The *Wagellus Cornubienſium* of Ray and Willughby is alſo conſidered as the young of one of theſe ſpecies, which is very probable. Lewin has divided the two ſpecies, but makes the ſmaller the male of the Herring Gull, and the mottled Gulls of the ſame ſize indiſcriminately the females, and erroneouſly conſiders the true Herring Gull as the young.

It is true no claſs of birds has occaſioned more perplexity than the Gulls, from the length of time moſt of them are arriving to maturity in plumage; and

and the ſpecies have been greatly multiplied. But we ſhall here again remark, that all the ſpecies we are acquainted with are, in their firſt feathers, mottled all over with brown and white; not to be diſcriminated but by ſize. Many of theſe errors have been corrected by the above authors in their laſt works; and it is certain there are no ſuch diſtinct ſpecies as the Wagel, the Winter Gull, Tarrock, or Red-legged Gull.

The firſt is the young of both the Black-backed and Herring, the ſecond that of the Common, the third that of the Kittiwake, and the laſt the young of the Black-headed Gull.

It is, however, very extraordinary that the two Black-backed Gulls ſhould have been to this day confounded, as the vaſt diſproportion in ſize is alone ſufficient to diſcriminate them. One ſpecies never exceeds two pounds and a half, the other is nearly double that weight, and as much ſtronger in the bill and legs in proportion; beſides, the legs of the greater ſpecies are invariably of a fleſh-colour, thoſe of the ſmaller are yellow; nor has this bird the dark ſpot in the orange mark on the bill.

Theſe two birds are never found to aſſociate, and always breed in diſtinct places. This congregates frequently with the Herring Gull, and breeds in the ſame places, where we have ſeen them ſitting on their neſts; but they are not near ſo plentiful. On Ramſey Iſland in Pembrokeſhire this and the Herring Gulls breed in great abundance; and where

we

we had an opportunity of examining them very attentively, this was not near ſo plentiful as the other ſpecies. The proportion was certainly not more than one in twenty, which muſt invalidate every idea of its being the male of the other ſpecies. Not a ſingle one of the Great black-backed Gull was to be ſeen on the iſland.

The eggs and young of this bird which we found in the neſts were ſo like thoſe of the Herring Gull that there was no diſcriminating mark; the eggs, indeed, were in general larger; the young were covered with a ſoft brown down, mottled with duſky.

This is by far a leſs numerous ſpecies than the Herring Gull, but much more plentiful than the Great black-backed Gull.

GULL-PEWIT, or PUIT. Vide *Gull-black-headed.*

GULL-RED-LEGGED.

Larus cinerarius. *Lin. Syſt.* i. p. 224. 4. *Gmel. Syſt.* ii. p. 597.

Larus ridibundus. *Ind. Orn.* ii. p. 812. 2. var. β.

Gavia cinerea minor. *Briſ.* vi. p. 178. 9. t. 17. f. 1.—*Ib.* 8vo. ii. p. 409.

Larus albus major. *Raii Syn.* p. 129. 9. *Will.* p. 264.

La petite Mouette cendrée. *Buf.* viii. p. 430.

Greater White Gull of Belon. *Will. Angl.* p. 348.

Red-legged Gull. *Arct. Zool.* ii. p. 533. E. *Lath. Syn.* vi. p. 381. 10.

This

This bird, ſpoken of by various authors as a diſtinct ſpecies, is no other than the Black-headed Gull in its different ſtages of change before it arrives to maturity.—*Vide* Gulls Black-headed and Brown-headed.

Is frequently known by the name of Red-legs.

GULL-SKUA.

Larus catarractes. *Lin. Syſt.* i. p. 226. 11. *Gmel. Syſt.* ii. p. 603. *Ind. Orn.* ii. p. 818. 12.

Catarractes et Catarracta. *Raii Syn.* p. 128. A. 6. —*Ib.* 129. 7. *Will.* p 265.—*Ib. Angl.* p. 348. 349. t. 67.

Larus fuſcus. *Briſ.* vi. p. 165. 4.—*Ib.* 8vo. ii. p. 405.

Le Goéland brun. *Buf.* viii. p. 408.

Brown Gull. *Albin.* ii. t. 85.

Skua Gull. *Br. Zool.* ii. N° 343.—*Ib.* fol. 140. t. L. 6. *Arct. Zool.* ii. N° 531. A. *Lath. Syn.* vi. p. 385. 14. *Lewin, Br. Birds*, vi. t. 211. *Walc. Syn.* i. t. 117.

This ſpecies is rather ſuperior in ſize to the Raven; weight three pounds; length two feet. The bill is an inch and three quarters long, black, and much hooked at the end; is covered for more than half its length with a kind of black cere.

The upper part of the head, neck, back, and wings, deep brown; the feathers margined with ferruginous; about the forehead and chin tinged with

with aſh-colour; the breaſt and all beneath pale duſky ferruginous; the quills are brown, white at the baſe; tail deep brown; roots and ſhafts white; the legs are black, rough, and ſcaly; talons black, ſtrong, and much hooked.

This is a bold rapacious bird; preys on the leſſer Gulls as well as fiſh; is ſaid to attack the Eagle, and even man if he approaches their neſt.

It breeds in the Orkney iſlands, and is much eſteemed in the iſle of Foula, from a ſuppoſition that it defends the flocks from the Eagle; is rarely ſeen in the ſouth; one in the muſeum of Dr. Latham was killed at Greenwich.

It prefers the colder climates; is common in Norway and Iceland; and alſo found in the ſouthern hemiſphere in ſeveral of the higher latitudes; has been met with at Falkland Iſland, and particularly at Port Egmont, where they were called by our circumnavigators Port Egmont Hens.

GULL-SMALL-BROWN. Vide *Tern-brown.*

GULL-SMALL-CLOVEN-FOOTED. Vide *Tern-black.*

GULL-TARROCK.

Larus tridactylus. *Lin. Syſt.* i. p. 224. 2. *Gmel. Syſt.* ii. p. 595. *Ind. Orn.* ii. p. 817. 11. β.

Larus nævius. *Lin. Syſt.* i. p. 225. 5.

Larus cinereus Bellonii. *Raii Syn.* p. 128. A. 4. *Will.* p. 263. t. 68.—p. 266. t. 66.

Gavia cinerea nævia. *Briſ.* vi. p. 185. 11. t. 17. f. 2.—*Ib.* 8vo. ii. p. 410.

La Mouette cendrée tachetée. *Buf.* viii. p. 424.

Tarrock. *Br. Zool.* ii. N° 251.—*Ib.* fol. 142. t. L 3. *Arct. Zool.* ii. 533. D.—*Ib. Sup.* p. 70. *Lath. Syn.* vi. p. 392. 18.—*Ib. Sup.* p. 268. *Lewin, Br. Birds,* vi. t. 213. *Walc. Syn.* i. t. 109.

This bird had long been conſidered as a diſtinct ſpecies, but was, in the laſt works of Mr. Pennant and Dr. Latham, given as the young of the Kittiwake, of which there is no doubt.

The ſize is the ſame as that bird; the bill is duſky black. In plumage it differs in having ſome black about the head; generally near each ear is a ſpot; ſometimes a little black under the throat; and others a creſcent of that colour on the hind part of the neck; the wing coverts duſky, edged with grey; the tail tipped with black, except the outer feather on each ſide; a ſmall protuberance inſtead of a hind toe.

This appears to be the plumage of the ſecond year, when they are found at their breeding-places with the Kittiwake. In the firſt year no doubt they are more mottled with brown, like all the young Gulls, and are not completed in feather till the third year. *Vide* Kittiwake.

GULL-WAGEL.

Larus nævius. *Gmel. Syſt.* ii. p. 598.

Larus marinus. *Ind. Orn.* ii. p. 814. 6. γ.

Wagellus

Wagellus Cornubienſium. *Raii Syn.* p. 130. A. 13. *Will.* p. 262. t. 66.

Le Goéland varié, Griſard. *Buf.* viii. p. 413. t. 33.

Wagel Gull. *Br. Zool.* ii. N° 247. A.—*Ib.* fol. ii. 422. *Artt. Zool.* N° 453. *Will. Angl.* p. 349. t. 66. *Lath. Syn.* vi. p. 375. 6. *Lewin, Br. Birds,* vi t. 209. *Walc. Syn.* i. t. 111. *Pult. Cat. Dorſet.* p. 18.

With reſpect to the plumage of this bird all authors pretty nearly agree; but vary conſiderably as to ſize. It is generally deſcribed to be from the ſize of the Herring Gull to that of the Great Black-backed Gull. The bill duſky; irides hazel. The whole plumage compoſed of a mixture of brown aſh-colour and white; quills black; tail mottled black and white, with a black bar near the end; tip white; legs duſky fleſh-colour.

Theſe birds may be ſeen in great abundance on all our ſhores in every ſeaſon of the year, and is in fact not a diſtinct ſpecies, but the young of both the Black-backed Gulls and the Herring Gull. We have ſeen all theſe birds in their different ſtages of changing their plumage, and at preſent poſſeſs a live ſpecimen which we have had upwards of two years, and now only in the autumn begins to ſhew the plumage of the Herring Gull, and probably will not be complete till this time next year. For further remarks we refer our readers to *Gull leſs-black-backed.*

GULL-WHITE-WEB-FOOTED. Vide *Gull-common.*

GULL-WINTER.

Larus hybernus. *Gmel. Syſt.* ii. p. 596.

Larus fuſcus, hybernus. *Raii Syn.* p. 130. A. 14. *Will.* p. 266. t. 66.

Guaca-guaca. *Raii Syn.* p. 130. 12. *Will.* p. 268. *Ib. Angl.* 352.

Gavia hyberna. *Briſ.* vi. p. 189. 12.—*Ib.* 8vo. ii. p. 411.

Larus canus. *Ind. Orn.* ii. p. 816. 9. β.

Mouette d'hyver. *Buf.* viii. p. 437.

Winter Mew, Çoddy Moddy. *Br. Zool.* ii. N° 248. t. 86.—*Ib.* fol. 142. t. L. 2. *Will. Angl.* p. 350. t. 66. *Albin.* ii. t. 87. *Lath. Syn.* vi. p. 384. 13. *Walc. Syn.* i. t. 114. *Lewin, Br. Birds,* vi. t. 210. *Don. Br. Birds,* iv. t. 77.

This is another of the Gulls which has been unneceſſarily multiplied into a diſtinct ſpecies, when in fact it is no other than the Common Gull in the ſecond year's plumage, the intermediate ſtage between the firſt or neſtling feathers and that of the adult. The bill is horn-colour, duſky at the tip; irides hazel. The top of the head, hind part and ſides of the neck, white, marked with duſky ſtreaks; back and wing coverts aſh-coloured, the latter mottled with brown; under parts white; quills

black, more or leſs tipped with white; the tail white, with a duſky bar near the end; legs whitiſh.

It is probable this, and ſome of the ſmaller ſpecies of Gulls, arrive at their full plumage in the ſummer of the ſecond year; and that this bird having only been obſerved in winter has occaſioned its name: but we have killed it pretty late in the ſpring. However, the brown mottled coverts of the wing, and the bar on the tail, ſeem to diſappear about the breeding ſeaſon; but frequently ſome few duſky ſtreaks on the head and neck are retained longer.

Theſe are common on all our ſhores, and are frequently ſeen together with the matured birds far inland in the winter ſeaſon, feeding on the ploughed lands.

GYRFALCON. Vide *Falcon-jer.*

H.

HAGISTER. Vide *Magpie.*

HAWK. A name commonly given to many ſpecies of the Falcon genus.

HAWK-BLUE. Vide *Henharrier* and *Falcon-peregrine.*

HAWK-DOR. Vide *Goatſucker.*

HAWK-DUCK. Vide *Falcon-peregrine* and *Moor Buzzard.*

HAWK-FISHING. Vide *Oſprey.*

HAWK-GOS.

Falco palumbarius. *Lin. Syſt.* i. p. 130. 30. *Gmel. Syſt.* i. p. 269. *Ind. Orn.* i. p. 29. 65. *Raii Syn.* p. 18. 1. *Will.* p. 51. t. 3. & 5.

Aſtur. *Briſ.* i. p. 317.—*Ib.* 8vo. p. 91.

L'Autour. *Buf.* i. p. 230. t. 12.

Goſhawk. *Br. Zool.* i. N° 52. t. 24. *Arct. Zool.* ii. N° 99. *Albin.* ii. t. 8. *Lewin, Br. Birds,* i. t. 9. *Lath. Syn.* i. p. 58.—*Ib. Sup.* p. 16. *Walc. Syn.* i. t. 9.

This is a large ſpecies of Falcon; is ſuperior in ſize to the Buzzard; length twenty-two inches or more. The bill is blue, tip black; cere yellow-green; irides yellow.

The head, hind part of the neck, back, and wings, deep brown; over the eye is a white line, and a broken patch of the ſame colour on the ſide of the neck; breaſt and belly marked with numerous tranſverſe bars of black and white; the tail is long, aſh-coloured, with four or five duſky bars; legs yellow; claws black.

Theſe birds vary a little in their markings, but the ſize and elegant ſlender ſhape at once point it out. The wing when cloſed does not reach near the end of the tail.

The Goſhawk is rarely found in England, but is

ſaid to be not uncommon in Scotland in the more wooded parts, where it breeds, and is a great deſtroyer of game. Is ſaid to build in trees, and to lay four white eggs. Is alſo found in North America.

It was held in high eſtimation in the days of falconry, and was uſed for the larger ſorts of game.

HAWK-NIGHT. Vide *Goatſucker.*

HAWK-PIGEON. Vide *Hawk-ſparrow.*

HAWK-SPARROW.

Falco Niſus. *Lin. Syſt.* i. p. 130. 31. *Gmel. Syſt.* i. p. 280. *Ind. Orn.* i. p. 44. 107.

Accipiter fringillarius. *Raii Syn.* p. 18. A. 2. *Will.* p. 51. t. 5.

Accipiter. *Brif.* i. p. 310. 1.—*Ib.* 8vo. p. 89.

Epervier. *Buf.* i. p. 225. t. 11.

Sparrow-Hawk. *Br. Zool.* i. N° 62. *Ib.* fol. t. A. 10. A. 11. *Arct. Zool.* ii. p. 25. N. *Hayes, Br. Birds,* t. 3. *Albin.* i. t. 5. *Will. Angl.* p. 86. *Lewin, Br. Birds,* i. t. 20. *Lath. Syn.* i. p. 99. 85.—*Ib. Sup.* p. 26. *Wat. Syn.* i. t. 20.

The weight of the male of this ſpecies is about five ounces, that of the female nine; the former meaſures in length about twelve inches, the latter fifteen.

The bill is bluiſh, duſky at the point; cere yellow; irides bright orange-yellow.

In

In ſome of both ſexes the plumage of the upper parts are of a deep bluiſh grey; in others brown, edged with ferruginous; the under parts of the female are more fully marked with minute undulated lines of deep brown; the male is inclined to ruſt-colour on the breaſt, which in the other is whitiſh; on the back of the head, in both ſexes, is an obſcure broken patch of white; quill-feathers duſky, barred with black on the outer webs, and ſpotted with white at the baſe of the inner; the tail like the back, with broad bars of duſky black, the extreme point whitiſh; legs long, ſlender, yellow.

This is a very common ſpecies in moſt of the wooded or encloſed parts of the kingdom, but leſs frequent in the more champaign parts. It ſeldom makes a neſt, but generally takes poſſeſſion of that which has been deſerted by a crow. It lays four or five eggs of a dirty white, ſometimes of a bluiſh tinge, blotched at the large, and ſometimes, though rarely, at the ſmaller end with ruſt-colour.

The female Sparrow Hawk is a very bold bird; has been trained for hawking with ſucceſs, though its flight is not ſo rapid as the longer-winged Hawks.

It is a great deſtroyer of game and young poultry; we have frequently known them carry away half a brood of chickens before the thief was diſcovered. They fly low, ſkim over a poultry yard, ſnatch up a chick, and out of ſight in an inſtant.

It is obſervable that the moſt generous Hawks (as

they were formerly termed), that is, the moſt tractable, have long and pointed wings, the ſecond feather being the longeſt. Of this diviſion the Falcons, properly ſo called, belong; the Hobby, Merlin, and Keſtril, are alſo of this kind.

This ſpecies, as well as the Goſhawk and all the Buzzards, are ſhort winged. Theſe have the third and fourth feather in the wing nearly of the ſame length, and longer than the ſecond; ſo that the wings when ſpread have a more rounded appearance at the end.

The more generous Hawks, we have frequently obſerved, kill their prey as ſoon as caught by eating the head firſt; whereas the Buzzards, in particular, begin eating their prey indiſcriminately. We have ſeveral times taken Partridges and other birds from them which had one ſide of the breaſt or a thigh devoured, and the bird ſtill alive.

HAY-BIRD. Vide *Pettychaps-leſſer.*

HEATH-COCK, or HEATH-FOWL. Vide *Grous-black.*

HEDGE-SPARROW. Vide *Warbler-hedge.*

HELLIGOG. Vide *Razorbill.*

HENHARRIER.

Falco cyaneus. *Lin. Syſt.* i. p. 126. 10. *Gmel. Syſt.* i. p. 276. *Ind. Orn.* i. p. 39. 94.

Pygargus accipiter (mas). *Raii Syn.* p. 17. A. 5. *Will.* p. 40. t. 7.

Falco torquatus (mas). *Brif.* i. p. 345.—*Ib.* 8vo. p. 100.

Lanarius

Lanarius cinereus. *Brif.* i. p. 365. 17.—*Ib.* 8vo. p. 106.

Oiſeau St. Martin. *Buf.* i. p. 212.

Hen Harrier. *Br. Zool.* i. N° 58. t. 28.—*Ib.* fol. p. 68. t. A. 6. *Will. Angl.* p. 72. *Edw.* t. 2. 225. *Albin.* ii. t. 5. *Hayes, Br. Birds,* t. 1. *Lewin, Br. Birds,* i. t. 18. *Lath. Syn.* i. p. 88.—*Ib. Sup.* p. 22. *Walc. Syn.* i. t. 17. *Pult. Cat. Dorſet.* p. 3. *Don. Br. Birds,* iii. t. 59.

FEMALE, or RING-TAIL.

Falco Pygargus. *Lin. Syſt.* i. p. 126. 11. *Gmel. Syſt.* i. p. 277. *Raii Syn.* p. 17. 5. *Will.* p. 40.

Falco torquatus (fem.) *Brif.* i. p. 345. 7.—*Ib.* 8vo. p. 100.

La Soubuſe. *Buf.* i. p. 215. t. 9.

Ring-tail. *Br. Zool.* i. N° 59.—*Ib.* fol. p. 68. t. A. 7. *Albin.* iii. t. 3. *Hayes, Br. Birds,* t. 2. *Lewin, Br. Birds,* i. t. 18. F. *Will. Angl.* p. 72. *Lath. Syn.* i. p. 89. 95.—*Ib. Sup.* p. 22. *Walc. Syn.* i. t. 18.

Falco Hudſonius. *Lin. Syſt.* i. p. 128. *Gmel. Syſt.* i. p. 277. *Brif.* vi. *Sup.* p. 18.—*Ib.* 8vo. p. 119.

Falco Buffoni. *Gmel. Syſt.* i. p. 277.

Ring-tail Hawk. *Edw.* t. 107. *Arct. Zool.* ii. N° 106.

White-rumped Bay Falcon. *Lath. Syn.* i. p. 54. 34. var. B.

Hudſon's Bay Ring-tail. *Lath. Syn.* i. p. 91. 76.
Cayenne Ring-tail. *Lath. Syn.* i. p. 91.

It will be manifeſt, by the above ſynonyms, that we conſider the Henharrier and the Ringtail to be of the ſame ſpecies.

In this we have followed the example of Dr. Latham, who, in his laſt ornithological works, has brought them together. We will not, however, take upon us abſolutely to determine this point, but ſhall give ſuch remarks as have recently inclined us to that opinion.

The Henharrier, in its moſt perfect ſtate, weighs about thirteen ounces; length eighteen inches and a half.

The bill is black; cere and irides yellow. Head, neck, breaſt, and whole upper parts, are of a fine light grey; round the head is a wreath of ſhort ſtiff feathers white at the baſe, ſlightly tipped with grey; the firſt ſix quill-feathers black, white at the baſe, and ſlightly tipped with grey; the reſt of the quills grey on their outer webs and tips, white on the inner webs; tail feathers white, except the two middle, on which are ſometimes a few markings; the exterior webs of all are grey, and not the leaſt appearance of any bars; vent, upper and under tail coverts pure white; legs long, ſlender, yellow.

In this bird now before us the wings when cloſed do not reach to the end of the tail by two inches; the firſt feather very ſhort, the third and fourth the longeſt, and nearly the ſame length.

Another

Another weighed twelve ounces; length eighteen inches. This is very like the firſt deſcribed, but rather inclined to brown on the ſcapulars; and the tail-feathers all, except the middle ones, barred on the inner webs more or leſs; the ſecond and third feathers from the outſide a little barred on the outer web near the ſhafts. In another ſpecimen the breaſt was ſtreaked with duſky, and ſeveral of the ſmaller coverts of the wings were ferruginous like the female; ſo that this part ſeems to be the laſt that arrives to perfection.

The Ringtail, or female, weighs about eighteen ounces; length twenty inches. Bill, cere, irides, and legs, the ſame as the Henharrier. Head and whole upper parts of a deep duſky brown; the feathers on the head, neck, and wing coverts, are margined more or leſs with rufous; round the head a wreath of ſhort feathers, rather lighter in colour; the under parts are pale rufous-brown, with large duſky ſtreaks; greater quills duſky, with a daſh of cinereous on the outer webs; upper tail coverts white; tail marked with three or four brown and duſky bars, the lighter bars ſhaded to rufous on the inner webs, which underneath appear whitiſh; the outer feather in ſome is of a light colour and plain; the tips of all whitiſh.

Various are the opinions concerning theſe birds. Some authors have ever conſidered them as male and female; others have ſhifted their opinion frequently. That males of the Ringtail are found there

there is no doubt, two inſtances of which have come under our inſpection lately; not only evident by their inferior ſize, but proved by diſſection. In one of theſe ſpecimens there are a few grey feathers on one ſide of the neck, and on one thigh, which indicates a change to the plumage of the Henharrier. But what is extraordinary, the Henharrier ſeems full as plentiful as the Ringtail; whereas if the young males are a conſiderable time arriving at maturity, we ought to ſee many more in the brown ſtate. During the whole of one ſummer we happened to be ſituated where we ſaw ſeveral Henharriers every day, frequently three or four on wing together; and yet, from the month of March to September, we never ſaw but one Ringtail. And Dr. Latham remarks, that no author has mentioned the Henharrier as a bird of the American continent, though the Ringtail and its varieties are common throughout. On the other ſide of the queſtion, the Doctor ſays that both males and females have been ſhot from their neſts in the north; and ſpeaks of it from the authority of Dr. Heyſham.

As both theſe birds are not uncommon in England, and continue with us the whole year, it is remarkable the young ones have never been taken and bred up to aſcertain the fact. But what ſeems to confirm the preſent opinion more ſtrongly than any other is, that the Henharriers are all nearly of the ſame ſize, and that no one inſtance has been given of that bird proving a female by diſſection;

tion; thoſe we have diſſected have always proved males.

Much might be extracted from various authors on the ſubject; but nothing appears to elucidate the matter more than has been here mentioned. The ſubject has, however, been more largely treated of by Dr. Latham in the *Supplement* to his *Synopſis*. To that work therefore we refer our curious readers.

HERON. A genus of birds, the characters of which are,

Bill long, ſtrong, ſharp-pointed.
Noſtrils linear.
Tongue pointed.
Toes connected by a membrane as far as the firſt joint; in ſome of the ſpecies the middle claw is pectinated.

HERON-AFRICAN.

Ardea caſpica. *Ind. Orn.* ii. p. 698. 73.
African Heron. *Lath. Syn. Sup.* p. 237. *Lewin, Br. Birds*, iv. t. 151. *Walc. Syn.* ii. t. 131.

This bird is ſmaller than the Common Heron; length three feet. Bill duſky yellow, blackiſh at the point, ſeven inches long.

The head and greater part of the neck are pale ferruginous; chin and throat white; the feathers on

on the top of the head long, black, and form a ſort of a creſt; from the head a liſt of black runs down the back of the neck for two thirds its length; from the eye on each ſide another liſt continues down to the breaſt; on the lower part of the neck the feathers are long, looſe, and of a deep aſh-colour; the breaſt ferruginous cheſnut; back very deep aſh-colour; quills and tail black; the lower feathers on the rump like thoſe on the fore part of the neck, but mixed with ferruginous; belly pale ferruginous aſh-colour; legs dull yellow; the fore part of them, the toes, and claws, black.

Not more than two of this ſpecies is mentioned to have been met with in this country, one of which is ſtated to have been ſhot in Aſhdown Park, near Lambourn, Berks, now in the Leverian muſeum.

Said to inhabit Africa and Aſia.

HERON-COMMON.

Ardea major. *Lyn. Syſt.* i. p. 236. 12. *Gmel. Syſt.* ii. p. 627. *Raii Syn.* p. 98. A. 1. *Will.* p. 203. t. 49. *Ind. Orn.* ii. p. 691. 54.

Ardea criſtata. *Briſ.* v. p. 396. 2. t. 35.—*Ib.* 8vo. ii. p. 313.

Le Heron huppé. *Buf.* vii. p. 342. & t. 19. (fem.)

Creſted Heron. *Albin.* i. t. 67.—*Ib.* iii. t. 78.

Common Heron. *Br. Zool.* N° 173. t. 61.—*Ib.* fol. 116. t. A. *Arct. Zool.* ii. N° 343. *Lath Syn.* v. p. 83. 50. *Lewin, Br. Birds,* iv. t. 118

Wal

Walc. Syn. ii. t. 129. *Pult. Cat. Dorſet.* p. 14.

Ardea cinerea. *Lin. Syſt.* i. p. 236. 11. (fem.) *Don. Br. Birds*, iv. t. 73.

PROVINCIAL.

Heronſhaw. Crane.

The weight of this ſpecies is about three pounds and a half; length, to the end of the tail, about three feet four inches.

Bill near ſix inches long, duſky; at the baſe of the under mandible yellow; irides bright yellow; round the eye the ſkin is bare and greeniſh. The forehead and crown of the head white; on the hind part of the head the feathers are of a gloſſy black, very long, and form a looſe pendant creſt; the neck is whitiſh, marked on the fore part with a double row of black ſpots; ſcapulars grey and white, which, with thoſe on the lower part of the neck, are long and looſe; wing coverts bluiſh grey; the baſtard wings and greater quill-feathers black; the ſides of the body, from the breaſt to below the thighs, black; middle of the breaſt and belly white; thighs white, tinged with ruſt-colour; the tail is ſhort, of a bluiſh aſh-colour; legs very long, of a dull greeniſh colour; the middle claw ſerrated.

The female wants the black and white feathers on the head, inſtead of which that part is bluiſh grey, and not much elongated into a creſt, as in the

the male; the feathers on the breaſt and ſcapulars are not ſo long and looſe.

The young male birds are like the female for ſome time.

Linnæus has made the two ſexes diſtinct ſpecies; others were long of the ſame opinion, but later obſervations have corrected the miſtake.

This bird is found in moſt parts of the known world; is common in England.

It is a great deſtroyer of fiſh, both ſea and freſh-water; enabled, by the great length of legs, to wade into ſome depth of water, where it ſtands motionleſs till ſome of the finny tribe approach, when in an inſtant it darts its bill into them. Its digeſtion is as quick as its appetite is voracious, and of courſe commits vaſt devaſtation in ponds and ſhallow waters.

They feed frequently by moonlight, at which time they become tolerably fat, being not only leſs diſturbed in the night, but it has been obſerved that fiſh then come into the ſhoaler waters. Beſides fiſh, frogs and toads have been found in their ſtomachs.

In the breeding ſeaſon they congregate, and make their neſts very near each other; Mr. Pennant mentions having ſeen eighty neſts on one tree. We once ſaw a heronry on a ſmall iſland in a lake in the north of Scotland, whereon there was only one ſcrubby oak-tree, which not being ſufficient to contain all the neſts, many were placed on the ground.

The

The neſt is large and flat, made of ſticks lined with wool and other ſoft materials. The eggs are four or five in number, of a greeniſh blue, about the ſize of thoſe of the Duck.

Heronries were held ſacred in the days of falconry, being eſteemed for the diverſion it gave. Some are yet to be ſeen in ſeveral parts of the kingdom.

HERON-DWARF. Vide *Egret.*

HERON-GARDENIAN.

Ardea Gardeni. *Gmel. Syſt.* ii. p. 645. *Ind. Orn.* ii. p. 685. N° 32. *Lin. Tranſ.* v. p. 276.

Botaurus nævius. *Briſ.* v. p. 462. 31.—*Ib.* 8vo. ii. p. 332.

Butor tacheté, *ou* Pouacre. *Buf.* vii. p. 427.

Gardenian Heron. *Arct. Zool.* ii. N° 355. *Lath. Syn.* v. p. 71. 32.

Spotted Heron. *Lath. Syn.* v. p. 70. 31.

An inſtance of this ſpecies having been killed in England enables us to add it to the liſt of Britiſh birds. It was ſhot in Oxfordſhire by John Horatio Dickins, Eſq. and communicated to the Linnean Society April 3d, 1798.

In ſize it ſcarce exceeds that of the Rook; length about twenty inches.

Bill duſky, beneath greeniſh yellow; lore greeniſh; the plumage above is duſky, ſpotted and ſtreaked with white, except the lower part of the back,

back, which is plain; the head, neck, breaſt, and belly, whitiſh, ſtreaked longitudinally with fine ſhort lines of black or duſky; the leſſer coverts marked with yellowiſh ſpots; greater coverts with a white ſpot at the tip of each feather, forming two rows acroſs the wing; the greater quills edged with dull white, tips the ſame; tail duſky; legs of a dirty yellow.

This ſpecies ſeems to vary ſomewhat in plumage, either from ſex or age. In ſome the duſky parts are more inclined to brown; and the under parts, which in ſome are whitiſh, in others are of a light brown.

Inhabits South Carolina and Cayenne; frequents ponds, marſhes, and rivers, in the interior parts remote from the ſea; and feed on frogs and fiſh.

HERON-GREAT-WHITE.

Ardea alba. *Lin. Syſt.* i. p. 239. 24. *Gmel. Syſt.* ii. p. 639. *Ind. Orn.* ii. p. 695. 65.

Ardea alba major. *Raii Syn.* p. 99. A. 4. *Will.* p. 205. t. 49.

Ardea candida. *Briſ.* v. p. 428. 15.—*Ib.* 8vo. ii. p. 322.

Le Heron blanc. *Buf.* vii. p. 365.

Great White Heron. *Br. Zool.* ii. p. 175. t. 62.—*Ib.* fol. 117. *Arct. Zool.* ii. N° 234.—*Ib. Sup.* p. 66. *Will. Angl.* p. 279. t. 49. *Lath. Syn.* v. p. 91. 60. *Lewin, Br. Birds*, iv. t. 150.

The

The weight is about two pounds and a half; length three feet four inches. The bill is near ſix inches long, and yellow; the bare ſkin between that and the eye green; irides and orbits pale yellow. The whole plumage is of a pure white; legs black; the middle claw ſerrated.

One of theſe birds (we are informed by Dr. Latham) was killed in Cumberland not many years ago.

It is ſaid to have been more plentiful in England formerly; is common on the European continent. In America it is found in the ſouthern parts in winter; breeds more northward. We have ſeen them about New York in the ſummer, as well as in the lower lands on the banks of ſmall ſtreams in Statin Iſland.

HERON-LESSER-ASH-COLOURED. Vide *Heron-night.*

HERON-LESSER-WHITE. Vide *Egret.*

HERON-NIGHT.

Ardea Nycticorax. *Lin. Syſt.* i. p. 235. 9. *Gmel. Syſt.* ii. p. 624. *Raii Syn.* p. 99. 3. *Will.* p. 204. t. 49. *Ind. Orn.* ii. p. 6 8. 13. *Briſ.* v. p. 493. 45. t. 39.—*Ib.* 8vo. ii. p. 341.

Le Bihoreau. *Buf.* vii. p. 435. t. 12.

Night Heron, *or* Night Raven. *Arct. Zool.* ii. N° 356. *Will. Angl.* p. 279. t. 49. *Albin.* ii. t. 67. *Lath. Syn.* v. p. 52.—*Ib. Sup.* p. 234. *Lewin, Br. Birds,* iv. t. 145. *Walc. Syn.* ii. t. 126.

HER

PROVINCIAL.

Leſſer aſh-coloured Heron.

FEMALE.

Ardea Griſea. *Lin. Syſt.* i. p. 239. 21. *Gmel. Syſt.* ii. p. 625. *Briſ.* v. p. 412. 9. t. 36. f. 1. —*Ib.* 8vo. ii. p. 317.

The length of this ſpecies is twenty inches.

The bill is ſtrong, three inches and three quarters long, black, with the baſe yellowiſh; irides orange; lore, and round the eyes, green. The crown of the head is greeniſh black, extending a little way down the back of the neck; on the back of the head are three very narrow white feathers near ſix inches long, with their tips duſky; the hind part of the neck and ſides are aſh-colour; upper part of the back dull green; the lower part, rump, wings, and tail, pale aſh-colour; the forehead and moſt of the body white; legs yellowiſh green; claws duſky.

The female is gloſſy brown on the head; the upper parts of the body the ſame, but tinged with grey; the hind part of the neck paleſt; the lower part of the back and rump almoſt grey; over the eye is a whitiſh ſtreak; chin white; fore part of the neck grey, ſtreaked with yellowiſh down the ſhaft of each feather; the reſt of the under parts grey, becoming white at the vent; the wings are greyiſh brown, ſtreaked with yellowiſh white; ſome of the

greater

greater coverts tipped with white; quill-feathers cinereous grey, moſtly tipped with white; tail nearly the ſame; legs greyiſh brown.

This bird is not uncommon in Europe, though one inſtance only of its being found in England is on record; this was ſhot near London, in the month of May 1782, now in the Leverian muſeum.

It is ſaid to be common in Ruſſia, particularly on the river Don, where it builds on trees; is found alſo in ſome parts of America. It is alſo ſaid to lay three or four white eggs.

HERON-SHAW. Vide *Heron-common.*

HERON-SPOTTED. Vide *Heron Gardenian.*

HEW-HOLE. Vide *Woodpecker-green.*

HICK-WALL. Vide *Woodpecker-leaſt-ſpotted.*

HIGH-HOE. Vide *Woodpecker-green.*

HOBBY.

Falco ſubbuteo. *Lin. Syſt.* i. p. 127. 14. *Gmel. Syſt.* i. p. 283. *Raii Syn.* p. 15. A. 14. *Will.* p. 49. t. 7. *Ind. Orn.* i. p. 47. 114.

Dendrofalco. *Briſ.* i. p. 375. 20.—*Ib.* 8vo. p. 109. *Raii Syn.* p. 14. 8. *Will.* p. 47.

Hobreau. *Buf.* i. p. 277. t. 17.

Hobby. *Br. Zool.* i. N° 61.—*Ib.* fol. p. 69. t. A. 9. *Arct. Zool.* ii. p. 227. C. *Albin.* i. t. 6. *Lath. Syn.* i. p. 103. 90.—*Ib. Sup.* p. 28. *Lewin, Br. Birds,* i. t. 21. *Pult. Cat. Dorſet.* p. 3. *Walc. Syn.* i. t. 21. *Don. Br. Birds,* iv. t. 91.

This ſpecies weighs about ſeven ounces; length twelve inches. Bill blue; cere and orbits yellow; irides duſky.

The head and upper parts of the body are of a dark duſky brown, almoſt black, daſhed with aſh-colour; the feathers margined with pale rufous-brown; over the eye a light ſtroke; beneath the eye a black patch, extending in a point from the under mandible down each ſide of the throat; chin and throat white, extending round each ſide of the neck, and partly encircles it, but is broken behind by duſky ſtreaks, and the white becomes more ferruginous as it inclines backward; the coverts of the wings like the back, but the feathers more ſlightly edged; quill-feathers duſky black, with oval ferruginous ſpots on the inner webs; the breaſt, belly, thighs, and under tail coverts, ferruginous, paleſt on the former, marked with duſky ſtreaks; the tail like the back; barred on the inner webs with ruſt-colour, except the middle feathers; tips whitiſh; legs yellow; claws black.

The female weighs about nine ounces, ſometimes more, and very much reſembles the male in plumage, but not ſo dark above, and the lighter parts beneath not ſo ferruginous.

The wings of this bird are long and pointed, but do not reach to the end of the tail when cloſed; the ſecond feather is longeſt.

This is a migrative ſpecies, at leaſt it has never been obſerved with us in winter, but leaves us th

latt

latter end of October, about the time the Merlin arrives in the ſouthern parts.

It builds in trees, and ſometimes takes poſſeſſion of a deſerted Crow's neſt; lays three or four eggs, which are ſaid to be white.

We have ſeen three young ones taken out of a neſt; theſe were not ſo dark-coloured as the old birds.

Small as this ſpecies of Falcon is, it is inferior to none in point of courage; will frequently pounce a partridge; but their favourite game ſeems to be the Lark, to which it is a great enemy, and is frequently taken in purſuit of them by the birdcatchers in their nets.

The Hobby was formerly trained for hawking, but more commonly uſed for taking Partridges and Larks with a net, which was termed daring; that is, the Hobby was caſt off, which ſo frightened the birds that they readily ſuffered a net to be drawn over them.

HONEY-BUZZARD.

Falco apivorus. *Lin. Syſt.* i. p. 130. 28. *Gmel. Syſt.* i. p. 267. *Raii Syn.* p. 16. 2. *Will.* p. 39. t. 3. *Ind. Orn.* i. p. 25. 52. *Briſ.* i. p. 410.

La Bondrée. *Buf.* i. p. 208.

Honey Buzzard. *Br. Zool.* i. N° 56.—*Ib.* fol. 67. t. A. 4. & A.* 4. *Arct. Zool.* ii. p. 224. I. *Will. Angl.* p. 78. t. 3. *Albin.* i. t. 2. *Lewin,*

HON

Br. Birds, i. t. 1. *Lath. Syn.* i. p. 52.—*Sup.* p. 14. *White, Selb.* p. 109. *Pult. Cat. Dorſet.* p. 2. *Walc. Syn.* i. t. 7.

PROVINCIAL.

Capped Buzzard.

This ſpecies of Falcon weighs about one pound ten ounces; length twenty-three inches; breadth four feet one inch. The bill and cere duſky; irides bright yellow.

The head is brown, tinged with aſh-colour; the feathers at the baſe of the bill ſmall and cloſe, without hairs, as in moſt of the Hawk tribe; all the other parts above are of a deep brown; the chin whitiſh; breaſt and belly light brown, marked with brown bars, tinged with ruſt-colour; the tail brown, with two broad bars of duſky brown, one of which is very near the end; legs ſtrong, and of a dull yellow.

The bird from which this deſcription is taken was killed in Lord Carnarvon's park at Highclere in Berkſhire, and preſented to us by that nobleman.

At the time it was ſhot it was ſkimming over a large piece of water; but whether it was male or female we have not noted, though we believe the latter.

The Honey Buzzard ſeems to vary conſiderably in their markings; that mentioned in the *Britiſh Zoology* was white beneath, with duſky ſtrokes pointing

pointing downwards; on the tail was three broad duſky bars, between which were two or three of the ſame colour but narrower.

Linnæus remarks, that the tail has but one cinereous bar, and the tip white.

Albin's bird has no bar on the tail; that deſcribed in Latham's *Synopſis* agrees pretty nearly with the above.

This appears to be a very rare ſpecies. Dr. Latham remarks, that in the many years he has been a collector, one ſpecimen only came to him freſh.

The name ſeems to have been given to this bird from its feeding on the larvæ of Waſps, and perhaps Bees, the firſt of which, Willughby informs us, he found in its neſt.

It builds in a tree, like the Common Buzzard. The egg is ſaid to be in colour like that of the Reſtril, dirty white, blotched all over with ruſt-colour. Mr. White mentions one egg only to be found in the neſt, and which contained the embryo young; ſays it is ſmaller, and not ſo round as that of the Buzzard, dotted at each end with ſmall red ſpots, and ſurrounded in the middle with a broad bloody zone.

HOOP. Vide *Finch-bul* and *Hoopoe.*

HOOPER. Vide *Swan-whiſtling.*

HOOPOE. A genus of birds, the characters of which are,

Bill long, ſlender, bending.

Noſtrils ſmall, placed near the baſe.

Toes, three before, one behind; the middle one connected at the baſe to the outmoſt.

HOOPOE-COMMON.

Upupa epops. *Lin. Syſt.* i. p. 183. 1. *Gmel. Syſt.* i. p. 466. *Raii Syn.* p. 48. A. 6. *Will.* p. 100. t. 24. *Ind. Orn.* i. p. 277. 1. *Briſ.* ii. p. 455. t. 43. f. 1.

La Huppe. *Buf.* vi. p. 439. t. 21.

Hoopoe. *Br. Zool.* i. N° 90. t. 39.—*Ib.* fol. 83. t. L. *Arct. Zool.* ii. p. 283. A. *Will. Angl.* p. 145. *Albin.* ii. t. 42, 43. *Edw.* t. 345. *Lewin, Br. Birds*, t. 54. *Lath. Syn.* ii. p. 687. 1.—*Ib. Sup.* p. 122. *Pult. Cat. Dorſet.* p. 7. *Walc. Syn.* i. t. 53. *Don. Br. Birds*, i. t. 9.

The weight of this beautiful bird is about three ounces; length twelve inches.

The bill is black, two inches and a half long, ſlender, and curved; irides hazel.

The crown of the head is furniſhed with a creſt compoſed of a double row of dull orange-coloured feathers tipped with black, lengthening from the forehead backwards, the longeſt of which is above two inches; the ſides of the head, neck, and breaſt, dull orange-colour daſhed with brown; upper part of the back browner; belly whitiſh; the greater quill-feathers are black, with a broad bar of white near the tips; on the ſecondaries the white decreaſes;

creaſes; thoſe next the body are marked with ſeveral black and white bars; the coverts are barred in the ſame manner, thoſe on the upper ridge of the wing like the neck; the rump is white; the tail conſiſts of ten black feathers, croſſed with a large ſemilunar bar of white; the exterior feather white on the outer margin, except at the tip; legs ſhort and black.

In ſome the breaſt is deſcribed to be white; in young birds this part is marked with narrow duſky lines.

The female reſembles the male.

This bird is only occaſionally met with in this country; every autumn perhaps produces a few; and inſtances have not been wanting to prove they have ſometimes bred with us. In the *Supplement* to the *General Synopſis* an account is given of a young one being ſhot in May. A pair is alſo mentioned to have begun a neſt in Hampſhire, but being diſturbed forſook it, and went elſewhere.

The neſt is ſaid to be made of bents, and lined with ſoft materials; the eggs, four in number, of bluiſh white, marked with pale brown ſpots. It builds in the hollow of a tree, and the neſt has been remarked to be extremely fetid; probably occaſioned by the fæces of the young, and not by the filthy food by which it has been ſuppoſed they feed their young.

Theſe birds have been ſeen in moſt parts of Great Britain, from Scotland to the moſt ſouthern parts,

parts, as we find on record; and we have known it killed in South Wales and in Devonſhire. With us they ſeem to prefer barren ſituations. Their food is inſects and worms. It is found plentiful in the deſerts of Ruſſia and Tartary; are ſeen in ſmall flocks at Gibraltar, in the month of March, on their paſſage north, ſuppoſed to come from Africa.

HORSEMAN-GREEN-LEGGED. Vide *Green-ſhank.*

HORSEMAN-RED-LEGGED. Vide *Gambet.*

HOWLET. Vide *Owl-white.*

HUCKMUCK-GROUND. Vide *Wren-yellow.*

HUCKMUCK-TREE. Vide *Titmouſe-long-tailed.*

I.

IBIS. A genus of birds, the characters of which are,

Bill long, incurved, thick at the baſe.
Face, and ſometimes the whole neck, bare of feathers.
Noſtrils linear.
Tongue ſhort.
Toes connected at the baſe by a membrane.

IBIS-GLOSSY.

Tantalus igneus. *Gmel. Syſt.* ii. p. 649. *Ind. Orn.* ii. p. 708. 16.

Gloſſy Ibis. *Lath. Syn.* v. p. 115. 14. *Walc. Syn.* ii. t. 132. *Levain, Br. Zool.* iv. t. 152. *Don. Br. Birds*, v. t. 118.

This is the only ſpecies that has ever been met with in England, and of that only one or two inſtances. Dr. Latham mentions one in the Leverian muſeum which was ſhot in Cornwall. As we have never had an opportunity of ſeeing a freſh ſpecimen, ſhall take the liberty of borrowing from the above author's Synopſis.

Length almoſt two feet. Bill five inches, ſmooth, roundiſh, much bent, green, fading to olive when dead; eyelids brown; irides olive; eyes placed in a white ſpace; under the chin a ſmall dilatable pouch; head and neck black; the feathers fringed with white; the reſt of the body variegated with blackiſh blue-green and vinaceous, and in general very gloſſy; hence the bird, in flying, appears gilded when the ſun ſhines upon it; quills green-gold, and when cloſed reach the end of the tail; wing coverts next the body reddiſh and blue mixed; the next ſeries black, red, and green; the laſt, and quills, green-gold; tail the ſame, gloſſed in different lights with red and violet; legs very long, of a bright green; claws crooked, black.

This inhabits Ruſſia, chiefly the ſhores of the Don,

Don, and about the Choper; lives on fish and insects; flies in flocks, and builds in trees.

IMBER. Vide *Diver Imber.*
IMMER. Vide *Diver Imber.*

J.

JACKDAW.

Corvus monedula. *Lin. Syst.* i. p. 156. 6. *Gmel. Syst.* i. p. 367. *Raii Syn.* p. 40. t. 5. *Will.* p. 85. t. 19. *Ind. Orn.* i. p. 154. 11. *Brisson.* ii. p. 24. 6.—*Ib.* 8vo. i. p. 160.

Le Choucas. *Buf.* iii. p. 69.

Jackdaw. *Br. Zool.* i. N° 81. t. 34.—*Ib.* fol. p. 78. *Arct. Zool.* ii. p. 251. C. *Will. Angl.* p. 125. t. 9. *Albin.* i. t. 14. *Lewin, Br. Birds,* i. t. 37. *Lath. Syn.* i. p. 378. 9.—*Ib. Sup.* p. 78. *Pult. Cat. Dorset.* p. 5. *Walc. Syn.* i. t. 36.

PROVINCIAL.

Chough. Daw.

This species of Crow weighs about nine ounces; length near thirteen inches.

Bill black; irides light grey.

The forehead is black; the hind part of the head and

and back of the neck cinereous-grey; upper parts of the body black, ſlightly gloſſed with blue; tail the ſame; the under parts duſky black; legs black.

This very common bird frequents old towers, ruined buildings, and high cliffs, where it builds, as well as in holes of trees. The neſt is made of ſticks, and lined with wool and other ſoft materials; the eggs are five or ſix in number; bluiſh ſpotted with black.

Theſe birds are gregarious, and frequently flock together with Rooks; feed in the ſame manner on grain and inſects; are fond of cherries, and will devour carrion in ſevere weather.

It is ſeen to frequently perch on the back of ſheep, not only to rob that animal of its wool to line its neſt, but alſo to pick out the ticks with which it is infeſted.

It is a very docile, tractable, and miſchievous bird, eaſily made tame, and may be taught to talk.

Some inſtances are mentioned of its breeding in rabbit-holes.

Several varieties of this ſpecies are given by different authors; ſome entirely black, without the grey on the head and neck; others quite white, or mixed black and white.

Is found in Denmark, France, and Germany; alſo in Ruſſia and the weſt of Siberia; but in moſt of theſe places are more or leſs migratory.

JAY.

Corvus glandarius. *Lin. Syſt.* i. p. 156. 7. *Gmel. Syſt.* i. p. 278. *Raii Syn.* p. 41. A. 2. *Will.* p. 88. t. 19. *Ind. Orn.* i. p. 157. 18.

Garrulus. *Briſ.* ii. p. 47. 1.—*Ib.* 8vo. i. p. 168.

Geay. *Buf.* iii. p. 107. t. 8.

Jay. *Br. Zool.* i. N° 79.—*Ib.* fol. t. D. *Arct. Zool.* ii. p. 252. E. *Will. Angl.* p. 130. t. 19. *Albin.* i. t. 16. *Hayes, Br. Birds*, t. 7. *Lewin, Br. Birds*, t. 38. *Lath. Syn.* i. p. 384. 19.—*Ib. Sup.* p. 79. *Pult. Cat. Dorſet.* p. 5. *Walc. Syn.* i. t. 37. *Don. Br. Birds*, i. t. 2.—*Ib.* ii. t. 34.

This beautiful ſpecies of Crow weighs ſeven ounces; length near thirteen inches.

The bill is duſky; irides whitiſh.

The forehead is white, ſtreaked with black; chin whitiſh; a broad ſtreak of black runs from the corners of the mouth under each eye, pointing downwards; on the crown of the head the feathers are long, and erectable in form of a creſt; the hind part of the head, the ſides, neck, breaſt, back, and ſcapulars, vinaceous; the leſſer wing coverts inclining to bay; the greater coverts are elegantly barred with a rich blue and black alternately, the reſt black; the greater quill-feathers duſky; the exterior webs aſh-colour, except the firſt; ſix of the ſecondary quills are black, white on their exterior webs near the baſe, and tinged with blue; the two next entirely black; thoſe neareſt the body bay, tipped with black; rump, upper and under tail coverts, white; the tail black; legs brown.

JAY

This bird is found in tolerable plenty in moſt of the wooded parts of this kingdom, but are never gregarious.

It makes a neſt moſt commonly in high coppice wood or hedges; ſometimes againſt the ſide of a ſcrubby tree; it is formed of ſticks, lined with fibrous roots; lays five or ſix eggs of a light brown colour, not very unlike thoſe of the Partridge, but ſmaller, and obſcurely marked with a darker ſhade of brown.

The Jay is a cunning, crafty bird; is a great devourer of fruit and grain, and ſeems particularly fond of cherries and peas; will frequently plunder the ſmaller birds neſts of their eggs and young, and ſometimes pounce the old birds, on which it preys, as well as on mice.

Its common notes are various, but harſh; will ſome time in the ſpring utter a ſort of ſong in a ſoft and pleaſing manner, but ſo low as not to be heard at any diſtance; and at intervals introduce the bleating of a Lamb, mewing of a Cat, the note of a Kite or Buzzard, hooting of an Owl, and even the neighing of a Horſe.

Theſe imitations are ſo exact, even in a natural, wild ſtate, that we have frequently been deceived.

In the autumn it feeds on acorns, which it has been ſaid to hoard for the winter; but this is certainly a miſtake; ſuch hoards found in our woods belong either to the Squirrel or ſome ſpecies of Mouſe.

JUDCOCK, or JUDDOCK. Vide *Snipe-jack*.

K.

KAE-CORNWALL. Vide *Crow-red-legged.*

KASTRIL, or KISTRIL. Vide *Keſtrel.*

KATE. Vide *Brambling.*

KESTREL.

Falco tinnunculus. *Lin. Syſt.* p. 127. 16. *Gmel. Syſt.* i. p. 278. *Raii Syn.* p. 16. 16.—*Ib.* p. 180. 2. *Will.* p. 50. t. 5. *Ind. Orn.* i. p. 41. 98.

La Creſſerelle. *Buf.* i. p. 280. t. 18. *Briſ.* i. p. 393. 27.

Accipiter alaudarius. *Briſ.* i. p. 379. 22.—*Ib.* 8vo. p. 100. *Gmel. Syſt.* i. p. 279.

Keſtrel, Stannel, *or* Windhover. *Will. Angl.* p. 84. t. 5.

Keſtrel. *Br. Zool.* i. N° 60.—*Ib.* fol. p. 68. t. A *Arct. Zool.* ii. p. 226. N. *Hayes, Br. Birds,* t. 4. *Albin.* iii. t. 5. & 7. *Lewin, Br. Birds,* i. t. 19. M. & F. *Lath. Syn.* i. p. 94. 79.—*Sup.* p. 25. *Walc. Syn.* i. t. 19. *Pult. Cat. Dorſet.* p. 3. *Don. Br. Birds,* iii. t. 51. (M.)—*Ib.* 63. (F.)

PROVINCIAL.

Kaſtril, or Kiſtril. Steingal, or Stonegall.

The

The male of this ſpecies of Falcon weighs about ſeven ounces; length thirteen inches. Bill lead-colour; cere yellow; irides duſky and large.

The crown of the head is of a fine cinereous-grey; throat whitiſh; under the eye a broad duſky ſtreak pointing downwards; the back, ſcapulars, and wing coverts are of a fine red-brown, ſpotted and barred with black; the under parts light ferruginous, ſpotted with black; thighs and vent generally plain; greater quill-feathers black, very ſlightly tipped whitiſh; the wing pointed; the ſecond feather the longeſt; tail fine cinereous-grey, with a broad black bar near the end; tip white; legs yellow.

The female is conſiderably larger and is diſtinguiſhed from the other ſex by the head and tail being of the ſame colour as the back, which is not ſo bright a red-brown as the male; the under parts are alſo lighter, and the black ſpots not ſo diſtinct, but more in ſtreaks of duſky; the tail is marked with tranſverſe duſky bars, with a broad one near the end.

This beautiful ſpecies of Hawk feeds principally on mice, in ſearch of which it is frequently ſeen hovering in the air, and quite ſtationary for a great length of time.

We never have ſeen the Keſtrel in purſuit of ſmall birds, nor have we ever found feathers in the ſtomach, but chiefly the fur of mice, and the exuviæ

of beetles; but no doubt it will ſometimes prey on ſmall birds, as it is occaſionally taken by bird-catchers in the act of pouncing their call-birds.

This is one of our moſt common ſpecies, eſpecially in the more rocky ſituations and high cliffs on our coaſts, where they breed.

The neſt is made of ſticks, and lined with wool and other ſoft materials; ſometimes they build in trees, or content themſelves with the deſerted neſt of a Crow or Magpie.

The number of eggs are four or five, of a dirty white, blotched over with ruſt-colour of various ſhades; ſometimes wholly covered with a deep ruſty red: theſe are rather inferior in ſize to thoſe of the Sparrow-hawk.

The young males reſemble the female in plumage till after the winter of the firſt year, when they aſſume the grey head and tail.

KIDDAW. Vide *Gaillemot-fooliſh.*

KILLEGREW. Vide *Crow red-legged.*

KINGFISHER. A genus of birds, the characters of which are,

Bill long, ſtrait, ſharp-pointed.
Tongue ſhort, broad, ſharp-pointed.
Legs ſhort.
Toes, three forward, one backward; three lower joints of the middle toe cloſely joined to the outmoſt.

KINGFISHER-

KINGFISHER-COMMON.

Alcedo Iſpida. *Lin. Syſt.* i. p. 179. 3. *Gmel. Syſt.* p. 448. *Raii Syn.* p. 48. A. 1. *Will.* p. 101. t. 24. *Ind. Orn.* i. p. 252. 20. *Briſ.* iv. p. 471. 1.—*Ib.* 8vo. ii. p. 176.

Martin pêcheur. *Buf.* vii. p. 164. t. 9.

Kingfiſher. *Br. Zool.* i. N° 88. t. 38.—*Ib.* fol. 82. t. 1. *Arct. Zool.* ii. p. 280. A. *Will. Angl.* p. 146. t. 24. *Albin.* i. t. 54. *Lewin, Br. Birds,* ii. t. 52. *Lath. Syn.* ii. p. 626.—*Ib. Sup.* p. 115. *Walc. Syn.* i. t. 52. *Pult. Cat. Dorſet.* p. 6. *Don. Br. Birds,* iv. t. 100.

The weight of this beautiful bird is an ounce and a half; length ſeven inches. The bill is two inches long and black, tinged with orange at the baſe of the lower mandible; irides hazel.

The crown of the head is of a dark changeable green-blue, with numerous ſmall tranſverſe bars of bright azure; from the upper mandible to the eye a duſky ſtreak; the ſides of the forehead rufous; behind the eye a broad ſtripe of red-orange, at the lower angle of which, on each ſide, commences a yellowiſh white ſtripe, which almoſt meet on the back part of the neck; from the baſe of the lower mandible ſprings a blue ſtreak, which runs to the ſide of the neck; the throat is buff-coloured; the under parts of a dull orange; darkeſt on the breaſt; the wing coverts like the upper part of the head, but not ſo much ſpotted; down the middle of the

back, the rump, and upper tail coverts, fine bright azure; tail deep blue; legs red-orange.

The bill of the female is not ſo long as in the other ſex.

This bird is found moſt frequently about clear running ſtreams and rivers, in the banks of which it generally takes poſſeſſion of a rat's-hole to depoſit its eggs. The many curious accounts which have been given of the neſt of this bird induced us to take ſome pains to diſcover the fact. The reſult of our reſearches are, that the hole choſen to breed in is always aſcending, and generally two or three feet in the bank; at the end is ſcooped a hollow, at the bottom of which is a quantity of ſmall fiſh-bones, nearly half an inch thick, mixed in with the earth. This is undoubtedly the caſtings of the parent birds, and not the young, for we have found it even before they have eggs, and have every reaſon to believe both male and female go to that ſpot for no other purpoſe than to eject this matter for ſome time before the female begins to lay, and that they dry it by the heat of their bodies, as they are frequently known to continue in the hole for hours long before they have eggs. On this diſgorged matter the female lays to the number of ſeven eggs, which are perfectly white and tranſparent, of a ſhort oval form, weighing about one dram. The hole in which they breed is by no means fouled by the caſtings; but before the young are able to fly it becomes extremely fetid by the fæces of the brood,

which

which is of a watery nature, and cannot be carried away by the parent birds, as is common with moſt of the ſmaller ſpecies. In defect of which inſtinct has taught them to have the entrance to their habitation aſcending, by which means the filthy matter runs off, and may frequently be ſeen on the outſide. We never could obſerve the old birds with any thing in their bills when they went in to feed their young; from which it may be concluded they eject from their ſtomach for that purpoſe.

Small fiſh, ſuch as Baniſtickles and Minnows, ſeem to be their principal food. But it is a miſtaken notion that theſe birds ſuſpend themſelves on wing, and dart on their prey like the Oſprey; the fact is, they ſit patiently on a bough over the water, and when a ſmall fiſh comes near the ſurface they dart on it, and ſeize it with their bill. The Kingfiſher is rarely ſeen about the rocky rapid waters where the Water Ouzel chiefly reſorts, but is frequently found about fiſh-ponds.

It flies with great rapidity notwithſtanding its wings are very ſhort; but the motion of the wings are ſo very quick as ſcarce to be perceptible.

When the young are nearly full feathered they are extremely voracious; the old birds not being capable of ſupplying them with food ſufficient to ſatisfy the calls of hunger, they are continually chirping, and may be diſcovered by their noiſe.

The ſuſpenſion of this bird by a thread, under the notion of its breaſt always turning to the north,

is as fabulous as that it will preſerve woollen cloth from the moth.

The poetic fiction of the older authors, with reſpect to the Halcyon, are too copious to be ſtated in this place. Thoſe who wiſh to ſee the marvellous accounts given by Ariſtotle, Virgil, Ovid, and others, may peruſe Mr. Pennant's hiſtory of this bird, where the eſſence is extracted from thoſe authors. The ſailors of the preſent day do not find it has the power to calm the ſtorm, or huſh the wind.

The Kingfiſher is found throughout Europe, Aſia, and Africa; the only ſpecies out of nearly fifty that is found in the colder parts.

KITE.

Falco Milvus. *Lin. Syſt.* i. p. 126. 12. *Gmel. Syſt.* i. p. 261. *Raii Syn.* p. 17. A. 6. *Will.* p. 41. t. 6. *Ind. Orn.* i. p. 20. 37.

Milvus regalis. *Brif.* i. p. 414. 35. t. 33.—*Ib.* 8vo. p. 118.

Milan royal. *Buf.* i. p. 197. t. 7.

Kite. *Br. Zool.* i. N° 53.—*Ib.* fol. t. A. 2. *Arct. Zool.* ii. p. 223. H. *Will. Angl.* p. 74. t. 6. *Albin.* i. t. 4. *Hayes, Br. Birds,* i. t. 5. *Lewin, Br. Birds,* i. t. 10. *Lath. Syn.* i. p. 61. 43.—*Sup.* p. 17. *Pult. Cat. Dorſet.* p. 3. *Walc. Syn.* i. t. 10. *Don. Br. Birds,* ii. t. 47.

PROVINCIAL.

Glead. Puttock.

This

KIT

This ſpecies of Falcon weighs about two pounds ſix ounces; length two feet two inches. The bill yellowiſh, point duſky; cere yellow; irides the ſame, but of a light colour. Head grey, ſtreaked down the middle of each feather with duſky; back and wing coverts duſky, edged with ferruginous; the under parts more or leſs ferruginous, ſtreaked with duſky, lighteſt on the breaſt; quill-feathers duſky black; from the fifth to the tenth daſhed with aſh-colour, with a few duſky bars, and white at the baſe and on the inner webs; the reſt are duſky, with obſcure bars; the tail is of a bright ferruginous colour, the two exterior feathers duſky on the outer webs, the firſt barred on the inner web the ſame; the legs are yellow; claws black.

The tail of this bird at once diſtinguiſhes it from all others of the genus, being much forked; the exterior feathers are twelve inches in length.

The female is ſomewhat larger, meaſuring in length two feet four inches; breadth five feet ſix inches; in colour much reſembling the other ſex, but in general not ſo ferruginous.

The Kite chiefly inhabits wooded ſituations, but frequently changes its abode in the winter, though it never wholly quits this country.

It makes a neſt early in the ſpring in a fork of ſome large tree, compoſed of ſticks, and lined with wool, the inner bark of ſome tree, hair, and other ſoft materials, ſuch as bits of cloth or rags.

The eggs are generally three in number, rarely

four; theſe are rather larger than thoſe of a Hen, of a dirty white, with a few ruſty ſpots at the larger end; ſometimes quite plain; their weight is nearly two ounces.

This bird, from its great length of wings and tail, is capable of ſupporting itſelf in the air with very little motion, and for a great continuance, but is ſlow in flight; its depredations therefore are confined to ſuch animals as are found on the ground, young Rabbits, Hares, and game of all kinds, poultry, and young birds incapable of flying.

It will alſo deſtroy young Lambs, and feed greedily on carrion; in defect of theſe it readily eats mice, worms, and inſects, and even ſnakes, the bones of which we have taken from the neſt.

It frequently reſorts to the environs of towns to feed on offal, and is ſeen to ſweep ſuch matter from the ſurface of water with great dexterity.

Is ſaid to be met with in Sweden, and even as far as Guinea and Senegal.

KITTIWAKE.

Larus Riſſa. *Lin. Syſt.* i. p. 224. 1. *Gme Syſt.* ii. p. 598.

Larus tridactylus. *Ind. Orn.* ii. p. 817. 11.

Le Goeland cendré. *Briſ.* vi. t. 14?

Kittiwake. *Br. Zool.* ii. N° 250. t. 89. *A Zool.* ii. N° 456. *Lath. Syn.* vi. p. 393. *Lewin, Br. Birds*, vi. t. 213. *Pult. Cat. Do* p. 18. *Walc. Syn.* i. t. 108.

KIT

This ſpecies of Gull is about fourteen or fifteen inches in length; weight ſeven or eight ounces. The bill is greeniſh yellow; irides duſky; inſide of the mouth deep orange. Head, neck, breaſt, and all beneath, white; back and wings cinereous grey; the firſt quill-feather has the exterior web black, and the four or five next are tipped the ſame; the tail is pure white; legs duſky, with a knob inſtead of a back toe.

It ſometimes has a duſky ſpot behind the ear; but this muſt be conſidered as a mark of immaturity, and but another remove from the ſtate in which it has been made a diſtinct ſpecies, under the denomination of Tarrock, to the perfect bird here deſcribed.

Theſe birds had long been conſidered as diſtinct ſpecies, but are now, without doubt, brought together as one and the ſame. Under the head of Gull Tarrock the ſynonyms of that bird will be found, with references as uſual to the various authors who had conſidered it as a different bird.

The Kittiwake is ſaid to inhabit and breed on the cliffs about Flamborough Head, the Baſs Iſle, the vaſt rocks near the Caſtle of Slains in the county of Aberdeen, and on Prieſtholm Iſle. In the firſt of theſe places it is called Petrel.

The Tarrock, or immatured birds, are always more or leſs found in company with them, and no doubt are two or three years arriving to their full plumage.

It

It ſeems to be a local ſpecies with us, rarely appearing in the more ſouthern parts. Is found in the arctic regions; not uncommon in Iceland and Greenland; has been met with at Kamtſchatka.

KNOT.

Tringa Canutus. *Lin. Syſt.* i. p. 251. 15. *Gmel. Syſt.* ii. p. 679. *Raii Syn.* p. 108. A. 5. *Will.* p. 224. t. 56. *Ind. Orn.* ii. p. 738. 44. *Briſ.* v. p. 258. 21.—*Ib.* 8vo, ii. p. 276.

Le Canut. *Buf.* viii. p. 142.

Knot. *Br. Zool.* ii. N° 193.—*Ib.* fol. 123. t. E. 2. f. 1. *Arct. Zool.* ii. N° 384. *Will. Angl.* p. 302. *Edw.* t. 276. *Lath. Syn.* v. p. 187. 36. *Lewin. B. Birds*, v. t. 178.

The weight of this ſpecies of Sandpiper is four ounces and a half; length about nine inches. Bill duſky brown, an inch and a quarter long; irides hazel.

The top of the head, neck, back, and wings, aſh-colour; from the bill to the eye a duſky ſtreak; over the eye a white one; coverts of the wings edged with white; the lower order deeply tipped and margined, forming a bar acroſs the wing; greater quills duſky, with white ſhafts; the rump and tail coverts white and duſky, tranſverſely marked in curved lines; the under parts, from throat to vent, white; the neck and breaſt ſtreaked with brown; the ſides and thighs croſſed with duſky lines;

lines; tail aſh-colour, the outer feather whitiſh; legs bluiſh aſh-colour.

In ſome the forehead, chin, and lower part of the neck, is brown, inclining to aſh-colour; back and ſcapulars deep brown, edged with aſh-colour. Theſe and other little variations in plumage are common in moſt of this genus of birds.

This ſpecies is chiefly confined to the fens of Lincolnſhire, the Iſle of Ely, and a few other places. In the firſt place they are taken in great numbers on the coaſt by means of nets, and when fattened are eſteemed equal to the Ruffs. It ſhould ſeem they firſt appear in the month of Auguſt, as from that time to November is the ſeaſon of catching them; but are ſaid generally to diſappear with the firſt froſt.

We have received from the coaſts of Devonſhire and Cornwall, in Auguſt, a ſpecies of Sandpiper in every reſpect like this deſcribed, except that the feathers on the back and wing coverts are margined with a ſmall ſemicircular line of black; the extreme edges light cinereous-brown.

This we have conſidered as the young of the Aſh-coloured Sandpiper, and cannot help expreſſing doubts whether the Knot may not hereafter prove to be that bird in one ſtate of its plumage, probably the moſt perfect.

Mr. Pennant and Dr. Latham both deſcribe their Aſh-coloured Sandpiper to have the back and wing coverts varied with concentric ſemicircles of black,

aſh-colour,

ash-colour, and white, which brings it near to the bird above mentioned. There is no genus of birds more obscure than the Sandpipers; and as few of them remain with us the whole year, it will in all probability be no inconsiderable time before we shall be able to clear up the many doubts respecting them. We cannot, however, help expressing our opinion that the species are unnecessarily multiplied.

The Knot is said to be found in various parts of Europe; has been met with also in Asia and America. It should seem they breed with us from Mr. Lewis's account, who says they appear with the Ruffs, and has given a figure of its egg, which is pale ferruginous, marked with spots and streaks of rust-colour and cinereous; in size rather inferior to that of the Lapwings.

L.

LANNER.

Falco lanarius. *Lin. Syst.* i. p. 129. 24. *Gmel. Syst.* i. p. 276. *Raii Syn.* p. 15. 13. *Will.* p. 48. *Ind. Orn.* i. p. 38. 92. *Brisson.* i. p. 363. 16.—*Ib.* 8vo. p. 105.

Le Lanier. *Buff.* i. p. 243.

Lanneret. *Albin.* ii. t. 7.

Lanner,

Lanner. *Br. Zool.* i. N° 51. t. 23. *Arct. Zool.* ii. p. 125. K. *Will. Angl.* p. 82. *Lewin, Br. Birds*, i. t. 17. *Lath. Syn.* i. p. 86. 72.—*Ib. Sup.* p. 21. *Wale. Syn.* i. t. 16.

This ſpecies of Falcon is rather leſs than the Buzzard. Bill bluiſh; cere greeniſh-blue; irides yellow.

The crown of the head brown, mixed with yellowiſh white; the reſt of the head, upper part of the neck, and the body above, are brown, the feathers edged with a paler colour; over the eye a broad whitiſh line; beneath the eye a black ſtroke pointing downwards; the throat, breaſt, and belly, whitiſh, tinged with dull yellow on the two laſt parts, which, with the thighs and vent, are marked with brown ſtrokes; quills and tail duſky, with oval ferruginous ſpots on the inner webs; legs bluiſh, ſhort, and ſtrong; claws black.

This is a very bold bird, and was formerly uſed in falconry. Is rarely met with in England, but is ſaid to breed in Ireland.

Mr. Pennant ſpeaks of one being caught in a decoy in Lincolnſhire, purſuing ſome Wild Ducks under the nets.

The Lanner is found in many parts of Europe; inhabits Iceland and the Ferroe Iſles, Denmark and Sweden; frequent in the Tartarian deſerts; ſaid to build among the low trees and ſhrubs in the deſerts about Aſtrachan.

LAPWING.

Tringa Vanellus. *Lin. Syſt.* i. p. 248. 2. *Gmel. Syſt.* ii. p. 670. *Raii Syn.* p. 110. A. 1. *Will.* p. 228. t. 57. *Ind. Orn.* ii. p. 726. 2. *Briſ.* v. p. 94. 1. t. 8. f. 1.—*Ib.* 8vo. ii. p. 236.

Le Vanneau. *Buf.* viii. p. 48. t. 4.

Lapwing, *or* Baſtard Plover. *Br. Zool.* ii. N° 190.—*Ib.* fol. 122. t. C*. f. 1. *Arct. Zool.* ii. p. 480. D. *Albin.* i. t. 74. *Will. Angl.* p. 307. t. 57. *Hayes, Br. Birds,* t. 11. *Lath. Syn.* v. p. 161. 2. *Lewin, Br. Birds,* v. t. 167. *Pult. Cat. Dorſet.* p. 15. *Walc. Syn.* ii. t. 145. *Don. Br. Birds,* ii. t. 25.

PROVINCIAL.

Pewit. Green Plover.

This ſpecies of Sandpiper weighs between ſeven and eight ounces.

Bill black, an inch long; irides hazel.

The upper part of the head is black, gloſſed with green; on the back part the feathers are elongated into a creſt, ſome of which are above three inches in length and very narrow, reflecting a little at the ends; the ſides of the neck and round the eye is white; beneath the eye is a black ſtreak; the fore part of the neck and upper breaſt black; the hind part brown, intermixed with white; back and coverts of the wings brown-green, gloſſed with purple and blue; quills black, with a white ſpot on

the tips of the firſt four; the ſecondaries are white half way from their baſe; breaſt and belly white; vent and upper tail coverts pale ferruginous; the baſe half of the tail white; the ends black; legs dull orange.

The female is leſs brilliant in colour, and the creſt much ſhorter.

This bird is common in moſt parts of the kingdom; it breeds early in the ſpring upon heaths and upland ſituations, as well as in fens and moiſt fields, and not unfrequently in old fallow land.

It lays invariably four eggs on the bare ground; theſe are of an olivaceous brown, blotched with black, and, what is remarkable, the eggs are always placed in a quadrangular manner, touching each other at the ſmaller ends, by which they occupy the leaſt poſſible ſpace. This is common to all the Sandpiper, Plover, and Snipe genus hitherto noticed, the eggs of which are commonly four, much tapering to the ſmaller end.

The young make uſe of their legs as ſoon as they are hatched, but are not capable of flying till nearly full grown; they are led by the parent birds in ſearch of food, but are not fed by them. At this time the old birds uſe every art to entice an intruder from their young; will ſtrike at a dog, and then flutter along the ground as if wounded, to entice him from the place where the young are concealed, and are very clamorous.

In the autumn theſe birds congregate into large flocks,

flocks, and frequent marſhy places. At this time they are eſteemed not unſavoury food; the eggs are alſo conſidered as a delicacy, and are frequently brought to the markets of London for ſale. Its principal food is worms, ſlugs, and various inſects; when partly domeſticated and kept in gardens, will eat bread or meat. It runs faſt, and has a ſingular motion with the head, frequently putting its bill to the ground without picking any thing up.

Its note has given riſe to one of its common names, as it reſembles the word *pe-wit*; the name of Lapwing has alſo been given from the conſtant flapping of its wings in flight.

LARK. A genus of birds, the characters of which are,

Bill ſtrait, ſlender, ſharp-pointed, bending a little towards the end.

Noſtrils covered partly with feathers and briſtles.

Tongue cloven at the end.

Toes divided to their origin; claw of the back toe very long, and little crooked.

Their motion running, not hopping.

LARK-BUNTING. Vide *Bunting-common.*

LARK-FIELD.

Alauda minor. *Gmel. Syſt.* ii. p. 793. *Ind. Orn.* ii. p. 494. 8.

Leſſer

Leſſer Field Lark. *Will. Angl.* p. 207.
Field Lark. *Br. Zool.* N° 139. *Arct. Zool.* ii. p. 395. D. *Lath. Syn.* iv. p. 375. 6. *Lewin, Br. Birds,* iii. t. 92. *Walc. Syn.* ii. t. 192. *Pult. Cat. Dorſet.* p. 8.

PROVINCIAL.

Short-heeled Field Lark, and Meadow Lark.

The length of this ſpecies is ſix inches and a half; weight five drams, forty-five grains. Bill duſky above, whitiſh beneath; irides hazel. The colour of the plumage on the upper parts is a light yellowiſh brown, the middle of each feather duſky brown; the wing coverts tipped whitiſh; rump plain light brown; throat and breaſt ochraceous yellow, the latter ſtreaked with black; belly yellowiſh white; the tail feathers are ſomewhat pointed; the exterior one half white; the next ſlightly tipped the ſame; the legs yellowiſh brown; claws horn-colour; hind claws ſhort and hooked.

No bird has been more confounded than this ſpecies of Lark. It viſits this country in the ſpring, but is rarely ſeen till the beginning of May, and is moſt frequently miſtaken for the Tit-Lark, to which it bears great reſemblance in plumage and habits; but as a ſpecial mark of diſtinction the baſe of the bill in this is broader, and the hind claw is much ſhorter and more hooked; the throat and breaſt is alſo much more inclined to yellow than the Tit-Lark

is found to be in the ſpring. But it muſt be remarked that bird aſſumes much of that colour in the winter, which has occaſioned Mr. Lewin and others to conclude this bird is found with us in that ſeaſon; and the ſame author has evidently given the figure of the Tit-Lark for the Field Lark, by the length and ſtraitneſs of the hind claw.

The bill and hind claw of this bird are unerring marks of diſtinction by which it may at once be diſcriminated from the other. The legs of this are alſo uniformly of the ſame pale yellowiſh brown colour, never becoming duſky as in the matured birds of the other ſpecies.

We have been more particular in deſcribing the difference between theſe two ſpecies as we have frequently received one for the other. Dr. Latham ſhewed us ſeveral Larks that were ſent to him out of Yorkſhire, amongſt which one of this ſpecies was marked Tit-Lark, and two of thoſe birds called Field Lark and Pipit.

Mr. Pennant ſays the birdcatchers about London take a bird in the autumn they call Pipit, but does not deſcribe it.

We have great reaſon to believe the bird ſo called is the Pipit Lark of this work, not the Tit-Lark, and certainly not the Graſshopper Warbler, which we believe has been called Pipit; nor is it likely to be this bird, which leaves us at that ſeaſon in the ſame ſolitary manner they come to us, and are never known to be gregarious in this country.

The

The Tit-Larks aſſemble in ſmall flocks in autumn, and remain with us the whole year, which Mr. Lewin denies; and we mention this as another proof that he has completely confounded the two ſpecies. But this need ſcarcely be inſiſted on, for it is obvious when he aſſerts that the hind claw of the Field Lark is longer and ſtouter than in the Tit-Lark.

This ſpecies is by no means plentiful, but appears to be thinly ſcattered over moſt of the encloſed parts of England; is never met with on the moors or downs where the Tit-Lark is moſt frequent. Its ſong is vaſtly ſuperior to that bird, though ſomething ſimilar: this it delivers from the branch of a tree, or on wing, as it is deſcending to the ground. From the beginning of May to July it may be ſeen mounting in the air in a fluttering manner, at the ſame time uttering a twittering note, and then deſcends to ſome neighbouring tree with motionleſs wing and the tail thrown up. At this time it ſings, but never when riſing. And it is obſervable that it rarely pitches on the ground again until it has perched, and it always mounts in the like manner from a tree before it deſcends to the ground.

The Tit-Lark, it is true, has much of theſe manners, but after mounting in the air either returns to the ground or pitches on ſome low buſh. The Field Lark generally makes a neſt amongſt the high graſs or green wheat, and reſides wholly in the more cultivated parts, and that only where there

are trees. The neſt is compoſed of dry graſs, fibrous plants, and ſometimes a little moſs, and lined with fine dry graſs and horſe-hair. The eggs are four in number, of a dirty bluiſh white, thickly blotched, and ſpotted with purpliſh brown.

We have found this bird as far weſt as Devonſhire, but rarely in Cornwall; alſo in the weſtermoſt parts of South Wales, and in moſt of the ſouthern parts of England; but no where ſo plentiful as in the north of Wiltſhire.

LARK-GRASS-HOPPER. Vide *Warbler-graſs-hopper.*

LARK-LESSER-CRESTED.

Alauda nemoroſa. *Gmel. Syſt.* ii. p. 797.

Alauda criſtata minor. *Raii Syn.* p. 69. A. 5. *Will.* p. 152. *Brif.* iii. p. 361. 9.—*Ib.* 8vo. i. p. 411.

Alauda criſtatella. *Ind. Orn.* ii. p. 499. 26.

Le Lulu. *Buf.* v. p. 74.

Creſted Lark. *Br. Zool.* i. N° 141.—*Ib.* fol. 95.

Leſſer Creſted Lark. *Will. Angl.* p. 207. *Lath. Syn.* iv. p. 391. 24. *Lewin, Br. Birds,* iii. p. 9.

This ſpecies has long remained in obſcurity Aldrovandus is the only author who ſeems to hav given an original deſcription of it, from which a others appear to have copied; and as we have nev been fortunate enough to meet with it muſt do t ſame.

That author obſerves, Vol. ii. p. 371, that it is not ſo brown as the Greater Creſted Lark, and the tuft on the head larger in proportion to the ſize of the bird, and that the legs are red.

Dr. Latham remarks that it is ſaid it flies in flocks; and that it is for the moſt part met with in woods and thickets, where it makes a neſt. The ſame author obſerves that it is ſaid to be found in plenty in Yorkſhire.

It ſeems, however, very unlikely that if it was a diſtinct ſpecies it ſhould remain ſo long in obſcurity, at a time when the reſearches in natural hiſtory are ſo general; eſpecially as it has been ſaid to be plentiful.

From the deſcription there is greater reaſon to believe the bird mentioned by Aldrovandus is no other than the Field Lark of this work. Mr. Bolton informed Mr. Pennant it was found plentiful in Yorkſhire.

Amongſt a variety of Larks ſent from that county to Dr. Latham (we believe from Mr. Bolton) one was marked Leſſer Creſted Lark, which was evidently the Field Lark.

Is ſaid to be met with in many parts of Europe, Italy, Auſtria, Poland, and Siberia.

Buffon and Briſſon have both mentioned it, and neither of them ſpeak of the Field Lark; but we have little doubt their bird is that ſpecies.

LARK-LESSER-FIELD. Vide *Lark-field.*

LARK-MEADOW. Vide *Lark-field* and *Lark-tit.*

LARK-PIPIT.

Alauda trivialis. *Lin. Syſt.* i. p. 288. 5. *Gmel. Syſt.* ii. p. 796. *Ind. Orn.* ii. p. 493. 6.

Alauda ſepearia. *Briſ.* iii. p. 349. 5. — *Ib.* 8vo. i. p. 407.

Alouette pipi. *Buf.* v. p. 39. t. 4.

Pipit Lark. *Albin.* i. t. 44.

The weight of this ſpecies is about five drams and a half; length ſix inches and a half.

Bill duſky; the ſides and baſe of the upper mandible dull yellow; irides hazel.

The top of the head and whole upper parts of the body of a ruſty olivaceous-brown, ſtreaked with duſky, the middle of each feather being of that colour; the rump plain; the coverts and quill-feathers duſky; the former margined the ſame as the back, but lighter; the primary quills ſlightly edged with olive-green; thoſe next the body margined like the coverts; the whole under parts pale ferruginous, darkeſt on the breaſt, vent, and under tail coverts, and lighteſt on the chin; the ſides of the neck, breaſt, and ſides of the body, marked with oblong ſpots of duſky; the two middle tail-feathers duſky brown, paler on the margins; the outer feather on each ſide is white, except towards the baſe of the inner web; and the outer web is browniſh towards the point; the ſecond is white only at the tip; the reſt duſky black, ſlightly edged

edged with greeniſh brown; legs dull yellow; claws horn-colour; hind claw rather longer than the toe, a little bent.

This we believe is the ſpecies known in the neighbourhood of London by the name of Pipit, and is frequently taken by the birdcatchers when they firſt appear in flocks in the autumn.

It has no doubt been frequently confounded with the Tit-Lark, which alſo flock at that ſeaſon of the year, and perhaps ſometimes together. They are only to be diſcriminated by their ſuperior ſize, olivaceous colour on the ſuperior parts, and being much more ferruginous yellow beneath.

We have never been able to procure one ſpecimen in the ſummer, nor could we till lately identify the ſpecies.

The Pipit Lark is probably not leſs common than the Tit-Lark with us in winter; it frequents the ſame places. We have killed it in Cornwall, in very ſevere weather, in the month of December.

Mr. Pennant in his *Britiſh Zoology*, and Dr. Latham in his *Synopſis*, have made the Alauda trivialis of Linnæus the Graſshopper Warbler; but the latter author has, in his *Index Ornithologicus*, placed all the ſynonyms to this which he had given to that bird.

LARK-RED.

Alauda rubra. *Gmel. Syſt.* ii. p. 794. *Ind. Orn.* ii. p. 494. 10.

Alauda Penſylvanica. *Briſ. Sup.* p. 94.—*Ib.* 8vo. i. p. 419. 13.

Alouette à joues brunes de Penſylvanie. *Buf.* v. p. 58.

Lark from Penſylvania. *Edw.* t. 297.

Red Lark. *Br. Zool.* N° 140. *Arct. Zool.* ii. N° 279. *Lath. Syn.* iv. p. 376. 8. *Lewin, Br. Birds,* iii. t. 93.

This ſpecies is rather ſuperior in ſize to the Sky Lark. The bill is duſky above, whitiſh beneath, except at the point; irides hazel. The upper part of the head, hind part of the neck, and back, rufous-brown, each feather a little duſky in the middle; over the eye a pale ferruginous ſtreak; chin and throat the ſame; the ear coverts inclining to duſky; from the bill under the eye a narrow duſky line; the ſides of the neck and breaſt ferruginous, with duſky ſpots; belly and under tail coverts ferruginous white; greater quill-feathers duſky, ſlightly edged with yellowiſh white; the reſt of the quills deeply margined with rufous; ſome of the larger coverts the ſame, but thoſe immediately impending the ſecondary quills have whitiſh tips, making a ſmall bar acroſs the wing; one row alſo of the ſmaller coverts are tipped with white, making another line of that colour acroſs the ſuperior part of the wing; the two middle feathers of the tail are duſky, deeply margined with rufous-brown; the outer feather is white, the next is white on the exterior

exterior web, and part of the inner web towards the tip; the ſhaft duſky; the reſt are duſky; legs yellowiſh brown: hind claws as long as the toe, and ſomewhat curved.

The above deſcription is taken from a ſpecimen which was killed in Middleſex, and now in our poſſeſſion.

It is a rare ſpecies in England, but perhaps has ſometimes been confounded with the Sky Lark, which occaſionally partakes of the ferruginous colour. The hind claw, however, if every other mark of diſtinction was wanting, is ſufficient to ſeparate it from that bird; it is not ſo long, and is conſiderably curved, whereas that of the other is nearly ſtrait.

One in the Leverian Muſeum differs a little from the above.

Said to inhabit North America, and is ſometimes found in the neighbourhood of London.

LARK-ROCK.

Alauda obſcura. *Ind. Orn.* ii. p. 494. 7.

Alauda Petroſus. *Lin. Tranſ.* iv. p. 41. egg t. 2.

Duſky Lark. *Lewin, Br. Birds,* iii. t. 94. *Don. Br. Birds,* iv. t. 76.

Sea Lark. *Walc. Syn.* t. 193.

The length of this ſpecies is ſix inches three quarters; weight about ſeven drams.

The

The bill is duſky, near ſeven eighths of an inch long from the apex to the corner of the mouth; irides hazel.

Upper part of the head, back of the neck, and tail coverts, are of a dark brown; back and ſcapulars of the ſame colour, obſcurely marked with duſky ſtrokes; above the eye and beneath the ear is a lightiſh-coloured ſtroke; the throat whitiſh; breaſt and belly yellowiſh white, the former blotched with large duſky ſpots; the ſides marked with ſtrokes of the ſame; under tail coverts light brown; the two middle feathers of the tail dark brown, the others duſky; outer one of a dirty yellow, white on the interior web and the point of the exterior; in the ſecond feather the light colour is juſt viſible at the end; the quill-feathers and coverts are duſky, ſlightly edged with light brown; legs and toes duſky; claws black; hind claw four tenths of an inch long, and ſomewhat crooked.

Both ſexes are alike.

The young birds are not maturely feathered till after the winter of the firſt year; till then the upper parts have a tinge of olivaceous aſh-colour; beneath the lighter parts are yellowiſh, and the coverts of the wings more deeply margined with light brown; the baſe of the under mandible and legs leſs duſky.

It is probable Mr. Pennant firſt noticed this ſpecies in his folio edition of *Britiſh Zoology*, Pl. P. 1. and calls it a variety of Tit-Lark with duſky legs.

Long

Long after the laſt works of that author on Zoology, in which nothing more is ſaid of it, Dr. Latham favoured us with this bird amongſt a parcel of other Larks, ſtating he had received it from Mr. Lewin. At that time we were unacquainted with the bird, and for diſtinction returned it by the name of Duſky Lark, which Mr. Lewin adopted in his publication on *Britiſh Birds*; and Dr. Latham, in his *Index Ornithologicus*, calls it Alauda obſcura. Soon after theſe works had been given to the public we diſcovered theſe birds in great plenty on the coaſt of South Wales, where it was known by ſome of the natives by the name of Rock Lark; and afterwards found it not uncommon on all the coaſts from Kent to the Land's-End in Cornwall, where the ſhores were abrupt; and have no doubt it inhabits moſt of the rocky ſhores throughout the kingdom. It ſeems wholly confined to the neighbourhood of the ſea, and is never found, even in winter, more remote than in the contiguous marſhes within the occaſional influx of the tide, depending chiefly on marine inſects for its ſubſiſtence, and has never been obſerved to be gregarious.

The ſong, the manner of flying, and its habits in general, are ſo like thoſe of the Tit-Lark that it is probably owing to this circumſtance it had laid ſo long in obſcurity.

It begins breeding early in the ſpring. The neſt is made of dry graſs, marine plants, and very little moſs externally, and lined with fine graſs, with a few

few long hairs. This is generally placed on the ſhelf of a rock near the ſea; ſometimes at a conſiderable height, where there are a few ſcanty buſhes or tufts of graſs. It lays four or five eggs, of a dirty white, ſprinkled with numerous ſpecks of brown, darker and confluent at the larger end, ſo as to appear on that part wholly of that colour; in ſize they rather exceed thoſe of the Tit-Lark, weighing about thirty-ſix grains.

LARK-SEA. Vide *Lark-rock*, *Plover-ringed*, and *Bunting-tawny*.

LARK-SHORT-HEELED. Vide *Lark-field*.

LARK-SKY.

Alauda arvenſis. *Lin. Syſt.* i. p. 187. 1. *Gmel. Syſt.* ii. p. 791. *Ind. Orn.* ii. p. 491. 1.

Alauda vulgaris. *Raii Syn.* p. 69. A. 1. *Will.* p. 149. t. 40. *Briſ.* iii. p. 335. 1.—*Ib.* 8vo. p. 404.

L'Alouette. *Buf.* v. p. 1. t. 1.

Common Field Lark, *or* Sky Lark. *Will. Angl.* p. 203.

Sky Lark. *Br. Zool.* i. N° 136.—*Ib.* fol. 93. t. S. 2. f. 7. *Arct. Zool.* ii. p. 394. A. *Lath. Syn.* iv. p. 368. 1. *Albin.* i. t. 41.—*Ib. Song Birds*, p. 38. *Lewin, Br. Birds*, iii. t. 89. *Walc. Syn.* t. 189. *Pult. Cat. Dorſet.* p. 7.

The length of this ſpecies is ſeven inches.

Bill duſky; the baſe of the under mandible yellowiſh.

lowiſh. The feathers on the top of the head are duſky, bordered with rufous-brown; they are rather long and erectable in form of a ſhort creſt; the hind part is plain, inclining to aſh-colour; on the upper parts of the body the feathers are reddiſh brown, darker in their middle, the edges pale; the under parts are dirty buff-colour, darkeſt on the neck and breaſt, which parts are ſtreaked with duſky; quills brown, lighter on the outer webs and tips; the tail is duſky brown, the two middle feathers darkeſt, with light rufous margins; the outer feather is white on the outer web and tip of the inner; the ſecond feather white on the outer web only; the third is inclining to white on the margin of the outer web; legs duſky in old birds, but lighter in young; claws duſky; the hind one very long and ſtrait.

This bird is common in the greater parts of this kingdom, but moſt plentiful in the more open and higheſt cultivated ſituations abounding with corn, and rarely ſeen on the extended moors at a diſtance from arable land.

The neſt is placed on the ground amongſt graſs or corn; it is formed of dry graſs and other vegetable ſtalks, lined with fine dry graſs. The eggs are generally four in number, rather larger than thoſe of a Tit-Lark, weighing above fifty grains, of a dirty white, blotched and ſpotted with brown.

It begins to breed in May, and will lay as late as September if their firſt neſts are deſtroyed.

The

The hiſtory of this delightful ſinging bird is ſo generally known that to ſay more of it is uſeleſs. It has been aſſerted that the Sky Lark never perches; but this is a miſtake, for we have frequently ſeen it perch on the top of a buſh, and ſometimes on the branch of a tree.

LARK-TIT.

Alauda pratenſis. *Lin. Syſt.* p. 287. 2. *Gmel. Syſt.* ii. p. 792. *Ind. Orn.* ii. p. 493. 5. *Raii Syn.* p. 80. 4. *Will.* p. 150. *Brif.* iii. p. 343. 3. —*Ib.* 8vo. i. p. 406.

Spipola altera Aldr. *Raii Syn.* p. 80. 4. *Will.* p. 153.—171.

L'Alouette de prés. *Buf.* v. p. 31. t. 3.

Tit-Lark. *Br. Zool.* N° 138.—*Ib.* fol. 94. t. Q. f. 6. *Arct. Zool.* ii. p. 395. C. *Will. Angl.* p. 110? 206. *Albin.* i. t. 43.—*Ib. Song Birds,* t. p. 53. *Lath. Syn.* iv. p. 374. 5. *Lewin, Br. Birds,* iii. t. 98. *Walc. Syn.* ii. t. 191. *Pult. Cat. Dorſet.* p. 8.

This ſpecies weighs about four drams forty grains; length nearly five inches and three quarters. Bill ſlender, duſky, except at the baſe of the under mandible; irides hazel.

The upper part of the head and whole plumage above is of a duſky brown, with paler margins; from each ſide the under mandible a duſky line paſſes down the ſide of the throat; the throat and under

under parts dirty white; the ſides of the neck and breaſt marked with oblong ſpots of duſky; quills duſky brown, ſlightly edged with a paler colour; tail the ſame, the outer feather white, except at the baſe of the inner web; the next has a little white at the point; legs browniſh.

No birds have cauſed greater confuſion than ſeveral of the ſmaller ſpecies of Larks; the Field Lark and Pipit having always been more or leſs confounded with this bird.

The Tit-Lark remains with us the whole year, but after the autumnal moulting aſſumes more of an olivaceous tinge on the upper parts, and beneath is of a yellowiſh white. In this ſtate it has been taken for the Field Lark by thoſe who have not attentively examined the bill and hind claw. In theſe parts it more reſembles the Pipip; but that bird is much ſuperior in ſize, and the whole plumage is much more ferruginous.

The Tit-Lark is a very common bird in moſt parts of the kingdom; is partial to barren ſituations, and is found equally on the mountainous parts as well as in the low and ſwampy places. In Scotland it is almoſt the only bird found upon the vaſt extended tracts of heath, amongſt which it breeds. The neſt is placed on the ground amongſt furze or long graſs; is compoſed of bents, dry graſs, and ſtalks of plants, lined with fine dry graſs, and ſometimes long horſe-hair. The eggs vary conſiderably in colour; ſometimes of a dark brown,

others

others whitiſh, thickly ſpeckled all over with rufous-brown, or of a pale brown tinged with red; theſe are generally ſix in number. In the winter theſe birds moſtly frequent the lower grounds in ſearch of inſects and worms, and fly in ſmall flocks; poſſibly they may congregate with the Pipits, who come to us at that ſeaſon, and may then be readily confounded.

We have found the eggs of this bird from different neſts weigh from twenty-four to thirty-four grains.

LARK-WILLOW. Vide *Warbler-ſedge.*

LARK-WOOD.

Alauda arborea. *Lin. Syſt.* i. p. 287. 3. *Gmel. Syſt.* ii. p. 793. *Raii Syn.* p. 69. A. 2. *Will.* p. 149. t. 40. *Ind. Orn.* ii. p. 492. 3. *Briſ.* iii. p. 340. t. 20. f. 1.—*Ib.* 8vo. i. p. 405.

L'Alouette de bois, *ou* le Cujelier. *Buf.* v. p. 25.

Wood Lark. *Br. Zool.* N° 137.—*Ib.* fol. 94. t. Q. f. 3. *Arct. Zool.* ii. p. 395. B. *Will. Angl.* p. 204. *Albin.* i. t. 42.—*Ib. Song Birds,* t. p. 46. *Lath. Syn.* iv. p. 371. 3. *Lewin, Br. Birds,* iii. t. 90. *Pult. Cat. Dorſet.* p. 8. *Walc. Syn.* ii. t. 190.

This ſpecies weighs about eight drams; length ſix inches. Bill duſky; whitiſh at the baſe of the under mandible; irides hazel.

The general plumage much reſembles the Sky Lark;

Lark; the feathers on the top of the head and whole upper parts are duſky, margined with a light reddiſh brown; on the crown of the head the feathers are long, and capable of being erected in form of a ſhort creſt; from the bill over the eye is a narrow band of yellowiſh white ſurrounding the crown of the head; the coverts of the ears brown, beneath which is another light-coloured ſtroke; the neck and breaſt yellowiſh white, tinged with brown, marked with narrow duſky ſpots; belly dirty white; quill-feathers duſky, ſlightly edged with brown; the tail is ſhort; the two middle feathers are brown, the next duſky, and the four outer ones on each ſide are black, tipped with dirty white; the tail coverts are brown, and reach within half an inch of the end of the tail; legs yellowiſh fleſh-colour; hind claw long, and nearly ſtrait.

The Wood Lark is by no means a plentiful ſpecies, but is met with in moſt parts of the kingdom ſparingly. It ſings delightfully on wing, but rarely when ſitting on the ground, though ſometimes when perched on a tree. The ſong is much more melodious than that of the Sky Lark, but does not conſiſt of ſo great a variety of notes; but then it ſings almoſt throughout the year, except in the months of June and July. It does not mount in the air in the perpendicular manner, and continue hovering and ſinging in the ſame ſpot like the Sky Lark, but will ſometimes ſoar to a great height, and keep flying in large irregular circles, ſinging

the whole time with little intermiſſion; and will thus continue in the air for an hour together.

It is a very early breeder, beginning to build in March. We have found the neſt with eggs as early as the fourth of April. It is placed on the ground, moſt commonly in rough and barren land, under a tuft of high graſs, furze, or ſome low buſh; is made of dry graſs, lined with finer, with ſometimes a few long hairs. The eggs are generally four in number, brown, mottled with duſky and cinereous, moſtly at the larger end; are ſomewhat leſs than thoſe of the Sky Lark; their weight from forty to fifty grains.

Theſe birds rarely aſſemble in larger flocks than ſix or ſeven; moſt probably the family, which aſſociate together till the returning ſpring. Their food is grain and ſeeds of various kinds, as well as inſects.

LAVY. Vide *Guillemot-fooliſh.*

LINNET-BROWN.

Fringilla Linota. *Gmel. Syſt.* ii. p. 916. *Ind. Orn.* i. p. 457. 81.

Linaria. *Raii Syn.* p. 90. A. 1. *Will.* p. 190. —*Ib. Angl.* p. 258. t. 46. *Briſ.* iii. p. 131. 29. —*Ib.* 8vo. i. p. 343.

La Linotte. *Buf.* iv. p. 58. t. 1.

Common Linnet. *Br. Zool.* N°130.—*Ib.* fol. 110. *Albin. Song Birds*, t. p. 31. *Lewin, Br. Birds*, ii. t. 83. *Lath. Syn.* iii. p. 402. 73. *Pult. Cat. Dorſet.* p. 12. *Walc. Syn.* ii. t. 221.

This

This bird is generally deſcribed to be of the ſame ſize as the Greater Redpole, or Red-headed Linnet. The plumage, on compariſon of the various authors who have given it as a diſtinct ſpecies, makes it nearly the ſame as that bird, but not quite ſo rufous on the upper parts. The principal diſtinction ſeems to be in the breaſt being of a fine crimſon colour, and none of that colour on the head.

Linnæus does not appear to have conſidered theſe birds as diſtinct; and we have no doubt he was perfectly right, for they are to be met with in all gradations with reſpect to the red markings on the head and breaſt; ſometimes on one of thoſe parts only, at other times on neither: this depends wholly on age and ſeaſon. From the vaſt number we have killed at all ſeaſons of the year, in which the greateſt variety of thoſe markings were obſerved, we do not heſitate to pronounce them one and the ſame ſpecies. For further remarks, ſee *Redpole-greater.*

LINNET-GREEN. Vide *Groſbeak-green.*

LINNET-MOUNTAIN. Vide *Twite.*

LINNET-RED. Vide *Redpole-greater.*

LINNET-RED-HEADED. Vide *Redpole-greater.*

LINNET-RED-HEADED-LESSER. Vide *Redpole-leſſer.*

LONG-NECK. Vide *Bittern-little.*

LONG-

LONG-TAILED-CAPON. Vide *Titmouſe-long-tailed.*

LONG-TAILED-MAG. Vide *Titmouſe-long-tailed.*

LOON. A common name given to ſeveral ſpecies of the Divers and Grebes.

LOON-ASH-COLOURED. Vide *Grebe-tippet.*

LOON-GREATER. Vide *Grebe-tippet.*

LOON-RED-THROATED. Vide *Diver-red-throated.*

LOON-SPRAT. Vide *Diver-ſpeckled.*

LUMME-SPRAT. Vide *Diver-black-throated.*

LUNDA. Vide *Puffin.*

END OF FIRST VOLUME.

T. Bensley, Printer, Bolt Court, Fleet Street, London.

Ornithological Dictionary.

Ornithological Dictionary;

OR,

Alphabetical Synopsis

OF

BRITISH BIRDS.

BY

GEORGE MONTAGU, F.L.S.

IN TWO VOLUMES.

VOL. II.

LONDON:

PRINTED FOR J. WHITE, FLEET STREET,

BY T. BENSLEY, BOLT COURT.

1802.

ERRATA, VOL. II.

Magpie, line 23, huſbandmen, *read* huſbandman
Martin, line 72, general warmth, *read* genial
Night-jar, vide Shriek-cinereous, *read* Shrike
Owl-great-eared, line 4, black-ferruginous, *read* black, ferruginous
Owl-Tawny, line 52, its egreſs, *read* its ingreſs and egreſs
Owl-White, Strix flaminea, *read* flammea
Pochard, line 27, tracheæ, *read* trachea
Puffin, line 58, Coulternel, *read* Coulterneb
Quail, line 47, looſe, *read* loſe
Rail, line 26, ſedge-ruſhes, *read* ſedge, ruſhes
Redpole-greater, gooſebery, *read* gooſeberry
Roller, generic character, characters or which, *read* of which
Sanderling, line 7, the ſhafts, *read* the ſhaft
Sandpiper-Greenwich, Grenovieenſis, *read* Grenovicenſis
Sandpiper-Spotted, line 3, upper parts, *read* upper part
Shrike-red-backed, line 37, clytra, *read* elytra
Stone-chat, line 29, Gold-Finch, *read* Cold-Finch
Warbler, generic character, Toes, the extreme, *read* exterior
Warbler-Dartford, line 15, alula ſpuriaræ, *read* ſpuria
Woodcock, line 22, northern counties, *read* countries
Woodpecker-green, line 3, tipped with cinereous, *read* crimſon
Ditto, line 27, colcopterous, *read* coleopterous.

Pult. Cat. Dorſet. p. 5. *Walc. Syn.* i. t. 39.
Don. Br. Birds, iv. t. 95.

PROVINCIAL.

Hagiſter.

This ſpecies of Crow is about eighteen inches in length; weight between eight and nine ounces. Bill black; irides hazel. The breaſt, upper part of the belly, and ſcapulars, white; the reſt of the plumage black; the wings and tail beautifully gloſſed with blue, green, and purple; the ten firſt quill-feathers are white in the middle on the inner web; the tail is nine inches and a half long, and very cuneiform, the outer feathers not being much above half as long as the middle ones; legs and claws black.

The female differs only from the other ſex in being rather leſs, and in the tail being ſhorter.

It has been very juſtly remarked, that England does not produce a more beautiful bird than the Magpie; but that thoſe who have only ſeen the dirty mutilated ſpecimens in captivity can form no idea of its native beauty.

Theſe birds generally continue in pairs the whole year; and though ſhy it rarely removes far from the habitation of man. Its attachment, however, is governed by ſelf-intereſt; is a great enemy to the huſbandmen, but has cunning enough to evade his wrath. No animal food comes amiſs to its carnivorous appetite; young poultry, eggs, young lambs, and even weakly Sheep, it will attempt to deſtroy by firſt plucking out their eyes; the young of Hares, Rabits, and feathered game, ſhare the ſame fate; fiſh, carrion, inſects, and fruit; and laſtly

laſtly grain, when nothing elſe can be got. It is an artful, clamorous bird, proclaiming aloud any apparent danger, and thereby gives notice to its aſſociates. Neither the Fox or other wild animal can appear without being noticed and haunted; even the fowler is frequently ſpoiled of his ſport, for all other birds ſeem to know the alarming chatter of this bird.

Its neſt is curiouſly built for the defence of its young; it is of an oval ſhape, made of ſticks, generally the black thorn, ſtrongly woven together, with only a ſufficient entrance on one ſide; the bottom is plaſtered with earth, and lined with fibrous roots.

It lays ſix or ſeven eggs early in the ſpring, of a yellowiſh white, ſpotted with brown and cinereous.

The place of nidification is various; ſometimes on the top of a high tree, in a thick buſh or hedge, and frequently at a ſmall diſtance from the ground. From theſe circumſtances it has been ſuppoſed there are two ſpecies, and have ſometimes been denominated the Tree-Mag and the Hedge-Mag.

In winter theſe birds aſſemble in great numbers to rooſt in ſome coppice or thicket, but ſeparate again in the day.

When reclaimed the Magpie is a very docile bird, and ſeems to have its natural faculties heightened by domeſtication, imitating the human voice, and various other ſounds.

MAGPIE-MOUNTAIN. Vide *Shrike-cinereous.*

MALLARD. Vide *Duck-common.*

MASH-HEN. Vide *Gallinule-common.*

MARROT. Vide *Puffin, Razor-bill, Guillemot-leſſer.*

MARTIN.

Hirundo urbica. *Lin. Syſt.* i. p. 347. 3. *Gmel. Syſt.* ii. p. 1017. *Ind. Orn.* ii. p. 573. 3.

Hirundo ruſtica, ſive agreſtis. *Raii Syn.* p. 71. A. 2. *Will.* p. 155. t. 39. *Brif.* ii. p. 490. 2. —*Ib.* 8vo. i. p. 295.

Hirondelle à cul blanc. *Buf.* vi. p. 614. t. 25. f. 2.

Martin, Martlet, *or* Martinet. *Br. Zool.* i. N° 169. —*Ib.* fol. 96. t. Q. f. 2. *Arct. Zool.* ii. N° 331. *Will. Angl.* p. 213. t. 39. *Albin.* ii. t. 56. a. *Lath. Syn.* iv. p. 564. 3. —*Ib. Sup.* p. 192. *Lewin, Br. Birds,* iii. t. 124. *Pult. Cat. Dorſet.* p. 13. *Walc. Syn.* ii. t. 252.

PROVINCIAL.

House Martin.

This ſpecies of Swallow is rather inferior in ſize to the Chimney Swallow; the length is about five inches and a half. Bill black; irides hazel. The whole upper parts are of a gloſſy blue-black, the rump excepted, which, with the under parts from chin

chin to vent, is white; the tail is blue-black, and forked; the legs are covered with a white down; the claws white.

This well-known ſpecies viſits England in the ſpring rather later than the Common Swallow. It firſt makes its appearance in low, warm ſituations, and if the weather is fine begins building early in May.

The neſt is generally placed under the eve of a houſe; ſometimes againſt rocks or cliffs contiguous to the ſea. It is built with mud externally, and lined with feathers, with a ſmall hole on one ſide for entrance.

The eggs are four or five in number, of a pure white. The young never quit the neſt till they are able to fly well, and are fed by the old birds frequently on wing. At the time of incubation, and when the young are firſt hatched, it is not uncommon to ſee both the old birds in the neſt together, in which place the act of conſummation is performed.

Theſe birds, like the reſt of the genus, are conſtantly on wing, are rarely ſeen to perch on a tree, and ſeldom ſettle at all during the day, except in the autumn, juſt before their winter migration, at which time they aſſemble in large flocks on the roofs of houſes and other buildings, expoſing themſelves to the influence of the ſun; but we have never been able to aſcertain where they rooſt at

night after they have forſaken their neſts; but moſt probably in holes under the tiles and thatch of houſes.

Much has been ſaid by various authors concerning the winter ſtate and abode of this bird and its congeneries. To attempt to refute the abſurd idea of their immerſion would be a treſpaſs on the patience of our readers. It muſt, however, be obſerved, that their internal formation abſolutely prevents it; nor is there any occaſion to reſort to ſuch unnatural means when no birds are furniſhed with more ample powers for migration. That a few inſtances of their having been found in old ſhafts of mines, and in ſimilar ſituations, we will not diſpute; though it is ſomewhat extraordinary that thoſe who have ſtated ſuch as facts do not mention which ſpecies of Swallow was ſo found; nor have we been able to find a ſingle perſon of good authority who ever ſaw the fact.

Why theſe birds ſhould ever have been denied their migrative powers we cannot conceive, when others, much leſs qualified for long flights, have not been doubted.

That this bird, as well as the Chimney Swallow, are now and then ſeen flying about long after the general diſappearance of theſe birds, we have more than once had ocular proof of. But theſe inſtances muſt be attributed to ſome accident or individual defect, which prevented them from performing their

their uſual autumnal flight. It is well known that in ſome animals, whoſe radical heat is ſenſibly affected by cold, the power of action is loſt when the fluids become languid, and animal life is as it were ſuſpended. The Batt, the Dormouſe, and Hedgehog, become torpid in winter, when the mean ſtate of the air is below 45 degrees; and their heat ſeems to keep pace nearly with the ſtate of the atmoſphere. But even at this ſeaſon the general warmth of the ſun regenerates their benumbed limbs, and the Batt is ſometimes ſeen flying about at mid-day, but returns again to its former ſtate for weeks, and perhaps months, if the air proves colder.

Thus it is with the Martin and Swallow who have been accidentally detained in this northern climate; they are rouſed by a certain degree of heat, and the calls of hunger induce them to fly abroad for food; and it is much to be doubted whether this ſudden return of all the animal functions does not prove fatal, from not being able to find ſufficient food to ſupply the natural excretions, which in a torpid ſtate have been obſerved to be little or none.

It cannot be ſuppoſed it was intended by nature that birds who have the power of ſhifting their quarters in ſo expeditious a manner, and with ſuch eaſe, ſhould ever repoſe during the colder ſeaſon. It is therefore moſt reaſonable to ſuppoſe ſuch accidental migrative birds, who are detained in a climate

unſuitable to their nature, periſh before the return of the warmer months.

We have known ſeveral inſtances of a ſingle bird of this ſpecies flying about in ſearch of food after the middle of November, but never for two days together, nor after the latter end of that month.

Found in moſt parts of Europe, and in Aſia.

MARTIN-BANK. Vide *Martin-ſand.*

MARTIN-BLACK. Vide *Swift.*

MARTIN-HOUSE. Vide *Martin.*

MARTIN-SAND.

Hirundo riparia. *Lin. Syſt.* i. p. 334. 4. *Gmel. Syſt.* ii. p. 1019. *Raii Syn.* p. 71. A. 3. *Will.* p. 156. t. 39. *Ind. Orn.* ii. p. 575. 10. *Brif.* ii. p. 506. 12.—*Ib.* 8vo. i. p. 299.

L'Hirondelle de rivage. *Buf.* vi. p. 632.

Sand Martin, *or* Shore-bird. *Br. Zool.* i. N°170. —*Ib.* fol. 97. t. Q. f. 1. *Arct. Zool.* ii. N°332. *Will. Angl.* p. 213. t. 39. *Albin.* ii. t. 56. b. *Lath. Syn.* iv. p. 568. 10. *Lewin, Br. Birds,* t. 125. *Walc. Syn.* ii. t. 253. *Pult. Cat. Dorſet.* p. 13.

PROVINCIAL.

Bank Martin.

This is the ſmalleſt ſpecies of Britiſh Swallow; length four inches and three quarters.

Bill duſky; irides hazel.

The

The whole upper parts of the plumage are of a mouſe-coloured brown; the under parts white, except acroſs the breaſt, which is brown; legs duſky, a little feathered behind.

The Sand Martin is not near ſo plentiful, and is more local than the other ſpecies.

It viſits England about the ſame time as the Swallow, reſorting only to ſuch places as are convenient for breeding; is frequently ſeen about rivers, where it makes a neſt in the banks, but moſt commonly in ſand-pits, where it can with more eaſe excavate the ſand in order to form a ſecure place for its neſt. The holes are generally horizontal, and their depth two or three feet. Sometimes they build in old walls, and, we are told, in hollows of trees. The neſt is compoſed of ſtraw and dried fibres, lined with feathers.

The eggs are four or five in number, quite white, like thoſe of the Martin, but rather ſmaller; weighing about twenty-two grains.

The manners of this ſpecies is much like that of the common Martin, and are often ſeen in company together ſkimming over water in purſuit of gnats and other ſubaquatic inſects. This, as well as the other ſpecies, have been ſuppoſed to lay torpid in their holes all the winter; and many fruitleſs attempts have been made to diſcover them at that ſeaſon by digging to the bottom of thoſe holes where they reſort in ſummer.

It is found in moſt parts of Europe, and is ſaid

to

to be common in America, where it is called Ground Swallow.

MARTINET. Vide *Martin.*

MARTLET. Vide *Martin.*

MATTAGES. Vide *Shrike-cinereous.*

MERGANSER A genus of birds, the characters of which are,

Bill ſlender, a little depreſſed, furniſhed at the end with a crooked nail; edges of each mandible very ſharply ſerrated.

Noſtrils, near the middle of the mandible, ſmall and ſuboval.

Toes three before, webbed, the outer one the longeſt; hind toe furniſhed with a fin.

MERGANSER-MINUTE.

Mergus minutus. *Lin. Syſt.* i. p. 209. 6? *Faun. Suec.* N° 138. *Ind. Orn.* ii. p. 575. 10. *Gmel. Syſt.* ii. p. 548. 6.

Mergus glacialis. *Raii Syn.* p. 135. A. 3. *Will.* p. 254.

Merganſer criſtatus minor. *Briſ.* vi. p. 243. 3. t. 24. f. 2.—*Ib.* 8vo. ii. p. 426.

Weeſel Coot. *Albin.* i. t. 88.

Red-headed Smew, *or* Lough-Diver. *Br. Zool.* ii. N° 263?—*Ib.* fol. 141. t. Addend.? *Arct. Zool.* ii. p. 540. A.—*Ib. Sup.* p. 741. *Will. Angl.* p. 338. *Walc. Syn.* i. t. 83. *Pult. Cat. Dorſet.* p. 19.

Minute

Minute Merganſer. *Lath. Syn.* vi. p. 429. 6. *Lewin, Br. Birds*, vi. t. 235. *Don. Br. Birds*, t. 52.

This bird is about the ſize of a Teal; the length rather exceeds fourteen inches. Bill black; irides hazel. The upper part and ſides of the head, taking in the eyes, duſky ferruginous, almoſt black about the eyes, eſpecially between that and the bill; on the back part of the head the feathers are long and ſlender; the chin, throat, and upper part of the neck before, are white; the lower part of the neck and breaſt duſky and cinereous-grey mixed; the under parts white; the back, rump, and ſcapulars, black; the laſt, as well as the upper part of the back, daſhed with cinereous; primary quills duſky; ſecondaries gloſſy black, ſlightly tipped with white; their coverts tipped the ſame, forming two ſlender bars acroſs the wing; on the upper coverts a large patch of white; tail ſomewhat cuneiform, duſky; legs duſky.

This bird is ſubject to vary a little in colour about the head; ſome are more ferruginous than others; and ſometimes that part is duſky with only a reddiſh tinge. No birds have perplexed the naturaliſt more than ſome ſpecies of Merganſers; and various are the opinions concerning the one before us. If we conſider that the young may not arrive at maturity in plumage for two or three years it may admit of great doubt whether this may not be the young or female of the Smew.

Dr.

Dr. Latham, in his Eſſay on the *Trachea* of Birds, publiſhed in the fourth volume of the *Linnæan Tranſactions*, conſiders that there are but three diſtinct ſpecies of Merganſer which migrate into this country; and ſays he has found the *labyrinth* of the ſuppoſed male of this bird in every reſpect ſimilar to that of the male Smew; from which he concludes it is that bird not arrived to maturity in plumage.

We have long had our ſuſpicions of what the Doctor has aſſerted; and we ſtrongly recommend attention to this new field of inveſtigation, particularly in this and the Duck tribe, from which conſiderable knowledge is to be expected.

It is not frequently met with in the ſouth of England, and only in winter when the weather is ſevere; and rarely otherwiſe than ſingle, or at moſt two together.

Its principal food is fiſh, after which it dives with great eaſe, and remains long under water.

MERGANSER-RED-BREASTED.

Mergus ſerrator. *Lin. Syſt.* i. p. 208. 3. *Gmel. Syſt.* ii. p. 546. 3. *Ind. Orn.* ii. p. 829. 4.

Mergus cirrhatus fuſcus. *Raii Syn.* p. 135. A. 4. *Will.* p. 255. t. 64.

Mergus criſtatus. *Brif.* vi. p. 237. 2. t. 23.— *Ib.* 8vo. ii. p. 424.

Le Harle huppé. *Buf.* viii. p. 273.

Red-breaſted Merganſer. *Br. Zool.* ii. N° 261. t. 93.

t. 93.—*Ib.* fol. 147. *Arct. Zool.* ii. N° 466. *Edw.* t. 95. *Albin.* ii. t. 101. *Lath. Syn.* vi. p. 423. 3. *Lewin, Br. Birds,* vi. t. 233. *Walc. Syn.* i. t. 81. *Pult. Cat. Dorſet.* p. 18. *Don. Br. Birds,* ii. t. 38.

PROVINCIAL.

Red-breaſted Gooſander. Leſſer-toothed Diver. Serula.

This ſpecies is about twenty-one inches in length; weight two pounds.

The bill is three inches long; the upper mandible duſky, the lower red; irides purpliſh red. The head and part of the neck black, gloſſed with green; on the back of the head the feathers are long, forming a ſort of pendant creſt; the reſt of the neck and under part of the body white; breaſt ferruginous, mixed with black and white; upper part of the back gloſſy black; rump marked with brown and cinereous tranſverſe ſtreaks; the ſcapulars and wing coverts are ſome black and ſome white; quills duſky; tail brown; legs orange; claws black.

The female has the upper part of the neck and head dull ferruginous; fore part of the neck and breaſt the ſame, mottled with black and white; the greater part of the neck behind, back, rump, and ſcapulars, cinereous; lower part of the breaſt and belly white. Both ſexes are ſubject to ſome variety in plumage.

Theſe

Theſe birds ſometimes appear in the ſouth of England in winter, but more frequently in the north; are ſaid to breed in Scotland in ſome of the lochs. Is found in the Ruſſian dominions, about the great rivers of Siberia.

Is ſaid to breed on the ſhores of Greenland, and are obſerved at Hudſon's Bay in large flocks; breeds there as well as at Newfoundland, chiefly on the iſlands. The neſt is ſaid to be made with dry graſs, lined with down; the eggs are white; ſometimes as many as thirteen in a neſt, about the ſize of thoſe of a Duck.

MERLIN.

Falco Œſalon. *Gmel. Syſt.* i. p. 284. *Ind. Orn.* i. p. 49. 119. *Raii Syn.* p. 15. 15. *Will.* p. 50. t. 3. *Briſ.* i. p. 382. 23.—*Ib.* 8vo. p. 111.

Merlin. *Br. Zool.* i. N° 63.—*Ib.* fol. t. A. 12. *Will. Angl.* p. 85. *Lewin, Br. Birds*, i. t. 22. *Lath. Syn.* i. p. 106. 93.—*Ib. Sup.* p. 29. *Pult. Cat. Dorſet.* p. 3. *Walc. Syn.* i. t. 22. *Don. Br. Birds*, iv. t. 94.

The weight of the male of this ſpecies of Falcon is about five ounces; length ten inches. Bill bluiſh lead-colour; cere greeniſh yellow; irides duſky.

The crown of the head is duſky brown, ſtreaked with black down the ſhafts of each feather; on the back of the head the feathers are white at the baſe, tipped

tipped with ferruginous; the middle of each feather, black at the point; the back, ſcapulars, rump, and wings, cinereous lead-colour, each feather marked with a long ſlender line of black down the ſhaft; greater quills black; the inner webs marked with many oblong white ſpots; thoſe next the body are coloured like the back on the outer webs; the inner webs ſpotted as the others; the two firſt feathers are much indented towards the point of the inner web, as if cut with a pair of ſciſſars; the third feather rather exceeds the ſecond in length, and is the longeſt; the throat is nearly white; breaſt, belly, ſides, and thighs, ferruginous, ſtreaked with duſky; vent and under tail coverts pale ferruginous; the under wing coverts are rufous-brown, with round white ſpots on each web; tail like the back, croſſed with ſix or ſeven bars of black; the end black for almoſt an inch, ſlightly tipped with white; legs yellow; claws black.

The female weighs about nine ounces; length twelve inches and a half. Bill, cere, and irides, the ſame. The whole upper parts of the plumage are brown, tinged with ferruginous, with duſky ſtreaks on the ſhafts; beneath yellowiſh white, with broad duſky brown ſtreaks; tail brown, like the back, with five or ſix narrow bars of yellowiſh white, tipped with the ſame.

The above deſcription is taken from the birds now before us; but theſe birds ſeem to vary a little in the markings. Mr. Pennant obſerves, the bars

on

on the tail are generally from thirteen to fifteen in number; but remarks in one ſpecimen there were only eight.

The male above deſcribed was recently taken alive in a trap-cage hung in a paſſage of a houſe, in which there was a Bulfinch; and what was extraordinary it had loſt an eye. We kept it alive for ſome time, and found it extremely docile; and what was remarkable, would drink freely whenever water was offered it, and ſhewed ſigns of diſtreſs when long kept without it. This remarkable thirſt, ſo unuſual in predaceous birds, which we have kept for years without ever attempting to drink, was certainly occaſioned by fever, for it died of an inflammation on its lungs.

The Merlin is exceedingly rapid on wing, and was uſed formerly in falconry, being eſteemed for its courage, though inferior in ſize. The female will kill a Partridge at a ſingle pounce; but the male is contented with humbler game. The wing of this ſpecies is not ſo long and pointed as that of the Hobby; when cloſed it does not reach to the end of the tail by an inch and a half.

It flies low, and is generally ſeen ſkimming along the ſide of a hedge, or over the ſurface of the ground, in purſuit of ſmall birds. Theſe birds viſit the ſouth of England in October, about the time the Hobby retires, but has never been obſerved to breed further ſouth than Cumberland, where, Dr. Latham informs us, it has been found more

than once with four young ones, placed on the ground. In the middle of a high clump of heath upon the moors in Northumberland we found three young ones about half grown, but no neſt. They were well concealed, and would not have been diſcovered but by a ſetting-dog making a point at them.

The eggs are ſaid to be of a plain chocolate-colour, and that an inſtance has been known of its depoſiting them in a deſerted Crow's neſt.

MEW. Vide *Winter-mew.*

MIRE-DRUM. Vide *Bittern.*

MISSEL-BIRD. Vide *Thruſh-miſſel.*

MOCK-NIGHTINGALE. Vide *Blackcap.*

MOOR-BUZZARD.

Falco œruginoſus. *Lin. Syſt.* p. 130. 29. *Gmel. Syſt.* i. p. 267. *Raii Syn.* p. 17. A. 4. *Will.* p. 42. t. 7. *Ind. Orn.* i. p. 25. 53.

Circus paluſtris. *Briſ.* i. p. 401.—*Ib.* 8vo. p. 115.

Buſard. *Buf.* i. p. 218. t. 10.

Moor Buzzard. *Br. Zool.* i. N° 57.—*Ib.* fol. 67. t. A. 5. *Arct. Zool.* ii. p. 225. L. *Will. Angl.* p. 75. t. 7. *Albin.* i. t. 3. *Lewin, Br. Birds,* i. t. 8. *Lath. Syn.* i. p. 53.—*Ib. Sup.* p. 15. *Walc. Syn.* i. t. 8. *Pult. Cat. Dorſet.* p. 3.

The ſpecimen now before us of this ſpecies of Falcon is a female; its weight twenty-eight ounces

and a half; length twenty-three inches and a half. Bill black; cere and irides yellow. The whole plumage is of a chocolate brown, tinged more or leſs with ferruginous; the crown of the head is of a dull yellow; legs long and yellow; claws black.

The male is ſomewhat leſs, weighing about twenty-one ounces; length twenty-one inches; plumage the ſame, but generally brighter in colour.

Theſe birds are ſubject to ſome variety in markings: ſometimes the crown of the head is white; others have the whole head white, or yellowiſh; the ſhoulders are ſaid to be ſometimes yellow; and we have ſeen a ſpecimen with the head, part of the wing coverts, and the four firſt quill-feathers, white. Theſe markings are conſidered as mere varieties, as ſome are ſaid to be found wholly of a chocolate-brown: ſuch, however, are by no means ſo common as thoſe with a yellow crown, or of various ſhades of that colour to white. In more than twenty ſpecimens we have examined this was the conſtant mark, except as above deſcribed.

This ſpecies appears to be local, moſtly frequenting ſwampy moors and barren ſituations; and though rarely met with in the more cultivated parts, it is the moſt common of the Falcon tribe about the ſandy flats on the coaſt of Caermarthenſhire, where they prey upon young Rabbits; and we have ſeen no leſs than nine feeding at one time upon the carcaſe of a ſheep.

It

It will ſometimes feed on frogs, lizards, worms, and even the larger inſects.

The neſt is moſt frequently made on the ground amongſt ſhort wood, furze, or fern. It is compoſed of ſticks, ruſhes, or coarſe graſs; ſometimes, though rarely, it builds in the fork of a large tree.

In both theſe ſituations we have found the neſt with eggs. Theſe are perfectly white, without any ſpots; conſiderably leſs than thoſe of the Common Buzzard.

It is by no means a bird of rapid flight, and therefore pounces its prey on the ground; for which purpoſe it is generally ſeen ſkimming over the ſurface of the ground like the Ringtail. It is ſaid to prey on fiſh occaſionally, and on young Ducks, and other water-fowl; from which circumſtance it has in ſome parts obtained the name of Duck Hawk.

In the breeding ſeaſon, when the female is ſetting, the male will ſoar to a conſiderable height, and remain ſuſpended on wing for a great length of time.

MOOR-COCK. Vide *Grous-red.*

MOOR-HEN. Vide *Gallinule-common.*

MOOR-TITLING. Vide *Stone-chat.*

MORILLON.

Anas Glaucion. *Lin. Syſt.* i. p. 201. 26. *Gmel. Syſt.* ii. p. 525. *Raii Syn.* p. 143. 11.—*Ib.* p. 144. *Will.* p. 282. *Ind. Orn.* ii. p. 868. 88.

Morillon. *Brif.* vi. p. 406. t. 36. f. 1. 2.—*Ib.* 8vo. ii. p. 468.

Glaucion, *or* Morillon. *Br. Zool.* ii. N° 277. *Arct. Zool.* ii. p. 573. F. *Will. Angl.* p. 367. *Lath. Syn.* vi. p. 537. 77. *Walc. Syn.* i. t. 70.

Although a bird of the Duck kind has been given under this denomination by a variety of authors of repute, we confeſs to be unacquainted with ſuch as a diſtinct ſpecies. The Golden-eye, Tufted, and Scaup Ducks, have been given the name of Morillon; that which Linnæus calls the Glaucion is ſuppoſed to be the Morillon of Briſſon. Dr. Latham, whoſe excellence and attention in the ſtudy of Ornithology is ſo well known, acknowledges, in his *Synopſis*, his doubts as to the identity of this bird, though he has continued it as a diſtinct ſpecies in his *Index Ornithologicus*. That gentleman mentions having received a pair by the name of Morillon, which were ſo like the female Golden-eye that he was ſtruck with the circumſtance; and we have every reaſon to believe the Morillon of Briſſon to be no other than a variety of the male Golden-eye, or Tufted Duck. The former of theſe birds is ſubject to great variety, probably from age; and we have ſeen it accord ſo nearly with the deſcription given of this bird, that there ſeems to be little doubt they are one and the ſame ſpecies.

The deſcription given of this bird is that it is rather

rather leſs than the Golden-eye; length near fifteen inches. Bill lead-colour; irides golden yellow; the head is ſomewhat creſted; that and moſt of the neck black, gloſſed with violet; the lower part of the neck rufous-brown; the back, ſcapulars, and rump, gloſſy blackiſh brown, with a ſlight tinge of violet; the feathers on the breaſt brown, deeply edged with white; belly white, mixed with brown near the vent; wing coverts blackiſh brown; moſt of the inner ones have a caſt of green; the ten firſt quills are blackiſh brown; of theſe from the fourth to the tenth are marked with grey more or leſs in the middle of the outer web near the ſhaft; the eleventh and twelfth pale grey, with brown tips; the eight following white, tinged as the laſt; the inner of theſe brown on the inner web; the ſix next the body dark brown, and the outer one of theſe has a white ſpot on the outer web; tail violet brown; legs lead-colour; claws black.

The deſcription given by Mr. Pennant differs conſiderably from that of Briſſon.

The bill is of a yellowiſh brown; irides gold-colour; the head of a duſky ruſt-colour; round the upper part of the neck is a collar of white, beneath that a broader of grey; the back and coverts duſky, with a few white lines; the greater coverts duſky, with a few great ſpots of white; the primaries black; the ſecondaries white; breaſt and belly white; tail duſky; the ſides above the thighs black; the legs yellow. The ſame author adds,

that the bird deſcribed was bought in the London market, but was doubtful of the ſex. And further remarks, that about the ſame time he was favoured with an account of two birds of this ſpecies ſhot in Eſſex; both agreed in colour, but one weighed twenty-ſix ounces, the other only nineteen.

In the ſecond edition of the *Britiſh Zoology* this bird is called Grey-headed Duck.

This ſeems to be the bird figured by Lewin, Vol. VII. t. 256, and is called Brown-headed Duck.

We believe there is little doubt this is no other than a variety of the Golden-eye in one ſtage of plumage before it arrives to maturity; many of which we have ſeen and proved to be ſuch by the ſingular ſtructure of the windpipe.

In Briſſon's bird the head being black and the legs lead-coloured, induces us to think it is a variety of the Tufted Duck, eſpecially as he mentions the head to be ſomewhat creſted; for many of them are without the long pendant feathers. We muſt, however, leave the ſubject to future ornithologiſts, who may hereafter be able to determine with more certainty if ſuch a bird as the Morillon actually exiſts as a diſtinct ſpecies. And we ſtrongly recommend to their attention a careful examination of the *trachia*, or windpipe, which ſubject is ſo ably handled by Dr. Latham in the fourth volume of the *Linnæan Tranſactions*, accompanied with excellent figures of the windpipes of a variety of birds;

birds; amongſt theſe are thoſe of the Golden-eye and Tufted Duck, the ſtructure of which may be conſulted.

MORROT. Vide *Guillemot-leſſer.*
MOUNTAIN-COCK. Vide *Grous-wood.*
MUMRUFFIN. Vide *Titmouſe-long-tailed.*
MURDERING-BIRD. Vide *Shrike-cinereous.*
MURRE. Vide *Guillemot-fooliſh* and *Razor-bill.*

N.

NETTLE-CREEPER. Vide *Whitethroat.*
NETTLEMONGER. Vide *Whitethroat.*
NIGHT-HAWK. Vide *Goatſucker.*
NIGHTINGALE.

Motacilla Luſcinia. *Lin. Syſt.* i. p. 328. 1. *Gmel. Syſt.* ii. p. 950. *Raii Syn.* p. 78. A. 2. *Will.* p. 161. t. 41. *Briſ.* iii. p. 397. 13.—*Ib.* 8vo. i. p. 420.

Sylvia Luſcinia. *Ind. Orn.* ii. p. 506. 1.

Le Roſſignol. *Buf.* v. p. 81. t. 6. f. 1.

Nightingale. *Br. Zool.* i. N° 154.—*Ib.* fol. 100. t. 8. 1. f. 2. *Arct. Zool.* p. 416. A. *Will. Angl.* p. 220. t. 41. *Albin.* iii. t. 53.—*Ib. Song Birds*, t. p. 67. *Lath. Syn.* iv. p. 408. 1.

—*Ib. Sup.* p. 180. *Lewin, Br. Birds,* iii. t. 99. *Walc. Syn.* ii. t. 229. *Pult. Cat. Dorſet.* p. 8. *Don. Br. Birds,* v. t. 108.

This ſpecies of Warbler is about the ſize of the Sky Lark, but of a more ſlender and elegant form. Its weight is ſix drams; length near ſeven inches. Bill brown; irides hazel.

The head and upper parts of the body pale tawny; the under parts cinereous-brown; tail deep tawny-red; quills brown; the outer webs reddiſh brown; legs long, and of a light brown.

The female is rather leſs, but in plumage both ſexes are nearly alike.

This is the largeſt ſpecies of the Warbler genus; it appears with us ſometimes in April, but moſt commonly not till the beginning of May.

The females do not arrive till a week or ten days after the males; ſo that on the firſt arrival of theſe birds none but males are caught, which has given riſe to a ſuppoſition that the proportion of males are greater than thoſe of the other ſex. But was this the caſe, thoſe males who were not fortunate enough to procure a mate would ſing all the ſummer through; whereas they are all ſilent by the latter end of June. If by accident the female is killed, the male aſſumes his ſong again, and will continue to ſing very late in the ſummer, or till he finds another mate. This we have proved by taking the female on her neſt, when the male aſſumed his

uſual

uſual vociferous notes, which attracted another female. In birds that pair there is no doubt nature has given an equal proportion of both ſexes; and yet, what is extraordinary and unaccountable, if either ſex is deſtroyed before the great demand of nature is perfected, the remaining ſex generally finds a ſecond mate. This we have obſerved in many ſpecies; and Mr. White, in his *Hiſtory of Selborne*, remarks the ſame with reſpect to Partridges.

The male Nightingale, as well as all the migrative ſpecies of Warblers, never quit the place they firſt reſort to, but attract the female by their ſong. It is probable, therefore, that ſuch females who have not at firſt paired, or by accident loſe their mate, are continually wandering in ſearch of the other ſex.

The local ſituation of this bird, as well as many others, is probably occaſioned by a peculiarity of food, which may be found in ſome places, and not in others. The Nightingale is ſaid to be found only as far north as Yorkſhire; and certainly not farther weſt than the eaſtern borders of Devonſhire; although they are plentiful both in Somerſetſhire and Dorſetſhire. Why they ſhould not be found in all the wooded parts of Devonſhire and Cornwall, which appear equally calculated for their reſidence, both from the mildneſs of the air and variety of ground, is beyond the naturaliſt's penetration. The bounds preſcribed to all animals, and even plants, is a curious and important fact in the

the great works of nature. It has been obſerved that the Nightingale may poſſibly not be found in any part but where cowſlips grow plentifully; certainly with reſpect to Devonſhire and Cornwall this coincidence is juſt.

This bird reſides wholly in woods and thickets, and ſo concealed that it is ſeldom ſeen. It prepares a neſt the latter end of May, made of dry leaves, generally of the oak, and lined with dry graſs, uſually placed on the ground amongſt the ſame materials with which it is compoſed; ſo that it is not eaſily diſcovered. The eggs are four or five in number, of a uniform dark brown colour, rather larger than thoſe of a Hedge-Sparrow; their weight about forty-ſeven grains. As ſoon as the young are hatched its ſong ceaſes, and it is no more heard during the remainder of its ſtay with us. It is a miſtaken notion that this or any of the later breeding birds have a ſecond brood in the ſame ſeaſon; and we may be aſſured whenever a later brood than uſual is found ſome accident has befallen the firſt.

We have before aſſerted that the ſong of birds is the effuſion of love; and though there are ſome who frequently ſing in the autumn, and even in winter if the weather is mild, it does not follow they have a neſt; but their teſtes will always be found enlarged: theſe are only ſuch as are early breeders, the Redbreaſt, Wren, and Wood Lark.

Mr. Pennant has been very diffuſe on the ſubject of this bird, in his uſual elegant manner; but as

we

we confine our pen to the facts of natural hiſtory, as far as relates to birds in this country, we muſt refer our readers to the *Britiſh Zoology* for the more elegant and claſſical information.

The variety of this bird's notes certainly exceeds all others. Of a ſtill ſummer's night, when all is huſhed in ſilence, the vocal powers of the Nightingale is moſt diſtinctly heard: in confinement it is much longer in ſong; but it is much to be queſtioned if its notes are ſo ſtrong and pure. The young are difficult to be raiſed by hand; at firſt they are fed with meal-worms, and afterwards with ſheep's heart boiled.

The winter reſidence of this bird is ſaid to be in Aſia, and that it migrates as far north as Siberia, and is found at Kamtſchatka.

Has not been noticed in Ireland.

NIGHT-JARR. Vide *Goatſucker* and *Shriek-cinereous.*

NIGHT-RAVEN. Vide *Heron-night.*

NOPE. Vide *Finch-bul.*

NUN. Vide *Titmouſe-blue.*

NUTCRACKER.

Corvus caryocatactes. *Lin. Syſt.* i. p. 157. 10. *Gmel. Syſt.* i. p. 370. *Raii Syn.* p. 42. 5. *Will.* p. 90. t. 20. *Ind. Orn.* i. p. 164. 39.

Nucifraga. *Brif.* ii. p. 59. 1. t. 5. f. 1.—*Ib.* 8vo. i. p. 171.

Caſſe-noix. *Buf.* iii. p. 122. t 9.

Nutcracker.

Nutcracker. *Br. Zool. App.* t. 1. *Arct. Zool.* ii. p. 252. D. *Will. Angl.* p. 132. t. 20. *Edw.* t. 240. *Lath. Syn.* i. p. 400. 38.—*Sup.* p. 82. *Lewin, Br. Birds*, t. 40. *Walc. Syn.* i. t. 38. *Don. Br. Birds*, iv. t. 80.

This ſpecies of Crow is about the ſize of a Magpie; length thirteen inches.

The bill is two inches long, nearly ſtrait, and black; irides hazel. The feathers that reflect over the noſtrils are bordered with brown; upper part of the head and nape of the neck black; the general colour of the plumage ruſty brown, marked with triangular white ſpots, which are larger on the under parts; the vent white; quills and tail black, the laſt tipped with white; legs black.

Theſe birds are ſaid to vary in ſize; and a variety is mentioned ſpotted black and white.

It is a rare ſpecies in England; two inſtances only on record: one ſhot in Flintſhire, the other in Kent. Moſt frequent in Germany; found alſo in Sweden and Denmark; ſaid to viſit Burgundy in vaſt flocks.

The Nutcracker is ſaid to lay up a ſtore of acorns and nuts for winter; but we much doubt the fact, as no animal but ſuch as become partly torpid in cold weather require ſuch a proviſion. Such ſtores are moſt probably the collection of Dormice, or ſome ſuch animals, which being found by this bird is plundered. The ſame faculty is attributed to the

the Jay and Nuthatch; but they only rob the granary of mice, who frequently depoſit their winter ſtore in the hollow of a tree; ſuch as beans, peas, corn, nuts, and acorns.

This bird, whoſe partial food ſeems to be the kernels of nuts, moſt probably breaks the ſhell in the manner of the Nuthatch, by hacking a hole in it, or ſplitting the ſhell by reiterated ſtrokes of the bill, for which that part ſeems better calculated than for cracking it by compreſſion, as the Groſbeak breaks the ſtones of the haw, whoſe bill is ſhort and ſtrong, and furniſhed with muſcles of prodigious ſtrength.

It is ſaid in ſome parts to keep chiefly in the pine foreſts, probably for the ſake of the ſeed of that tree. Is alſo ſaid to make its neſt in the hole of a tree, which it perforates, or at leaſt enlarges what has already been begun by the Woodpecker: the bill ſeems not ill ſuited to this purpoſe.

Is found common in the pine foreſts of Ruſſia and Siberia, and all over Kamtſchatka.

NUTHATCH. A genus of birds, the characters of which are,

Bill ſtrait, ſtrong, ſharp pointed.
Noſtrils covered with reflected briſtles.
Tongue ſhort, horny, and jagged at the end.
Toes three forward, one backward; middle toe cloſely joined at the baſe to the others; back toe as large as the middle one.

NUTHATCH-

NUTHATCH-EUROPEAN.

Sitta Europæa. *Lin. Syſt.* i. p. 177. 1. *Gmel. Syſt.* i. p. 440. *Raii Syn.* p. 47. A. 4. *Will.* p. 98. t. 23. *Ind. Orn.* i. p. 261. 1. *Briſ.* iii. p. 588. 1. t. 29.—*Ib.* 8vo. i. p. 474.

La Sittelle, *ou* Torchepot. *Buf.* v. p. 460. t. 20.

Nuthatch. *Br. Zool.* i. N° 89. t. 38.—*Ib.* fol. 81. t. H. *Arct. Zool.* ii. N° 170. *Will. Angl.* p. 142. t. 23. *Albin.* ii. t. 28. *Lewin, Br. Birds,* ii. t. 53. *Lath. Syn.* ii. p. 648.—*Sup.* p. 117. *Walc. Syn.* i. t. 51. *Pult. Cat. Dorſet.* p. 5. *Don. Br. Birds,* iv. t. 81.

PROVINCIAL.

Nutjobber. Woodcracker.

This is the only ſpecies met with in England. It is about the ſize of a Sparrow; length near ſix inches; weight rather more than ſix drams. The bill is about three quarters of an inch long, both mandibles equally convex, and a little compreſſed ſideways; the upper one duſky, lower one whitiſh at the baſe; irides hazel. The crown of the head and whole upper parts of the body are of a bluiſh grey; from the upper mandible through the eye is a black ſtreak paſſing backwards down the neck; chin and cheeks whitiſh; breaſt and belly buff-colour; ſides and thighs ferruginous-cheſnut; quills duſky; under coverts of the tail white, margined with ferruginous; tail ſhort, compoſed of twelve feathers,

feathers, but not ſtiff, as in the Woodpeckers; the two middle ones bluiſh grey; the outer one is black, tipped with grey, ſeparated by a white bar; the ſecond nearly the ſame, but the ſpot of white is only on the inner web; the reſt are black, more or leſs marked with a little grey and white at the ends; legs pale yellowiſh; claws large; the hind one very ſtrong.

The female is lighter coloured beneath, eſpecially about the ſides and thighs.

The ſingular noiſe produced by ſome ſpecies of Woodpecks, by reiterated ſtrokes of the bill againſt the decayed limb of a tree, has been erroneouſly aſcribed to this bird.

It remains with us the whole year, but is a local bird, and not to be found in ſeveral parts of the kingdom. We have never obſerved it far north, nor ſo far weſt as Cornwall. It chiefly affects wooded and incloſed ſituations, chooſing the deſerted habitation of a Woodpecker in ſome tree for the place of nidification. This hole is firſt contracted by a plaſter of clay, leaving only ſufficient room for itſelf to paſs in and out. The neſt is made of dead leaves, moſt times that of the oak, which are heaped together without much order. The eggs are ſix or ſeven in number, white, ſpotted with ruſt-colour, ſo exactly like thoſe of the Great Titmouſe in ſize and markings, that it is impoſſible to diſtinguiſh a difference. If the barrier of plaſter at the entrance is deſtroyed when they have eggs, it

it is ſpeedily replaced; a peculiar inſtinct to prevent their neſt being deſtroyed by the Woodpecker and other birds of ſuperior ſize who build in the ſame ſituation. No perſecution will force this little bird from its habitation when ſitting; it defends its neſt to the laſt extremity, ſtrikes the invader with its bill and wings, and makes a hiſſing noiſe; and, after every effort of defence, will ſuffer itſelf to be taken in the hand rather than quit.

The Nuthatch is more expert in climbing than the Woodpecker, for it runs in all directions up and down a tree; whereas the other is never obſerved to deſcend. The ſtiff tail of thoſe birds ſupport them in the act of climbing and hacking.

The flexible tail of the Nuthatch gives it no ſuch advantage, nor does it ſeem to want it, for its moſt favourite poſition, when breaking a nut, is with the head downwards. In the autumn it is no uncommon thing to find, in the crevices of the bark of an old tree, a great many broken nut-ſhells, the work of this bird, who repeatedly returns to the ſame ſpot for this purpoſe. When it has fixed the nut firm in a chink, it turns on all ſides in order to ſtrike it with moſt advantage. This, with the common hazel-nut, is a work of ſome labour; but it breaks a filbert with eaſe. In defect of ſuch food, inſects and their *larvæ* are ſought after amongſt the moſs on trees and old thatched buildings. It is commonly met with about orchards, and is ſometimes

times ſeen in the cyder ſeaſon picking the ſeeds from the refuſe of the preſſed apples.

The note is various; in the ſpring it has a remarkable loud, ſhrill whiſtle, which ceaſes after incubation; in the autumn a double reiterated cry.

NUTJOBBER. Vide *Nuthatch.*

O.

OLIVE. Vide *Oyſter-catcher.*

ORIOLE. A genus of birds, the characters of which are,

Bill ſtrait, conic, ſharp-pointed; edges cultrated, inclining inwards; mandibles of equal length.

Noſtrils ſmall, partly covered.

Tongue divided at the end.

Toes, three forward one backward; middle one joined near the baſe to the outmoſt.

ORIOLE-GOLDEN.

Oriolus Galbula. *Lin. Syſt.* i. p. 160. 1. *Gmel. Syſt.* i. p. 382. *Ind. Orn.* i. p. 186. 45.

Galbula. *Raii Syn.* p. 68. 5. *Will.* p. 147. t. 36. 38.

Oriolus. *Brif.* ii. p. 320. 58.—*Ib.* 8vo. i. p. 247.
Le Loriot. *Buf.* iii. p. 254. t. 17.
Yellow Bird from Bengal. *Albin.* iii. t. 19.
Witwall. *Will. Angl.* p. 198.
Golden Thruſh. *Edw.* t. 185.
Golden Oriole. *Br. Zool. App.* 4. t. 4. *Lewin, Br. Birds*, ii. t. 43. *Lath. Syn.* ii. p. 449. 43. —*Ib. Sup.* p. 89. *Don. Br. Birds*, i. t. 7.

This is the only ſpecies ever found in England, a few inſtances of which are only on record.

It is about the ſize of a Blackbird; length nine inches and a half.

The bill is browniſh red; irides red. General colour of the plumage fine golden yellow; between the bill and eye a ſtreak of black; the wings black, marked here and there with yellow, and a patch of the ſame in the middle of the wing; the two middle feathers of the tail are black, inclining to olive at the baſe, the very tips yellow; the baſe half of the others black, the reſt yellow; legs lead-colour; claws black.

The female is of a dull greeniſh brown in thoſe parts where the male is black; wings duſky; tail dirty green; all but the two middle feathers yellowiſh white at the ends.

This beautiful bird is not uncommon in France, where it breeds. The neſt is curiouſly conſtructed, in ſhape like a purſe: it is faſtened to the extreme forked branches of tall trees, compoſed of fibres of hemp

hemp or ſtraw, mixed with fine dry ſtalks of graſs, and lined with moſs and liverwort.

The eggs are four or five in number, of a dirty white, marked with ſmall dark brown ſpots, which are thickeſt about the larger end. She is ſaid to be ſo tenacious of her eggs as to ſuffer herſelf to be taken with the neſt.

Their food is figs, grapes, and cherries, in the ſeaſon; at other times inſects.

Their note is a loud, ſhrill cry, which may be heard at a great diſtance.

It is found in various parts of the European continent during the ſummer months, but rarely ſo far north as England or Sweden; has been obſerved in Malta in September on its paſſage ſouthward, and on its return in the ſpring northward; ſuppoſed to winter in Africa and Aſia.

Is met with in China and Bengal, as well as at the Cape of Good Hope.

OSPREY.

Falco Haliæetus. *Lin. Syſt.* i. p. 129. 26. *Gmel. Syſt.* i. p. 263. *Ind. Orn.* i. p. 17. 30.

Morphnos, ſeu Clanga. *Raii Syn.* p. 7. 6. *Will.* p. 32.—*Ib. Angl.* p. 63.

Baldbuſardus Anglorum. *Raii Syn.* p. 16. A. 3. *Will.* p. 37.—*Ib. Angl.* p. 69. t. 6.

Baldbuzard. *Buf.* i. p. 103. t. 2. *Brif.* i. p. 440. 10. t. 34.—*Ib.* 8vo. p. 126.

Oſprey. *Br. Zool.* i. N° 46.—*Ib.* fol. p. 65.

t. A. 1. *Arct. Zool.* ii. N° 91. *Lewin, Br. Birds*, i. t. 5. *Lath. Syn.* i. p. 45.—*Sup.* p. 13. *Walc. Syn.* i. t. 5. *Pult. Cat. Dorſet.* p. 2. *White, Hiſt. Selb.* p. 97. *Don. Br. Birds*, iii. t. 70.

PROVINCIAL.

Fiſhing Hawk. Fiſhing Eagle. Bald Buzzard.

This large ſpecies of Falcon weighs between four and five pounds; length near two feet.

The bill is black; cere blue; irides yellow. The feathers on the head are brown, with white margins; the back part of the head, throat, and neck, white, with a little mixture of brown; beneath the eye is a band of brown reaching almoſt to the ſhoulder; the body is brown above, the under parts are white; the feathers of the tail are tranſverſely barred with white on the inner webs, except the two middle ones, which are wholly brown; legs ſhort, ſtrong, naked, of a bluiſh aſh-colour; claws long, much hooked, and black; the outer toe turns eaſily backwards; and, what is remarkable, the claw belonging to it is larger than that of the inner toe.

Theſe birds ſeem to vary a little in plumage; that from which Mr. Pennant took his deſcription had a ſpot of white on the joint of the wing next the body; the breaſt ſpotted with dull yellow; the greater quills black; the interior webs varied with brown and white.

This

This ſpecies is now rarely met with in England; it reſides chiefly near water, eſpecially large rivers and lakes.

Its principal food is fiſh, which it catches with great dexterity, by pouncing on them with vaſt rapidity, and carrying them off in its talons.

We are informed it is frequently ſeen about the Lake of Killarney in Ireland at particular ſeaſons, no doubt breed there. It is ſaid to make its neſt generally on the ground by the ſide of water, compoſed of flags and ruſhes; but we once ſaw the neſt of this bird on the top of a chimney of a ruin in an iſland on Loch Lomond in Scotland: it was large and flat, formed of ſticks laid acroſs, and reſting on the ſides of the chimney, lined with flags.

It is ſaid to lay three or four white eggs, of an elliptical form, rather leſs than thoſe of a Hen.

Many of the ancient writers have deſcribed this bird to have one foot ſubpalmated; a circumſtance that has never occurred in any animal; each ſide always correſponds in ſize and ſhape. The Oſprey can neither ſwim nor dive, but takes its prey as they approach the ſurface of the water.

OUZEL-MOUNTAIN. Vide *Ouzel-ring.*

OUZEL-RING.

Turdus torquatus. *Lin. Syſt.* i. p. 296. *Gmel. Syſt.* ii. p. 832. *Ind. Orn.* i. p. 343. 56.

Merula torquata. *Briſ.* ii. p. 235. 12.—*Ib.* 8vo. i. p. 223.

p. 223. *Raii Syn.* p. 65. A. 2. *Will.* p. 143. t. 37.

Le Merle à plaſtron blanc. *Buf.* iii. p. 340. t. 31.

Ring Ouzel, *or* Amſel. *Br. Zool.* N° 110. t. 46. —*Ib.* fol. 92. t. P. 1. f. 1. *Artt. Zool.* ii. p. 344. H. *Will. Angl.* p. 194. *Albin.* i. t. 39. *Lewin, Br. Birds,* ii. t. 62. *Lath. Syn.* iii. p. 46. 49.—*Ib. Sup.* p. 141. *Walc. Syn.* ii. t. 202. *Pult. Cat. Dorſet.* p. 10. *White, Hiſt. Selb.* p. 34. 56. 66. 71. 84. 96. *Don. Br. Birds,* iii. t. 61.

PROVINCIAL.

Rock, or Mountain-Ouzel. Michaelmas Blackbird. Tor-Ouzel.

This ſpecies of Thruſh is rather larger than the Blackbird; length about eleven inches.

Bill duſky; irides hazel. The general plumage is black; the feathers on the upper parts are ſlightly margined with aſh-colour, thoſe of the rump and belly deeply bordered with grey; on the breaſt is a large patch of white, ſomewhat in the ſhape of a creſcent, with the horns pointing upwards. This mark, however, is ſubject to much variation; in ſome it is of a light brown, in others wholly wanting: the former is generally the female, the latter young birds not arrived at maturity.

This is not a common ſpecies in England, but

are

are ſometimes ſeen in ſmall flocks in different parts in the ſpring and autumn, migrating from one part to another: in ſuch caſes they remain but a ſhort time in a place. It is generally believed they do not winter with us, but are known to breed in many of the barren and mountainous parts, particularly in Scotland and the north of England, as well as in ſome parts of Wales, on Dartmoor in Devonſhire, and near the Land's-end in Cornwall. We have alſo received it from the mountainous parts of Ireland.

It is ſaid to be found in many parts of the old continent, both in the warmer as well as colder regions; alſo in Africa and Aſia; and in all theſe places noticed as migratory.

The neſt is generally placed on the ground, under ſome ſmall buſh; is formed like that of a Blackbird; and the eggs, in number, ſize, and colour, are much like thoſe of that bird. It is rarely that more than one pair are ſeen near the ſame ſpot in the breeding ſeaſon. They are very clamorous when diſturbed, eſpecially when they have young. Their food is ſnails, inſects, and berries, particularly thoſe of the juniper.

The young birds, before the white on the breaſt appears, have been conſidered as a different ſpecies, under the name of Rock-Ouzel; and, in the *Catalogue of Dorſet Birds*, we are told theſe birds appear in Portland in their ſpring and autumnal flights, and are there called Michaelmas Blackbirds.

OUZEL-ROCK. Vide *Ouzel-ring.*

OUZEL-ROSE-COLOURED.

Turdus roſeus. *Lin. Syſt.* i. p. 294. 15. *Gmel. Syſt.* ii. p. 819. *Ind. Orn.* i. p. 344. 59.

Turdus Seleucis. *Gmel. Syſt.* ii. p. 837.

Merula roſea. *Raii Syn.* p. 67. 9. *Will.* p. 143. *Brif.* ii. p. 250. 20.—*Ib.* 8vo. i. p. 228.

Merle couleur de roſe. *Buf.* iii. p. 348. t. 22.

Roſe-coloured Thruſh, *or* Ouzel. *Br. Zool. App.* N° 5. t. 5. *Arct. Zool.* ii. p. 344. G. *Will. Angl.* p. 194. *Edw.* t. 20. *Lewin, Br. Birds,* ii. t. 64. *Lath. Syn.* iii. p. 50.—*Ib. Sup.* p. 142. *Walc. Syn.* ii. t. 203. *Pult. Cat. Dorſet.* p. 11.

This beautiful ſpecies of Thruſh is rather leſs than the Blackbird; length near eight inches. Bill three quarters of an inch long, a little bent, of a fleſh-colour, blackiſh at the baſe; irides pale. The head, which is creſted, neck, wings, and tail, are black, gloſſed with blue, purple, and green, as viewed in different lights; back, rump, breaſt, belly, and leſſer wing coverts, pale roſe-colour, with a few irregular dark ſpots; legs pale red; claws crooked and brown.

Theſe birds vary conſiderably with reſpect to the ſhades of the roſeate colour; ſome are much darker than others; the females are pale.

It is rarely met with in England. There are, however, ſeveral inſtances on record: about Ormſkirk in Lancaſhire one or two is ſaid to be ſhot almoſt every ſeaſon.

In

In France it is more frequent, eſpecially about Burgundy in its paſſage to other parts.

Is found alſo in many other parts of Europe, and in Aſia. Said to viſit Aleppo in great numbers in July and Auguſt in purſuit of the ſwarms of Locuſts; from which it is called Locuſt-bird, and is held ſacred by the Turks. It is alſo ſeen in abundance in the ſouth of Ruſſia and in Siberia, where it is ſaid to breed in the rocks about the rivers Don and Irtiſch; is mentioned as a bird of Switzerland and Lapland; but probably is rare in the latter.

OUZEL-TOR. Vide *Ouzel-ring*.

OUZEL-WATER.

Sturnus cinclus. *Lin. Syſt.* i. p. 290. 5. *Gmel. Syſt.* ii. p. 803. *Ind. Orn.* i. p. 343. 57.

Merula aquatica. *Raii Syn*, p. 66. A. 7. *Will.* p. 104. t. 24. *Briſ.* v. p. 252. 19.—*Ib.* 8vo. ii. p. 275.

Le Merle d'eau. *Buf.* viii. p. 134. t. 11.

Water Ouzel. *Br. Zool.* N° 111.—*Ib.* fol. 92. t. P. 1. f. 2. *Arct. Zool.* ii. p 332. B. *Will. Angl.* p. 149. t. 24. *Albin.* ii. t. 39. *Lath. Syn.* iii. p. 48.—*Sup.* p. 142. *Lewin, Br. Birds*, ii. t. 63. *Walc. Syn.* ii. t. 196. *Don. Br. Birds*, t. 24.

PROVINCIAL.

Water Crake. Water Crow. Piet.

This

OUZ

This ſingular bird, according to the lateſt authors, is placed in the Thruſh genus, to which it certainly has more affinity than to the Stare.

It is leſs than the Blackbird; length ſeven inches and a half.

The bill is three quarters of an inch long, nearly ſtrait, black; the upper mandible a little turned down at the point; irides hazel.

The upper part of the head and neck deep brown; the eyelids, chin, fore part of the neck, and breaſt, white, beneath which is a band of rufous-brown; the reſt of the upper parts, the belly, vent, and tail, are black; the feathers on the back and wings are edged with aſh colour; legs black; the tail much ſhorter than is uſual in the Thruſhes.

This ſpecies is a retired ſolitary bird, rarely ſeen but on the banks of rapid rocky rivers, or ſtreams of water, particularly in the mountainous parts, as in Scotland and Wales; is not unfrequent in Devonſhire.

In theſe places it breeds, and continues the whole year. The neſt is very large, formed of moſs and water-plants externally, and lined with dry oak-leaves; in ſhape it reſembles that of the Wren, but not ſo deep, with a dome or covering; it is uſually placed in ſome moſſy bank impending the water, in which ſituation we have frequently found it. The eggs are five or ſix in number, of a ſemitranſparent white. The tinge of bluiſh colour which they are ſaid to have is occaſioned by the yolk, and diſ-

appears

appears when they are blown. Theſe are conſiderably leſs than thoſe of the Blackbird; their weight rather more than one dram.

A pair of theſe birds, which had for many years built under a ſmall wooden bridge in Carmarthenſhire, we found had made a neſt early in May. It was taken, but had no eggs, although the bird flew out of it at the time. In a fortnight after they had completed another neſt in the ſame place, containing five eggs, which was taken; and in a month after we took a third neſt under the ſame bridge, with four eggs; undoubtedly the work of the ſame birds, as no others were ſeen about that part. At the time the laſt neſt was taken the female was ſitting, and the inſtant ſhe quitted her neſt plunged into the water, and diſappeared for a conſiderable time; at laſt ſhe emerged at a great diſtance down the ſtream. At another time we found a neſt of this bird in a ſteep projecting bank over a rivulet clothed with moſs. The neſt was ſo well adapted to the ſurrounding materials, that nothing but the old bird flying in with a fiſh in its bill would have led to a diſcovery. The young were nearly full feathered, but incapable of flight, and the moment the neſt was diſturbed, they fluttered out and dropt into the water, and to our aſtoniſhment inſtantly vaniſhed, but in a little time made their appearance at ſome diſtance down the ſtream; and it was with difficulty two out of five were taken, as they dived on being approached.

The

The aquatic habits of this bird have not eſcaped the notice of many ornithologiſts, ſome of which ſpeak of their flying under water. If, indeed, the wings being in motion can be called flying, it certainly does; but this is no more than is common to all diving birds, who, in purſuit of fiſh, or to eſcape danger, always uſe their wings to accelerate their motion. In this caſe, however, the wings are not extended, for that would retard their progreſs; but it is effected by ſhort jerks from the ſhoulder joint. Whether theſe birds can run at the bottom of the water, as ſome have aſſerted, is much to be doubted, as it is requiſite all birds ſhould uſe a conſiderable exertion to keep them under water, by reaſon of their ſpecific gravity being ſo much leſs.

It is certainly a moſt curious and ſingular circumſtance that a bird, not apparently in the leaſt formed for diving, ſhould purſue its prey under water, living chiefly on ſmall fiſh and aquatic inſects. It cannot, however, ſwim on the ſurface.

The young birds which were taken ſhewed no inclination to dive in a tub of water, but ſhewed great uneaſineſs by ſtruggling on the ſurface. They refuſed all food, and ſoon periſhed.

Theſe birds will ſometimes pick up inſects at the edge of the water; when diſturbed it uſually flirts up the tail, and makes a chirping noiſe. It ſings prettily in the ſpring.

Their

Their flight is even and rapid, like the Kingfiſher, as their wings are ſhort.

It is ſaid to be met with in many parts of Europe, and even as far as Kamtſchatka, and in ſome places is ſuppoſed to be migratory.

OWL. A genus of birds, the characters of which are,

Bill ſtrong, much hooked, but not furniſhed with a cere.

Noſtrils oblong; that and great part of the bill covered with briſtly feathers.

Head large; both ears and eyes large.

Tongue bifid.

Face ſurrounded by a ſeries of ſmall cloſe-ſet feathers.

Toes four, the outmoſt capable of being turned backwards.

Claws hooked, and very ſharp-pointed.

The exterior web of the outer quill-feather ſerrated.

The Owl is a nocturnal bird, moſtly preying by night, or at leaſt when moſt other birds are at rooſt. The pupil of the eye is capable of great extenſion; it is alſo furniſhed with a ſtrong nictitating membrane, with which it frequently covers the eye when expoſed to a ſtrong light, inſtead of cloſing the eyelid.

Some

Some ſpecies occaſionally prey by day when the weather is cloudy, but moſtly by twilight, or by the light of the moon; for they can no more ſee in total darkneſs than any other animal.

OWL-BARN. *Vide Owl-white.*

OWL-BLACK. *Vide Owl-brown.*

OWL-BROWN.

Mr. Pennant has given, in his *British Zoology*, a ſpecies under this denomination, and makes it diſtinct from the Tawny Owl; but ſays that both theſe kinds agree entirely in their markings, and differ only in the colours. In this the head, wings, and back, are of a deep brown, ſpotted with black in the ſame manner as the Tawny Owl; the coverts of the wings and the ſcapulars are adorned with ſimilar white ſpots; the exterior edges of the four firſt quill feathers in both are ſerrated; the breaſt in this is of a pale aſh-colour, mixed with tawny, and marked with oblong jagged ſpots; the feet too are feathered down to the very claws; the circle round the face is aſh-coloured, ſpotted with brown.

The colour of the irides is not mentioned by the above author; but as he ſeemed to be well acquainted with this bird, as he ſpeaks of the manners and habits to be the ſame as the Tawny Owl, certainly he would have informed us if they had differed in that particular.

Various have been the opinions concerning this bird; but we have no heſitation in declaring it to be

be a mere variety of the Tawny Owl, and not the female of that ſpecies, as ſome have conjectured; for we have found both ſexes perfectly according with the above deſcription, but moſt commonly females. What ſeems to place the matter beyond doubt is, that we have killed this bird with the Tawny Owl from the ſame neſt. The young birds are alſo ſome tawny, others brown; but the latter is moſt properly the variety, not being ſo common.

Dr. Latham has made the Tawny and Brown Owls diſtinct; but in his *Index Ornithologicus* he has expreſſed his doubts.

OWL-CHURCH. Vide *Owl-white.*

OWL-CHURN. Vide *Goatſucker.*

OWL-EAGLE. Vide *Owl-great-eared.*

OWL-FERN. Vide *Goatſucker.*

OWL-GOAT. Vide *Goatſucker.*

OWL-GREAT. Vide *Owl-great-eared.*

OWL-GREAT-EARED.

Strix Bubo. *Lin. Syſt.* i. p. 131. 1. *Gmel. Syſt.* i. p. 286. *Ind. Orn.* i. p. 51. 1. *Raii Syn.* p. 24. 1. *Will.* p. 63. t. 12. *Brif.* i. p. 471. 1.—*Ib.* 8vo. p. 139.

Le grand Duc. *Buf.* i. p. 332. t. 22.

Great-eared Owl. *Br. Zool.* i. N° 64. t. 29.—*Ib.* fol. p. 4. t. 6. *Artt. Zool.* ii. N° 114. *Will. Angl.* p. 99. t. 12. *Albin.* iii. t. 6. *Lewin, Br. Birds*, i. t. 23. *Lath. Syn.* i. p. 116. 1.—*Sup.* p. 40. *Walc. Syn.* t. 23.

PROVINCIAL.

OWL

PLUMICAL.

Great, or Eagle Owl. Great Horned Owl.

This ſpecies is almoſt equal to an Eagle in ſize. Irides bright yellow.

The plumage of the whole bird is a mixture of black-ferruginous, brown, and cinereous, elegantly varied with lines, ſpots, and ſpecks; the wings are very long, the ſecond and third quill-feathers being the longeſt; tail ſhort, marked with duſky bars; legs ſtrong, covered thickly with a light-coloured down quite to the claws, which are ſtrong, much hooked, and duſky.

This bird is rarely met with in England; a few inſtances only are on record. It has been ſhot in Yorkſhire and in Suſſex, as well as in Scotland, but is more plentiful in Norway and other parts of Europe.

Is ſaid to inhabit mountainous and rocky ſituations, and not woods, being rarely known to perch on trees.

It preys on Hares, Rabbits, Moles, and other inferior animals; and even Snakes and Toads. The eggs are ſaid to be larger than thoſe of a Hen, mottled like the bird; and that only two are found in a neſt.

Dr. Latham has mentioned three varieties, one of which has the legs bare of feathers; and he conſiders the Black-winged Horn Owl of Albin to be a mere variety.

OWL-GREAT-HORNED. Vide *Owl-great-eared.*

OWL-GREY. Vide *Owl-tawny.*

OWL-HISSING. Vide *Owl-white.*

OWL-HORN. Vide *Owl-long-eared.*

OWL-HORN-BLACK-WINGED. Vide *Owl-great-eared.*

OWL-IVY. Vide *Owl-tawny.*

OWL-LITTLE.

Strix paſſerina. *Lin. Syſt.* i. p. 133. 12. *Gmel. Syſt.* i. p. 296. *Ind. Orn.* i. p. 65. 46.

Noctua minor. *Raii Syn.* p. 26. 6. *Will.* p. 69. t. 13. *Briſ.* i. p. 514. 5.—*Ib.* 8vo. p. 150.

La Chevêche, *ou* petite Chouette. *Buf.* i. p. 377. t. 28.

Little Owl. *Br. Zool.* i. N° 70.—*Ib.* fol. 73. t. B. 5. *Arct. Zool.* ii. N° 126. *Will. Angl.* p. 105. t. 13. *Edw.* t. 228. *Albin.* ii. t. 12. *Lewin, Br. Birds,* i. t. 29. *Lath. Syn.* i. p. 150. 40. *Walc. Syn.* i. t. 28.

This elegant little ſpecies of Owl is the ſmalleſt that has been found in England.

The length little more than eight inches; ſize not much ſuperior to that of a Blackbird. The bill is duſky, with a yellowiſh tip; irides pale yellow.

The head and upper parts are brown, tinged with olive; the former, with the wing coverts, are ſpotted with white; the feathers that compoſe the circle round the face are white, tipped with black;

under part of the body white, ſpotted with brown; the tail is brown, barred tranſverſely with rufous, and tipped with white; but in this part they are ſubject to vary, as Mr. Pennant and Dr. Latham both ſay the tail is barred with white likewiſe: the legs are covered with down of a grey colour; claws brown. Other varieties of this bird are alſo ſpoken of with the colours darker, as well as the irides being black. It is perhaps difficult to ſay what changes climate may produce. We know ſeaſon alone in the ſame country effects extraordinary changes; but we have ever held the colour of the eyes as a characteriſtic mark, which in adults never change. It is true, many birds, before they arrive at maturity, have dark irides, which afterwards become yellow. This might be the caſe with the bird here mentioned. The Peregrine Falcon has been taken with yellow irides, but for the firſt two or three years are duſky. A young Herring Gull, which we have at this time, only begins to appear yellowiſh in that part at two years and a half old. The eye, therefore, being ſubject to ſuch a certain change by age, muſt be conſidered as the effect of maturity when they become of a lighter colour.

The Little Owl is a very rare ſpecies in England. In France it is ſaid to frequent ruined edifices. It makes a neſt in the holes of rocks and walls, and lays five or ſix eggs, ſpotted with yellowiſh and white.

It is ſaid to fly by day, and to give chace to

ſmall

ſmall birds; but its principal food is mice. Is ſaid to build in chimnies in Carniola; and Mr. Edwards mentions two having been taken in England by coming down chimnies. It is found at Hudſon's Bay, and has been received from the Weſt Indies.

OWL-LONG-EARED.

Strix otus. *Lin. Syſt.* i. p. 132. 4. *Gmel. Syſt.* i. p. 288. *Raii Syn.* p. 25. A. 2. *Will.* p. 64. t. 12. *Ind. Orn.* i. p. 53. 7.

Aſio. *Briſ.* i. p. 486. 4.—*Ib.* 8vo. p. 142.

Le Hibou, *ou* moyen Duc. *Buf.* i. p. 342. t. 22.

Horn Owl. *Albin.* ii. t. 10. *Will. Angl.* p. 100.

Long-eared Owl. *Br. Zool.* i. N° 65. t. 30.—*Ib.* fol. t. B. 4. f. 1. *Arct. Zool.* ii. N° 115. *Lewin, Br. Birds,* i. t. 24. *Lath. Syn.* i. p. 121.—*Sup.* p. 42. *Walc. Syn.* i. t. 24. *Pult. Cat. Dorſet.* p. 3.

This beautiful ſpecies is nearly fifteen inches in length; weight nine or ten ounces.

The bill is black; irides orange-yellow. The feathers, of a hair-like form, that cover the bill, are white, with black ſhafts; over the eye and round the angle next to the bill is black; cheeks pale rufous; over the eyes are two tufts of feathers, erect like ears, compoſed of ſix feathers that appear in front, gradually lengthening from the firſt to the laſt; the hindmoſt is an inch and a half long, black,

bordered with dull yellow; the circle round the face is white, ſpeckled with black and rufous; thoſe that immediately cover the ears are tipped only with black, forming a ſemicircular line of that colour; the general colour of the bird is an ochraceous yellow, elegantly ſtreaked above with black, and ſpeckled with the ſame, aſh-colour, and white; beneath, the feathers are tinged with light ferruginous, ſtreaked with black down the ſhafts; the quills are barred with black and cinereous; on the primores are two bars of dull yellow; the tail is barred, and ſpeckled with duſky and cinereous; legs and toes are covered with down of a yellowiſh buff-colour; claws duſky.

This deſcription is taken from a female; the male differs in nothing but in being rather leſs. The wings of this ſpecies are very long, reaching beyond the tail when cloſed, and croſſing each other at the points; the ſecond feather is the longeſt.

This is by no means ſo common as the Tawny or White Owls; and though it is frequently taken in England, very little is known of its habits. It is ſaid to make no neſt, but to take poſſeſſion of that of a Magpie or Crow; and that it lays four or five eggs. It remains with us the whole year, having killed them both in ſummer and winter; of courſe they muſt breed with us. An ingenious friend informed us he found an Owl's neſt in a tree covered with ivy, which had three white eggs; and that from the appearance of the bird as it flew out he had

had no doubt it was this ſpecies. It muſt be obſerved, the other Owls of this country are never known to build amongſt the branches of trees; from which we may conclude the neſt above mentioned was of this ſpecies.

This bird frequents large woods, and wooded tracts; is partial to fir, box, or holly plantations, where it more readily conceals itſelf by day amongſt the evergreen foliage, as well in winter as in ſummer.

The one before us was killed in the winter as it was flying out of a large holly-buſh in Cornwall; and we have ſeen others in ſeveral parts of the kingdom.

Its principal food is mice, and ſometimes ſmall birds taken at rooſt. Is never ſeen to fly in the day except diſturbed.

The note of this ſpecies is unknown; moſt of the genus make a ſcreeching noiſe, but the Tawny Owl is the only one with us that is known to hoot, and is ſo commonly heard in the evening.

It is ſaid to be far from uncommon in France, and many other parts of Europe, as far as the northern parts of Ruſſia.

It is alſo found in ſome parts of America, and is common at Hudſon's Bay.

OWL-SCREECH. Vide *White* and *Tawny-Owls.*

OWL-SHORT-EARED.

Strix brachyotos. *Gmel. Syſt.* i. p. 289. *Ind. Orn.* i. p. 55. 11.

Short-eared Owl. *Br. Zool.* i. N° 66. t. 31.—*Ib.* fol. p. 71. t. B. 3. & B. 4. f. 2. *Artt. Zool.* ii. N° 116. *Lath. Syn.* i. p. 124. 9.—*Sup.* p. 43. *Lewin, Br. Birds,* i. t. 25. *Walc. Syn.* i. t. 25. *Don. Br. Birds,* t. 35. *Pult. Cat. Dorſet.* p. 4.

PROVINCIAL.

Mouſe Hawk. Woodcock Owl. Hawk Owl.

This ſpecies weighs eleven ounces; length near fifteen inches. Bill duſky; irides bright yellow. The feathers immediately ſurrounding the eyes are black; thoſe that cover and ſurround the bill white; the wreath round the face beautifully ſpeckled with light ferruginous, black, and white, except at each ear, where it is wholly black; on the top of the head, above each eye, is a tuft of feathers, which it can erect at pleaſure, the foremoſt of which are black on the outer webs, and white on the inner; the reſt of the head, neck, back, and ſcapulars duſky, bordered more or leſs with light ferruginous; breaſt and belly yellowiſh white, ſtreaked with duſky down the ſhafts; the greater quills are light ferruginous on the outer webs; the three firſt have a ſingle bar of black each, and deeply tipped the ſame; the others have two bars each, their tips brown, inclining to greyiſh; the inner webs have one, or part of an irregular bar; the coverts of the primores black; on the coverts

of

of the ſecondaries are ſeveral large ſpots of white; the ſecond feather in the wing is the longeſt; the feathers of the tail are light ferruginous, croſſed with four duſky bars on the ſix middle ones, and marked with duſky ſpots on the yellow bars of the two middle feathers; the bars on the outer feathers are not ſo numerous or ſo perfect, and the yellow is ſhaded off to almoſt white on the exterior feathers, which have only irregular circles of duſky brown on the inner webs; the legs are covered down to the claws with light yellow feathers.

The above deſcription is taken from a male killed near Briſtol.

The female is rather leſs bright in colour, and ſomewhat ſuperior in ſize.

This bird is diſtinguiſhed from all the other ſpecies by the ſmallneſs of its head; which has occaſioned it to be called in ſome places by the name of Hawk Owl, or Mouſe Hawk.

To Mr. Pennant the Britiſh *Fauna* is indebted for the firſt diſcovery of this bird. But that excellent naturaliſt has deſcribed it to have only one feather on each ſide the head which is erectable; which miſtake has been followed by all other ornithologiſts.

It muſt, however, be remarked, that theſe tufts, or ears, are never erected but when the bird is in a quieſcent ſtate. A few years ſince one of this ſpecies was taken alive in a Lark-net, making a pounce at the decoy bird; and we bought it of the

birdcatcher in order to observe its manners. In a few months it became tolerably tame, and when hungry would take food from the hand. It was mostly fed with small birds and mice, but would eat any raw meat, which it first took in its bill, and immediately placed in its talons, and devoured it by piecemeal.

When it was asleep or undisturbed the aurated feathers were very distinguishable, standing above the rest about half an inch; but on being disturbed they were instantly depressed, and the head apparently enlarged by the feathers round the face being somewhat raised. Upon minute examination these tufts were found to consist of a series of feathers very little if any thing longer than the rest. While we were in possession of this we received a dead specimen, in which one feather was visibly longer than the rest in the tuft: this might have led to the mistake before mentioned.

The Short-eared Owl comes to us in October, about the time the Woodcock makes its appearance, and departs at the same time with that bird in March; hence the name of Woodcock Owl.

With us this bird is observed never to perch on a tree, but generally hides itself in long grass, fern, or the like; and seems partial to open, barren situations. When disturbed it flies a little way, and lights again on the ground. In dusky weather it will prey by day, and sometimes fly at small birds as well as mice. It is a bold bird, but seems contented

contented in confinement. That above mentioned never drank for the ſix months it was in our poſſeſſion. How long it lived after it was given away, we never heard. It may, however, be preſumed, contrary to the general opinion, that Owls, as well as all predacious birds, bear confinement, whether taken young or old; which we have experienced as well in the Tawny and White Owls as in this.

Is ſuppoſed to breed in the Orknies, and probably in Norway; viſits Hudſon's Bay in May; makes a neſt of dry graſs on the ground, and lays three or four white eggs.

We have great reaſon to believe this bird is the Chouette, or Grande Chereche, of Buffon, and perhaps has been noticed by other authors; but from the circumſtance of the ears not being mentioned, which is not diſcoverable in a dead ſpecimen, confuſion has ariſen.

OWL-TAWNY.

Strix ſtridula. *Lin. Syſt.* i. p. 133. 9. *Gmel. Syſt.* i. p. 294. *Ind. Orn.* ii. p. 58. 25.

Strix Aldrov. *Raii Syn.* p. 25. A. 2. *Will.* p. 65. t. 14. *Briſ.* i. p. 500. 1.—*Ib.* 8vo. p. 146.

Le Chathaunt. *Buf.* i. p. 362. t. 25.

Common Brown, *or* Ivy Owl. *Will. Angl.* p. 102. t. 14. *Albin.* i. t. 9.

Tawny Owl. *Br. Zool.* i. N° 68.—*Ib.* fol. 7. t. B. 3. *Arct. Zool.* ii. p. 237. B. *Lewin, Br. Birds*, i. t. 27. *Walc. Syn.* i. t. 27. *Don. Br. Birds*, v. t. 121. *Pult. Cat. Dorſet.* p. 4.

Strix

ſometimes in barns, which laſt it frequents for the ſake of mice; and as it is a better mouſer than a cat, the farmer holds it in great eſtimation, and leaves a hole in his barns and granary for its egreſs.

It prepares very little neſt, and ſometimes depoſits its eggs on the decayed wood, which are two in number, and rarely three, of a dull white, not gloſſy, and of a roundiſh form. The young are covered with a light-coloured down; are at firſt very ſhy, but ſoon become tame if fed by hand. If put out of doors within the hearing of the parent birds, retain their native ſhyneſs, as the old ones viſit them at night, and ſupply them with ample proviſion; amongſt which we have found young Hares, young Rats, and Mice; but the laſt is their principal food.

We have taken this bird in its mature ſtate as well as young, and found no difficulty in either caſe of preſerving them alive. They were never obſerved to drink; and indeed for many months together had no water offered them.

This is the only ſpecies known to hoot; beſides which it makes a diſagreeable ſcreaming noiſe.

By ſome it is called Wood-Owl and Screech-Owl. It is a great deſtroyer of young Pigeons, and frequently reſorts to pigeon-houſes for that purpoſe.

OWL-WHITE.

Strix flaminea. *Lin. Syſt.* i. p. 133. 8. *Gmel. Syſt.* i. p. 293. *Raii Syn.* p. 25. A. 1. *Will.* p. 104. t. 13. *Ind. Orn.* i. p. 60. 28. *Briſ.* i. p. 503. 2.—*Ib.* 8vo. p. 147.

Aluco minor Aldrov. *Will.* p. 67. t. 13.

L'Effraie. *Buf.* i. p. 366. t. 26.

White Owl. *Br. Zool.* i. N° 67.—*Ib.* fol. p. 71. t. B. *Arct. Zool.* N° 124. *Will. Angl.* p. 104. t. 13. *Albin.* ii. t. 11. *Lewin, Br. Birds,* i. t. 26. *Lath. Syn.* i. p. 138. 26.—*Sup.* p. 46. *Walc. Syn.* i. t. 26. *Don. Br. Birds,* v. t. 113. *Pult. Cat. Dorſet.* p. 4.

PROVINCIAL.

Barn-Owl. Gillihowter. Howlet. Madge-Howlet. Church-Owl. Hiſſing-Owl. Screech-Owl.

The weight of this ſpecies is about eleven ounces; length between thirteen and fourteen inches. The bill is light coloured; irides duſky. The feathers round the eyes yellowiſh; the circle round the face white; the upper parts of the body, coverts, and ſecondary quills, pale dull yellow, with two white and two grey ſpots placed alternately on each ſide the ſhafts; prime quills dull yellow on the outer webs, the inner white, marked on each ſide with four black ſpots; the whole under ſide white; the interior webs of the tail-feathers are white, the exterior marked with obſcure duſky bars; legs covered

Angl. p. 297. *Albin.* i. t. 78. *Hayes, Br. Birds,* t. 12. *Lath. Syn.* v. p. 219. t. 84. *Lewin, Br. Birds,* v. t. 188. *Walc. Syn.* ii. t. 166. *Don. Br. Birds,* iii. t. 62. *Pult. Cat. Dorſet.* p. 15.

PROVINCIAL.

Pienet. Olive.

The weight of the Pied Oyſter-catcher is about ſeventeen ounces; length ſixteen inches. Bill three inches long, ſtrait, compreſſed, the tip wedge-ſhaped, the colour orange; irides crimſon; orbits orange-yellow.

The head, neck, upper part of the back, ſcapulars, and leſſer wing coverts, black; a ſmall ſpot of white under the eyelid, and a creſcent of the ſame acroſs the throat; but this laſt is by no means a general character, for in more than twenty ſpecimens we never have found it, although ſuch are to be ſeen in different collections. The greater coverts are white, the next above them tipped with white; the quills are black, with more or leſs white on the inner webs; the lower part of the back, rump, breaſt, and under parts, are white; the baſe half of the tail is white, the end black; legs red-orange; claws black.

This bird is common on our ſhores, but never quits the ſea-coaſt. In winter they aſſemble in ſmall flocks. Their principal food is marine

inſects and ſhell-fiſh; and from their adroitneſs in getting at an oyſter they have taken this name. This, however, can only be done when the ſhells are partly open; it is then the bill is inſerted to kill the oyſter.

It makes no neſt, but depoſits its eggs on the bare ground above high-water mark; generally four, of an olivaceous-brown, blotched with black, ſomewhat larger than that of the Lapwing. At the time of incubation the male is very watchful, and upon the leaſt alarm flies off with a loud ſcream, upon which the female inſtantly runs from her eggs to ſome diſtance, and then takes wing. It is a ſhy bird, but when the young are hatched becomes bolder. The young is capable of running as ſoon as it quits the egg, and is led by the parent birds to its proper food.

Theſe birds are ſubject to ſome variation in plumage. In moſt the white under the chin is wanting; in ſome that mark is mottled with black, and the white ſpot under the eyelid wanting. Whether theſe are varieties, or depending on age or ſex, has not been thoroughly aſcertained; perhaps both: but we have killed both ſexes at all ſeaſons without the white on the throat. Dr. Latham obſerves it is eaſily tamed when taken young, and have been known to attend the Ducks and other poultry to feed and ſhelter at night.

The only known ſpecies of this genus.

P.

PARTRIDGE. A genus of birds, the characters of which are,

Bill convex, ſtrong and ſhort.

Noſtrils covered above with a callous prominent rim.

No naked ſkin above the eyes; but in ſome ſpecies more or leſs warty excreſcences round the eyes.

Legs naked, ſome ſpecies furniſhed with ſpurs behind.

Tail ſhort.

PARTRIDGE-COMMON.

Tetrao Perdix. *Lin. Syſt.* i. p. 276. 13. *Gmel. Syſt.* ii. p. 757.

Perdix cinerea. *Raii Syn.* p. 57. A. 2. *Will.* p. 118. t. 28. *Ind. Orn.* ii. p. 645. 9. *Brif.* i. p. 219. 1.—*Ib.* 8vo. i. p. 61.

La Perdrix griſe. *Buf.* ii. p. 401.

Common Partridge. *Br. Zool.* i. N° 96.—*Ib.* fol. 86. t. M. *Arct. Zool.* ii. p. 319. A. *Will. Angl.* p. 166. t. 28. *Albin.* i. t. 27. *Lath. Syn.* iv. p. 762. 8. *Lewin, Br. Birds,* iv. t. 136. *Walc. Syn.* ii. t. 184. *Pult. Cat. Dorſet.* p. 7.

This

This ſpecies is ſo well known as to require very little deſcription. Length about thirteen inches; weight fifteen ounces. The bill is bluiſh brown; irides hazel. General colour of the plumage cinereous brown and black mixed; ſome of the back and coverts ſtreaked with buff; ſides of the head bright ruſt-colour; behind the eye a naked, red warty ſkin; on the breaſt a deep bay-coloured mark, in form of a horſeſhoe.

The tail conſiſts of ſixteen feathers of a bright ruſt-colour, except the four middle ones, which are like the back; legs bluiſh grey.

The female weighs about fourteen ounces; the head is leſs bright, and the coverts of the ears inclining to grey; the horſeſhoe on the breaſt is white for the firſt year, afterwards more or leſs like the male, and by the third year is no longer a mark of diſtinction; whereas by the head the ſexes may always be known; the bare ſkin behind the eye is leſs conſpicuous, and very little red.

It has been long an eſteemed opinion amongſt ſportſmen as well as naturaliſts, that the female Partridge had none of the bay feathers on the breaſt like the male. This, however, is a miſtake, as we have proved by the unerring rule of diſſection; for happening to kill nine old birds one day, with very little variation as to the bay markings on the breaſt, we were led to open them all, by which we diſcovered five of them were females; and by re-examining the plumage found the males could only

be known by the ſuperior brightneſs of colour about the head, which alone ſeems to be the mark of diſtinction after the firſt or ſecond year.

This bird is found in all parts of Great Britain where corn is cultivated, but never at any great diſtance from arable land: upon the barren mountains of the north it is never ſeen. In Scotland the Partridge, the Grous, and the Ptarmigan, each have their diſtrict: the firſt is only found in the glens or vallies; the ſecond on the firſt hills; the laſt on the ſummits of the higheſt mountains; and it is very rare that they intrude upon each other; though we have killed the three ſpecies in the ſame day.

The Partridge is very prolific, laying from twelve to twenty eggs. It makes no neſt, but ſcrapes a ſmall hollow in the ground, placing a few contiguous fibres therein to depoſit its eggs on; theſe are of a light brown colour. The old birds ſit very cloſe on their eggs when near hatching; and we have been informed by a gentleman of veracity that he ſaw this bird taken with her eggs on the point of hatching, and carried in a hat to ſome diſtance; and that ſhe continued to ſit in confinement, and brought out her young. The time of incubation is three weeks; the young leave the neſt in twelve hours; the parent birds are equally tenacious of their young, and lead them immediately to ant-hills, on the eggs of which they principally feed at firſt.

Theſe

Theſe birds flock together in broods till the returning ſpring. Sometimes three or four coveys will aſſemble in winter, and are then exceedingly ſhy. In vain may the ſportſman purſue them, unleſs by ſurpriſe he can break and ſcatter the covey.

About the middle of February they begin to pair. In June they lay, and the young are excluded about the middle of July; and in about three weeks are capable of flying.

This is one of the few birds under the denomination of game, and protected by the legiſlature. But notwithſtanding many ſevere penal laws are enacted for its preſervation, it is a query if the breed is not decreaſed thereby. The great demand of the luxurious and wealthy, and the high price given for theſe birds, is too great a temptation to the poacher; and he riſks his liberty to ſupply the tables of the rich. Thus the ſeveral laws reſpecting game are ineffectual; they only ſerve to enhance the price, and hold out a temptation for the huſbandman to ruin himſelf and family; whereas if the penalty was made much greater, and that wholly confined to the buyer, it would fall upon the tempters, who ſhould be the only perſons obliged by law to pay for their luxuries. The late act of parliament for confining the killing of this ſpecies between the fourteenth of September and the twelfth of February only tended, in ſome degree, to preſerve them from the fowler, but not from the nets of the poacher.

There are ſeveral varieties mentioned of this bird, but they are merely accidental defects. Four entirely white were taken alive out of one covey, by order of the late Lord Courtenay, at Powderham in Devonſhire; three of which we ſaw. Theſe were intended to be given their liberty in the ſpring, in hopes of propagating the breed; but we never heard if it had the effect, or what became of them.

PARTRIDGE-GUERNSEY.

Tetrao rufus. *Lin. Syſt.* i. p. 276. 12. *Gmel. Syſt.* ii. p. 756. *Ind. Orn.* ii. p. 647. 12.

Perdix græca. *Raii Syn.* p. 57. A. 5. *Will.* p. 121. t. 29. *Briſ.* i. p. 241. 12. t. 23. f. 1. —*Ib.* 8vo. i. p. 67.

La Bartavelle. *Buf.* ii. p. 420.

Greek, *or* Red Partridge. *Lath. Syn.* iv. p. 767. 12. *Will. Angl.* p. 169. *Albin.* i. p. 27.

Perdix rubra. *Briſ.* i. p. 236. 10.—*Ib.* 8vo. i. p. 66.

—— ruffa. *Raii Syn.* p. 57. A. 5. *Will.* p. 167. 29.

Perdix rouge. *Buf.* ii. p. 431. t. 15.

Guernſey Partridge. *Will. Angl.* p. 167. t. 29. *Albin.* i. t. 29. *Lath. Syn.* iv. p. 768.—*Sup.* p. 220.

Perdix rubra barbarica. *Briſ.* i. p. 239. 11.—*Ib.* 8vo. i. p. 67.

Perdrix rouge de Barbarie. *Buf.* ii. p. 445.

Barbary

Barbary Partridge. *Edw.* t. 70. *Lath. Syn.* iv. p. 770.

Guernſey Partridge. *Lewin, Br. Birds*, iii. t. 137. *Pult. Cat. Dorſet.* p. 7.

It is much to be doubted whether this ſpecies has any claim in the Britiſh *Fauna*, for it does not ſeem even naturaliſed like the Pheaſant; and moſt probably the few that have been met with at large had only eſcaped from the *ornithones* of the curious. As it has, however, been killed in a ſtate of liberty we ſhall give it a place in this work.

This bird is rather larger than the common ſpecies. Bill, irides, and legs, red. The upper part of the head is red-brown; greyiſh on the forehead; chin and throat white, encircled with black; over each eye a band of white; fore part and ſides of the neck cinereous, ſpotted with black; back, wings, and rump, grey-brown; breaſt pale aſh-colour; belly rufous; ſides marked with lunular ſtreaks of black, white, and orange; the tail compoſed of ſixteen feathers of a rufous-colour, except the ſix middle ones, which are more or leſs grey-brown.

The habits of this ſpecies differ ſomewhat from the preceding. This frequently perches on a tree, and will breed in confinement, which the other is never known to do.

Why this ſhould be called Guernſey Partridge we cannot imagine, ſince we are credibly informed

it is very rare in that iſland. The common ſpecies breed there, but are ſcarce; whether this ever bred there is uncertain, though they are known to breed in the iſland of Jerſey. It is alſo found in various parts of the European continent, Aſia, and Africa, and is called by the name of Red-legged Partridge.

PARTRIDGE-WHITE. Vide *Ptarmigan.*

PELICAN. A genus of birds, whoſe characters are,

Bill long, ſtrait, hooked at the end.

Noſtrils none; face and gullet deſtitute of feathers, the latter capable of great diſtention.

Toes four, all webbed together.

The only Britiſh ſpecies of this genus are the Corvorant, Shag, and Gannet.

PENGUIN. Vide *Auk-great.*

PETREL. A genus of aquatic birds, the characters of which are,

Bill ſtrait, but bent at the end.

Noſtrils contained in one tube.

Legs naked a little above the knee.

Toes three, forward, webbed; a ſpur behind inſtead of a back-toe.

PETREL-LITTLE. Vide *Petrel-ſtormy.*

PETREL-MANKS. Vide *Shear-water.*

PETREL-

PETREL-STORMY.

Procellaria pelagica. *Lin. Syſt.* i. p. 212. *Gmel. Syſt.* ii. p. 561. *Ind. Orn.* ii. p. 826. 19. *Briſ.* vi. p. 140, 1. t. 13. f. 1.—*Ib.* 8vo. ii. p. 398.

Oiſeau de tempête. *Buf.* ix. p. 327. t. 23.

Stormfinch. *Will.* p. 306.—*Ib. Angl.* p. 395.

Stormy Petrel. *Br. Zool.* ii. N° 259. t. 91.—*Ib.* fol. 146. t. L. 5. *Arct. Zool.* ii. N° 464. *Edw.* t. 90. *Albin.* iii. t. 92. *Borlaſ. Cornw.* p. 247. t. 29. *Lath. Syn.* vi. p. 411. 18.—*Sup.* p. 269. *Lewin, Br. Birds,* vi. t. 219. *Pult. Cat. Dorſet.* p. 19. *Walc. Syn.* i. t. 91.

PROVINCIAL.

Little Petrel. Witch. Mother Cary's Chicken.

This ſpecies is about the ſize of a Swallow, and in its general appearance and flight not unlike that bird. Length about ſix inches; colour of the plumage above is black; beneath ſooty brown, or duſky; ſome of the ſecondary quills are white at the ends; rump and vent white; tail white at the baſe of the inner webs; legs black.

The Stormy Petrel is rarely ſeen on our ſhores, except in ſome of the northern iſlands, where it is ſaid to breed in the holes of rocks. It is a pelagian bird, living at all other ſeaſons remote from land. They are ſeen in numbers all over the vaſt Atlantic Ocean, and will frequently follow a ſhip for

for many days; ſuppoſed for the purpoſe of ſheltering themſelves in the wake of the veſſel, but more probably for the ſake of the various matter thrown overboard, for they will ſtoop and pick up bits of biſcuit or meat. Suppoſed to be ſeen only before ſtormy weather, and of courſe not a welcome viſitor to the ſailors. Are ſaid to be excellent divers, and capable of remaining under water a great while. The body is of ſo oily a nature that if a wick is drawn through from the mouth to the vent and lighted will burn as a lamp; and is ſaid to be actually uſed for that purpoſe in the Ferroe iſlands. Some few inſtances are recorded of its being killed far inland: one is mentioned in Latham's *Synopſis* to have been ſhot at Oxford. We are alſo informed that ſome are annually ſeen on the weſtern part of the peninſula of Cornwall, about Marazion and Penzance; in the former of which we ſaw one that was taken.

PETTYCHAPS-GREATER.

Motacilla hortenſis. *Gmel. Syſt.* ii. p. 955.
Sylvia hortenſis. *Ind. Orn.* ii. p. 507. 3.
Curruca. *Brif.* iii. p. 372. 2.—*Ib.* 8vo. i. p. 414.
La Fauvette. *Buf.* v. p. 117. t. 7.
Pettychaps. *Lath. Syn.* iv. p. 413. 3. *Lewin, Br. Birds,* iii. t. 100. *Walc. Syn.* ii. t. 230.

Length ſix inches; weight about five drams. Bill duſky above; baſe of the under mandible

yellowiſh; irides hazel; orbits white. General colour of the plumage above light brown, inclining to olive; below the ear is a daſh of aſh-colour, throat, neck, breaſt, and ſides, dirty white, inclining to brown on the two laſt; belly and under tail coverts white; quills and tail feathers duſky, edged with olive; legs bluiſh brown.

This ſpecies of Warbler, which is not very plentiful in England, was firſt diſcovered in Lancaſhire, and ſent from thence to Dr. Latham by Sir Aſhton Lever. However, ſince it has been better known, it is found to arrive in ſeveral of the ſouthern counties about the latter end of April or beginning of May. Its ſong is little inferior to that of the Nightingale. Some of the notes are ſweetly and ſoftly drawn; others quick, lively, loud, and piercing, reaching the diſtant ear with pleaſing harmony, ſomething like the whiſtle of the Blackbird, but in a more hurried cadence: ſings frequently after ſunſet. This bird chiefly inhabits thick hedges, where it makes a neſt compoſed of gooſe graſs, and other fibrous plants, flimſily put together, like that of the common White Throat, with the addition ſometimes of a little green moſs externally: the neſt is placed in ſome buſh near the ground. It lays four eggs, about the ſize of a Hedge Sparrow's, weighing about thirty-ſix grains, of a dirty white, blotched all over with light brown, moſt numerous at the larger end, where ſpots of aſh-colour alſo appear.

In Wiltſhire, where we have found this ſpecies not

not uncommon, it reſorts to gardens in the latter end of ſummer, together with the White Throat and Blackcap, for the ſake of currants and other fruit.

The Pettychaps of Mr. Pennant ſeems to correſpond with this, except that the inſide of the mouth of this is more inclined to yellow than red. That author alſo ſays, in Yorkſhire his is called the Beambird, from its neſting under beams in outbuildings. But as he quotes the *Motacilla Hippolais* of Linnæus, we muſt conclude he means the Leſſer Pettychaps of Latham's *Synopſis*, and of this work. But in either he muſt have been deceived as to its making a neſt in outbuildings; and we are rather inclined to think the habits of the Spotted Flycatcher has been given him with his Pettychaps; that bird having obtained the name of Rafter in ſome parts from the circumſtance of building its neſt on rafters and beam-ends in old buildings and outhouſes.

PETTYCHAPS-LESSER.

Motacilla Hippolais. *Lin. Syſt.* i. p. 330. 7. *Gmel. Syſt.* ii. p. 954.

Sylvia Hippolais. *Ind. Orn.* ii. p. 507. 4.

Ficedula ſeptima Aldrovandi. (Pettychaps.) *Raii Syn.* p. 79. A. 7. *Will.* p. 158.—*Ib. Angl.* p. 216.

Leſſer Pettychaps. *Br. Zool.* i. N° 149? *Arct. Zool.* ii. p. 418. G.—*Ib. Sup.* p. 64. *Lath. Syn.* iv. p. 413. 3*. *Lewin, Br. Birds*, iii. t. 101.

t. 101. *Walc. Syn.* ii. t. 231. *Pult. Cat. Dorſet.* p. 9.

Weighs about two drams; the length exceeds four inches and a half.

This ſpecies is inferior in ſize to the Yellow Wren. In its plumage it ſo much reſembles that bird that we ſhall only make mention here of any eſſential marks of diſtinction, and refer our reader to the Yellow Wren.

Its general colour is not ſo much tinged with yellow, and the legs are duſky, which in the other are brown.

Theſe two birds have been, and are frequently confounded, and with them the Wood Wren of this work; but this laſt is at once diſtinguiſhed by the under tail coverts being a pure white, and the plumage of a more lively green on the upper parts than either of the others. The neſt, eggs, and notes, will be found alſo different by conſulting and comparing the hiſtory of each. This is the firſt ſpecies of Warbler that pays its annual viſit to us in the ſpring, and is generally heard on or before the firſt of April repeating its ſong, if it may be ſo called, for it conſiſts only of two notes, which ſeem to expreſs the words *chip, chop*, four or five times ſucceſſively. It is a buſy, reſtleſs bird, always active amongſt the trees and buſhes in ſearch of inſects.

The plumage of the ſexes are alike.

The

the cheek; quills black; tail duſky, edged with aſh-colour; legs compreſſed, like the Diver's, of a light colour; toes ſcallopped; membranes ſerrated on their margins.

This deſcription is taken from a ſpecimen killed at a pond at Alderton in Wiltſhire, now in my muſeum. The ſex my notes do not inform me; but it ſeems to be ſubject to ſome variety, either from ſex or age. Dr. Latham mentions one to have the back, rump, and ſcapulars, deep dove-colour; legs black.

It ſeems to be a ſolitary and ſcarce bird with us. We once had an opportunity of ſeeing one ſwimming in a ſmall pool of water left by the tide on the Suſſex coaſt. It was continually dipping its bill into the water, as if feeding on ſome inſects, and ſo intent as to ſuffer us to approach within a few yards. It never attempted to dive, and when diſturbed flew only a ſmall diſtance, very like the Purre. Is ſaid to inhabit the northern parts of Europe, and to congregate about the borders of the Caſpian Sea.

PHALAROPE-RED.

MAS.

Tringa hyperborea. *Lin. Syſt.* i. p. 249. 9. *Gmel. Syſt.* i. p. 675.

Phalaropus hyperboreus. *Ind. Orn.* ii. p. 775. 1.

——— cinereus. *Briſ.* vi. p. 15, 2.—*Ib.* 8vo. ii. p. 362.

Larus fidipes alter noſtras. *Raii Syn.* p. 132. A. 7.
Will. p. 270.
Le Phalarope cendré. *Buf.* viii. p. 224.
Cock Coot-footed Tringa. *Edw.* t. 143.
Red Phalarope. *Br. Zool.* ii. N° 219. t. 76.
Lath. Syn. v. p. 270. 1. (maſ.) 271. 2. (femin.)
Lewin, Br. Birds, v. t. 193. *Walc. Syn.* ii. p. 157.

FEM.

Tringa fulicaria. *Lin. Syſt.* ii. p. 249. 20. *Gmel. Syſt.* ii. p. 676. 6.
Phalaropus rufeſcens. *Briſ.* vi. p. 20. 4. *Ib.* 8vo. ii. p. 363.
Le Phalarope rouge. *Buf.* viii. p. 225.
Red Coot-footed Tringa. *Edw.* t. 142.

Size of the preceding ſpecies; length eight inches; bill black, an inch long, ſlender, ſtrait, except at the end, which bends downwards. Top of the head, hind part of the neck, back, and ſcapulars, deep aſh-colour; through the eye, from the baſe of the bill, a duſky ſtreak paſſing backwards; quill-feathers duſky, ſome of the ſecondaries tipped white; wing coverts aſh-colour, the greater darkeſt, and tipped with white; the whole under parts, from chin to vent, white; in ſome the under part and ſides of the neck are ferruginous, and the breaſt cinereous; in others the rump and upper tail coverts are banded duſky and white; tail duſky, daſhed with cinereous; legs and feet black, or lead-colour.

Female. The head, throat, hind-neck, back, ſcapulars, and upper tail coverts, black, margined rufous; over the eye a pale rufous-brown ſtreak; rump white, ſpotted with duſky; under parts from the throat duſky red; wings and tail as in the other ſex.

From the authority of Dr. Latham we give this as the female. It is ſo rare a ſpecies in England that few come under examination in a freſh ſtate. Their manners and habits are probably the ſame as the other ſpecies before mentioned.

Inhabits the north of Europe; ſaid to breed at Hudſon's Bay, and lays four eggs. Sometimes called Small Cloven-footed Gull, Scallop-toed Sandpiper.

PHEASANT. A genus of birds, the characters of which are,

Bill ſhort, ſtrong, convex.

The ſides of the head bare, carunculated.

Legs moſt commonly furniſhed with a ſpur.

PHEASANT-COMMON.

Phaſianus colchicus. *Lin. Syſt.* i. p. 270. 3. *Gmel. Syſt.* i. p. 741. *Raii Syn.* p. 56. A. 1. *Will.* p. 117. t. 28. *Ind. Orn.* ii. p. 629. *Briſ.* i. p. 262. 1.—*Ib.* 8vo. p. 73.

Le Faiſan. *Buf.* ii. p. 328. t. 11.

Common Pheaſant. *Albin.* i. t. 25. 26. *Hayes, Br. Birds*, t. 20. *Will. Angl.* p. 163. t. 28.

Lath.

Lath. Syn. iv. p. 712. 4. *Lewin, Br. Birds*, iii. t. 31. *Walc. Syn.* ii. t. 178. *Don. Br. Birds*, v. t. 101. *Pult. Cat. Dorſet.* p. 7.

Weight of a full-grown bird near three pounds; length three feet; bill pale yellowiſh horn-colour; irides yellow; ſides of the head bare, granulated, crimſon, minutely ſpeckled with black. This part is conſiderably brighter and much dilated in the courting ſeaſon; at which time alſo it erects a tuft of ear-like feathers on each ſide of the head. The various tints of green-gold, blue, and violet, in the plumage, exceeds deſcription, and is too well known to require it. The tail conſiſts of eighteen feathers, very cuneiform, the two middle ones about eighteen or twenty inches long; the legs are furniſhed with a ſpur three quarters of an inch long; toes joined by a ſtrong membrane at the baſe.

The female is not ſo large, of a rufous-brown colour, mixed with grey and duſky; the bill is brown; irides hazel; and the ſides of the head covered with feathers; tail of the ſame ſhape, but not ſo long as in the male.

It is difficult perhaps to trace the origin of this ſpecies, where it came from, or when firſt introduced into this country. It is now found in a ſtate of nature in almoſt every part of the old continent, except the northern, but is not known in America. It bears confinement tolerably well, and produces a great many eggs in that ſtate. The female will

ſometimes incubate if not diſturbed by the male, which is too often the caſe; on which account the eggs are generally put under a common fowl to be hatched: by this means a great many are reared and given their liberty every year by gentlemen of property. Were it not for this probably the breed would be extinct in a few years in ſpite of the ſeverity of the game laws. The demand for them at the tables of the luxurious, and the irreſiſtible mark they offer to the ſportſman, would ſoon cauſe their deſtruction in this age of excellence in the art of ſhooting flying.

It is a fooliſh bird; when rouſed will frequently perch on the firſt tree, and is ſo intent upon the dogs as to ſuffer the ſportſman to approach very near. At the time they perch they moſt frequently crow, or make a chuckling noiſe, by which the unfair ſportſman is led to their deſtruction. The poacher will catch them in nooſes made of wire, horſe-hair twiſted, and even with a briar ſet in the like manner at the verge of a wood, for they always run to feed in the adjacent fields morning and evening. Beſides this they are taken by a wire faſtened to a long pole, and by that means taken off their rooſt at night; or by fixing a bunch of matches lighted at the end of a pole are ſuffocated, and drop off the perch. Foxes alſo deſtroy a great many, in particular the females when ſitting on their neſt.

The Pheaſant is found partially in moſt parts of England, but not ſo plentiful in the north, and

rarely

rarely in Scotland. Wood and corn land ſeem neceſſary to its exiſtence: it is partial to oak and beech woods, on the ſeed of which they feed; buckwheat is alſo a favourite food. In the autumn they frequent turneps. Large wooded tracts only produce them in abundance, and they moſtly frequent the thickeſt and moſt impenetrable coverts, or ſuch as are covered with long graſs, which the female generally makes choice of to depoſit her eggs, ſcraping together a few ſurrounding dry vegetables for a neſt, and laying from eight to twelve eggs. The male is frequently heard to crow in ſpring, clapping his wings at the ſame time.

In confinement the female ſometimes aſſumes the plumage of the male; at this time ſhe becomes barren, and is equally buffetted by both ſexes. This ſtrange change of plumage does not ſeem to be the effect of age, for we have been aſſured by our noble friend Lord Carnarvon, who has had ſeveral in that ſtate, that it takes place at three or four years old. In the one that nobleman favoured us with the colours were not ſo bright as generally found in the other ſex. Whether barrenneſs is the occaſion of this change, or whether the want of commerce with the other ſex, by reaſon of the male plumage, is the occaſion of her not breeding, is yet to be diſcovered by diſſection; for if in the breeding ſeaſon there ſhould appear any eggs in the ovarium, and thoſe diſtended, there can be no doubt of the latter cauſe. In a ſtate of nature this circumſtance probably does not take place.

This bird does not appear to pair; but the female carefully hides her neſt from the male; and we are informed where they are in plenty, and food provided for them, the two ſexes do not in general feed together.

Domeſtication generally produces variety, and we find this bird mixed more or leſs with white; ſometimes wholly ſo.

In the *Naturaliſt's Calendar*, publiſhed in 1795 from the papers of the late Rev. Gilbert White of Selborne, a plate and deſcription is given of a ſingular bird killed in Hampſhire, which he conſidered as an hybrid between the Pheaſant and domeſtic fowl. The head, neck, breaſt, and belly, gloſſy black; the back, wing-feathers, and tail, pale ruſſet, ſtreaked ſomewhat like the upper parts of a Partridge; the tail was even at the end, and not very long; legs bare of feathers, and deſtitute of a ſpur; round the eye the ſkin was bare and ſcarlet; weight three pounds three ounces and a half. By the drawing there alſo appears to be ſome white on the ſhoulder, and ſome dark feathers in the middle of the tail beneath.

It is generally believed the Pheaſant and domeſtic fowl will breed together; if ſo, poſſibly the colour of the ſpurious breed would depend much on that of the common fowl. Some authors have given a deſcription of ſuch, but not at all like Mr. White's bird. One in the Leverian Muſeum is almoſt throughout of a dingy reddiſh brown colour, with very

very few markings. Surely if it was common (as ſome have aſſerted) for male Pheaſants to mix breed with our tame fowls in farm-yards contiguous to their natural haunts, more would be known of the ſpurious breed; at preſent no one has determined whether ſuch a produce are mules or not. A variety called the Ring Pheaſant is not uncommon in this country. *See* that bird.

PHEASANT-RING, or RING-NECKED.

Phaſianus torquatus. *Gmel. Syſt.* ii. p. 742.
——— colchicus. *Ind. Orn.* p. 629. 4. β.
Ring Pheaſant. *Lath. Syn.* iv. p. 715.—*Ib. Sup.* p. 208. *Oſt. Menag.* p. 57. 58. (M. F.)

This ſeems to be a mere variety of the common ſpecies, differing only in having a white ring round the neck; and the plumage in general thought to be more brilliant, and the markings more diſtinct.

It is ſaid to have been firſt introduced by the late Duke of Northumberland, and was called the Barbary Pheaſant. His Grace bred and turned out many at his ſeat at Alnwick in Northumberland. Lord Carnarvon alſo turned out ſeveral at his ſeat at Highclere in Berkſhire; at both which places we have ſeen them, and the mixed produce between that and the common Pheaſant, in which the ring on the neck is very indiſtinct; ſometimes only a few white feathers are to be found. Suppoſed to be hardier and eaſier reared in confinement than the

other. It is ſaid to be found in ſome parts of China, and in Tartary.

The eggs no ways differ from the common ſort, which are of a light brown colour.

PHEASANT-SEA. Vide *Duck Pintail.*

PIANET. Vide *Magpie, Oyſtercatcher.*

PICARANI. Vide *Avoſet.*

PIE. Vide *Oyſtercatcher.*

PIET. Vide *Ouzel-water.*

PIGEON-ROCK. Vide *Dove-ſtock.*

PIGEON-STOCK. Vide *Dove-ſtock.*

PIGEON-WOOD. Vide *Dove-ring.*

PINK. Vide *Finch-chaf.*

PIRENET. Vide *Shieldrake.*

PLOVER. A genus of birds, the characters of which are,

Bill ſtrait, not longer in general than the head.

Noſtrils linear.

Toes three, all placed forwards.

PLOVER-BASTARD. Vide *Lapwing.*

PLOVER-CREAM-COLOURED.

Charadrius gallicus. *Gmel. Syſt.* ii. p. 692.

Curſorius europæus. *Ind. Orn.* ii. p. 751. 1.

Le Coure-vite. *Buf.* viii. p. 128.

Cream-coloured Plover. *Lath. Syn.* v. p. 217. 25. —*Ib. Sup.* p. 254. t. 116. *Lewin, Br. Birds,* v. t. 187. *Walc. Syn.* ii. t. 164.

Length

Length of this ſpecies ten inches. Bill black, three quarters of an inch long, ſlender, and bent at the tip. Plumage in general cream-colour, paleſt beneath; behind the eye a patch of black; through them a pale ſtreak paſſing back to the hind-head, and dividing the black; quills black; tail cream-colour, marked with black near the tip; legs yellowiſh.

This appears to be a rare bird. One was killed in France, as mentioned by Buffon; another was killed near St. Alban's in Kent, at the ſeat of William Hammond, Eſq. who preſented it to Dr. Latham, in whoſe inſtructive muſeum we ſaw it, and who has given a figure of it in the Supplement to his *Synopſis*. In this ſpecimen the markings about the head do not appear to be ſo conſpicuous.

It is ſaid to be a bold bird, and to run with great ſwiftneſs; but its habits ſeem to be much in obſcurity.

The ſhape of the bill in this ſpecies ſeems to divide it from the Plovers, although it has no back-toe like that genus.

Dr. Latham in his *Synopſis* made it a Plover, but in his *Index Ornithologicus* has removed it, and calls it a Curſorius.

PLOVER-GREATER. Vide *Greenſhank.*

PLOVER-GREEN. Vide *Lapwing.*

PLOVER-GREY. Vide *Sandpiper-grey* and *Plover-golden.*

PLOVER-GOLDEN.

Charadrius pluvialis. *Lin. Syſt.* i. p. 254. 7. *Gmel. Syſt.* ii. p. 688. *Ind. Orn.* ii. p. 740. 1. *Briſ.* v. p. 43. 1. t. 4. f. 1.—*Ib.* 8vo. ii. p. 222.

Pluvialis viridis. *Raii Syn.* p. 111. A. 2.—190. 9. *Will.* p. 229. t. 57.

Pluvialis aurea minor. *Briſ.* v. p. 47. 2.—*Ib.* 8vo. ii. p. 223.

Le Pluvier doré. *Buf.* viii. p. 81.

Golden, *or* Green Plover. *Br. Zool.* ii. N° 208. t. 72.—*Ib.* fol. 128. *Arct. Zool.* ii. N° 399. *Will. Angl.* p. 308. *Albin.* i. t. 75. *Lath. Syn.* v. p. 193. 1.—*Sup.* p. 252. *Lewin, Br. Birds*, iii. t. 181. *Wale. Syn.* ii. t. 158. *Don. Br. Birds*, ii. t. 45. *Pult. Cat. Dorſet.* p. 16.

PROVINCIAL.

Grey Plover. Whiſtling Plover.

Weight of this ſpecies between ſeven and eight ounces; length ten inches and a half. Bill one inch, duſky; irides hazel. The general plumage above is duſky, ſpotted with greeniſh yellow, brighter on the back and ſcapulars, paleſt on the wing coverts; ſides of the head and neck, and ſides of the body, lighter; middle of the belly and vent white; quills duſky, ſlightly margined at the tips with grey; tail duſky, ſpotted with yellow and dull white, ſomewhat in form of bars; legs black.

A variety

A variety is ſaid to poſſeſs a ſmall claw behind in lieu of a back-toe.

In the breeding ſeaſon both ſexes appear black on the lower part of the breaſt; theſe feathers begin to ſhew themſelves in March, and are perfected in May, at which time the female begins to lay. It is a common bird, found in moſt parts of the known world. With us it chiefly inhabits open ground, ſuch as heaths, moors, and downs; in ſevere weather the ſea-coaſt; but repairs to the more uncultivated waſtes of the northern mountains to breed. We have ſeen them in various parts of Scotland at that ſeaſon on ſwampy ground upon the higher hills, and even on the lower lands covered with heath, amongſt which they lay their eggs, four in number, about the ſize and ſhape of that of the Lapwing's; colour cinereous-olive, blotched with duſky.

The young run as ſoon as they are excluded from the egg, and follow the old ones to the moiſt places in ſearch of worms. At firſt they are covered with down of a duſky colour, and are incapable of flying for a conſiderable time. The parent birds are very tenacious of their young; become very bold at this time; will light juſt before a dog, and run on the ground to entice him from their neſt.

PLOVER-KENTISH. Vide *Plover-ringed.*

PLOVER-LONG-LEGGED.

Charadrius himantopus. *Lin. Syſt.* i p. 255. 11. *Gmel. Syſt.* ii. p. 690. *Ind. Orn.* ii. p. 741. 3. Himantopus.

Himantopus. *Raii Syn.* p. 106. 9.—190. 7.—*Ib.* 193. 1. t. 1. f. 3. *Will.* p. 219. t. 54. *Brif.* v. p. 33. 1. t. 3. f. 1.—*Ib.* 8vo. ii. p. 220. 1.

L'Echaſſe. *Buf.* viii. p. 114. t. 8.

Long-legged Plover. *Br. Zool.* ii. N° 209.—*Ib.* fol. 128. Addenda. *Arct. Zool.* ii. N° 405. *White, Hiſt. Selb.* t. p. 258. *Lath. Syn.* v. p. 195. 3.—*Ib. Sup.* p. 252. *Lewin, Br. Birds,* iii. t. 182. *Walc. Syn.* ii. t. 159. *Don. Br. Birds,* iii. t. 55.

PROVINCIAL.

Longlegs.

This extraordinary ſpecies is certainly the longeſt-legged bird, in proportion to its bulk, hitherto known. Length, from the apex of the bill to the end of the tail, thirteen inches; from that to the end of the toes five inches more.

Bill two inches and a half long, ſlender, and black; irides red. Forehead, round the eyes, and rump, white; crown of the head, back, and wings, gloſſy black; tail the ſame, inclining to grey; outer feathers white; neck and under parts white; the hind part of the neck marked with duſky ſtreaks; in ſome theſe ſtreaks are wanting, the effect probably of maturity; the thigh is bare of feathers three inches and a half from the knee; legs four inches and a half long, red; the outer and middle toes connected by a membrane at the baſe.

Six

Six of this ſpecies were ſhot out of ſeven in a flock in the month of April at the verge of a lake not very diſtant from Farnham in Surrey. One of them was preſerved by the late Rev. Mr. White of Selborne, and is now in the poſſeſſion of Mr. White in Fleet-ſtreet, where we ſaw it. This bird is wholly white, except the wings and back as far as the rump, which is black. Of this bird there is a good figure in White's *Natural Hiſtory of Selborne*, who ſays it weighed, when drawn and ſtuffed with pepper, four ounces and a quarter.

This is a rare bird in England; but Dr. Latham informs us it is ſufficiently plentiful in the Eaſt and Weſt Indies, in Egypt, on the ſhores of the Caſpian ſea, and in the warmer parts of America. Specimens received from the two firſt places had the crown and all the hind part of the neck black. The above author gives another variety in his *Supplement*,

PLOVER NORFOLK. Vide *Buſtard thick-kneed*.

PLOVER RINGED.

Charadrius Hiaticula. *Lin. Syſt.* i. p. 256. *Gmel. Syſt.* ii. p. 683. *Raii Syn.* p. 121. A. 6. *Will.* p. 190. 13. *Will.* p. 230. t. 57. *Ind. Ornith.* ii. p. 743. 8.

Pluvialis torquata minor. *Briſ.* v. p. 63. 8. t. 5. f. 2. *Ib.* 8vo. i. p. 227.

Petit Pluvier à collier. *Buf.* viii. p. 190. t. 6.

Sea

Sea Lark. *Alm.* i. t. 80. *Will. Angl.* p. 300. t. 57.

Ringed Plover. *Br. Zool.* ii. N° 211.—*Br. fol.* 129. t. Addenda. *Arct. Zool.* ii. N° 400. *Lath. Syn.* v. p. 201. 8. *Lewin, Br. Birds,* v. i. 184. *Walc. Syn.* ii. t. 161. *Don. Br. Birds,* i. t. 18. *Pult. Cat. Dorſet.* p. 16.

PROVINCIAL.

Dulwilly.

This ſpecies weighs about two ounces; length between ſeven and eight inches. Bill half an inch long, the baſe half orange, the other black; irides hazel. At the baſe of the upper mandible the feathers are black, which paſſes in a broad ſtreak under the eye, taking in the coverts of the ears; forehead white; behind which, on the top of the head, is a black band from eye to eye; over the eye a ſtreak of white paſſing backwards; chin and throat white, continuing in a circle round the neck; beneath this, on the lower part of the neck, is a broad black band encircling that part; the back of the head and upper parts of the body and wing coverts pale brown; under parts white; quills duſky, with ſome white at their baſe; ſhafts partly white; tail conſiſts of twelve feathers; the two middle brown, duſky towards the tips; the three next black towards the end; in the next is only a brown band on the inner web; the outer one quite white;

white; the whole tipped the ſame: legs orange; claws black.

When flying this bird ſhews a white bar on the wing, by reaſon of the baſe of ſome of the quills and tips of ſome of the greater coverts being of that colour.

The Ring Plover is a plentiful ſpecies in moſt parts of the known world. In England every part of the coaſt is enlivened with their ſhrill note. It has been ſaid to leave us in the autumn; but this is certainly not the caſe, as we have frequently procured them throughout the ſevereſt winters in Devonſhire, Cornwall, and other places; but at this time they quit the open ſhores, and ſeek ſhelter in creeks and inlets.

Early in May they pair, and we have found their eggs as early as the 20th of that month. It makes no neſt, but lays four eggs in a ſmall cavity in the ſand, juſt above high-water mark. Theſe are of a cinereous-brown, marked all over with ſmall black and aſh-coloured ſpots; weight three drams. It is remarkable that theſe, as well as moſt if not all ſpecies of birds that lay invariably four eggs only on the ground, place them ſo as to occupy the leaſt poſſible ſpace; that is, with their ſmall ends touching each other as a centre. It is greatly attached to its young; will uſe various deceptions to ſave them from men or dogs; ſometimes will flutter along the ground as if crippled, and if purſued will fly to a little diſtance, diſtend

all

all its feathers; and ſeem to tumble head over heels repeatedly, till it has enticed its enemy to a diſtance from its young, and then it flies off.

In the autumn they become gregarious, and continue in ſmall flocks all winter, mixing ſometimes with Purrs and Dunlins. We have frequently obſerved a variety of this bird without any black about the head and breaſt, and the bill and legs duſky; others ſeem inclined to thoſe markings, and a tinge of orange on the bill and legs. From theſe gradations it ſhould appear ſuch are birds of the firſt year not maturely feathered; and are not unfrequently ſhot in company with the others. We ſuſpect the *Alexandrinus* of Linnæus to be this bird in one of its changes.

Lewin has figured a bird under the title of Kentiſh Plover, Vol. V. t. 185, which he conſiders as a diſtinct ſpecies. This, however, is much to be doubted, as the principal difference from the young of this ſpecies appears to be in the colour on the back of the head, which he deſcribes to be of a clay-colour, though in his figure it is of an orange-red.

PLOVER-STONE. Vide *Buſtard-thick-kneed.*

PLOVER-WHISTLING. Vide *Plover-golden.*

POCHARD.

Anas ferina. *Lin. Syſt.* i. p. 203. 31. *Gmel. Syſt.* i. p. 530. *Ind. Orn.* ii. p. 862. 77.

Anas fera fuſca. *Raii Syn.* p. 143. A. 10. *Will.* p. 288. t. 72.—*Ib.* p. 282. 12. Fem.?

Penelope.

Penelope. *Briſ.* vi. p. 384. 19. t. 35. f. 1.—*Ib.* 8vo. ii. p. 462.

Millouin. *Buf.* ix. p. 216.

Poker, Pochard, Red-headed Widgeon. *Br. Zool.* ii. N° 284.—*Ib.* fol. 156. t. Q. 5. *Arct. Zool.* ii. N° 491. *Will. Angl.* p. 367. t. 72. *Albin.* ii. t. 98. *Lath. Syn.* vi. p. 523. t. 68. *Lewin, Br. Birds*, vii. t. 253. *Pult. Cat. Dorſet.* p. 20.

A ſpecies of Duck about the ſize of a Widgeon, but not ſo ſlender and elegantly formed. Weight about twenty-eight ounces; length nineteen inches.

Bill broad, blue, tip black; irides in ſome orange, others hazel; head and neck deep cheſnut; lower part of the neck, breaſt, and upper part of the back, black, with ſmall undulated lines of grey; back and ſcapulars cinereous and duſky, diſpoſed in ſmall undulated lines; ſmaller wing coverts darker; greater coverts and ſecondary quill-feathers blue-grey, primary quills duſky; rump and under tail coverts black; under part of the body duſky white, marked with numerous ſmall duſky lines, darkeſt at the vent; the tail conſiſts of fourteen feathers, duſky, daſhed with aſh-colour; legs lead-colour; feet the ſame, very broad.

The female differs in having the head and neck ferruginous-brown; breaſt and belly duſky white, clouded with brown; under tail coverts duſky and white; in other reſpects like the male, but the markings leſs diſtinct.

This ſpecies is frequently caught in our decoys in winter, but is not known to breed with us. It is commonly called Dunbird, or Red-headed Widgeon.

The male poſſeſſes a *labyrinth*, or enlargement of the *trachea*, near the junction with the lungs. This is well figured by Dr. Latham in the fourth volume of the *Linnean Tranſactions*, amongſt many others. This ſingular formation is peculiar to the male only of ſeveral ſpecies of the Duck tribe: what the uſe of it can be has not yet been diſcovered.

POKER. Vide *Pochard.*

POPE. Vide *Puffin.*

POPPINJAY. Vide *Woodpecker-green.*

PORT-EGMONT-HEN. Vide *Gull-ſkua.*

PTARMIGAN.

Tetrao Lagopus. *Lin. Syſt.* i. p. 274. 4. *Gmel. Syſt.* ii. p. 749. *Raii Syn.* p. 55. 5. *Will.* p. 127. t. 32. *Ind. Orn.* ii. p. 639. 9. *Briſ.* i. p. 216. 12.—*Ib.* 8vo. i. p. 60.

Lagopede. *Buf.* ii. p. 264. t. 9.

White Game. *Will. Angl.* p. 176. t. 32.

Ptarmigan. *Br. Zool.* i. N° 95. t. 43.—*Ib.* fol. 86. t. M. 4. 5. *Arct. Zool.* ii. p. 315. D. *Lath. Syn.* iv. p. 741. 10. *Lewin, Br. Birds*, iv. t. 134. *Walc. Syn.* ii. t. 182. *Don. Br. Birds*, i. t. 12.

This ſpecies of Grous weighs about twenty ounces; length fifteen inches. Bill black; irides hazel.

hazel. The ſummer plumage is a mixture of light brown and aſh-colour, marked with minute bars and ſmall duſky ſpots; the head and neck with broad bars of black, white, and ruſt-colour; belly white; wings white, with black ſhafts to the greater quills. Some are more rufous on the head, ſuppoſed to be the male ſex. In the month of September it begins to change its plumage, and about the middle of October it is of a pure white all over, except the ſhafts of the wings and the tail, which laſt conſiſt of ſixteen feathers; the two middle ones white, the reſt black, with a little white on the tips of the ſecond feathers from the middle. In the male alſo there are black feathers covering the noſtrils, and from thence to the eyes.

This is taken from the Ptarmigan of the Scottiſh highlands; but in thoſe received from Norway all the black feathers of the tail were tipped with white, largely ſo in the middle feathers, but gradually decreaſing till almoſt loſt on the exterior ones. When the tail is cloſed the black is completely concealed by the coverts, which are white, and reach to the end.

It is a very local ſpecies with us, confined to the loftieſt mountains of the north. Some few are yet found to the ſouth of the Tweed, but it is more plentiful on ſome of the highlands of Scotland, from which it rarely or never deſcends, even in the ſevereſt ſeaſon, when nothing but ſnow is to be ſeen.

It makes no neſt, but depoſits ten or twelve eggs on the bare ground, amongſt the rocks. Theſe are of a dirty white, ſpotted and blotched with rufous-brown, ſomething larger than thoſe of the Partridge. It is by no means a ſhy bird, but will ſuffer the ſportſman to approach very near. The herdſmen frequently knock them down with ſticks.

In winter they aſſemble together in flocks. They are called Birch Partridges in Nova Scotia; with us White Game, or White Partridge.

We are inclined to believe the White Partridge of Mr. Edwards is a mere variety of this bird, contrary to the opinion of that great naturaliſt Mr. Pennant, who ſays he has received both from Norway. But with the moſt diligent inquiry we cannot find more than one ſpecies known in that country, where they are called Rype, but pronounced Reuper; and in the many we have ſeen no ſpecific diſtinction could be diſcovered.

PUCKERIDGE. Vide *Goatſucker.*

PUFFIN.

Alca arctica. *Lin. Syſt.* i. p. 211. 4. *Gmel. Syſt.* ii. p. 549. *Ind. Orn.* ii. p. 792. 3.

Anas arctica. *Raii Syn.* p. 120. A. 5. *Will.* p. 244. t. 65.

Fratercula. *Briſ.* vi. p. 81. 1. t. 6. f. 2.—*Ib.* 8vo. ii. p. 380.

Le Macareux. *Buf.* ix. p. 358. t. 26.

Puffin. *Br. Zool.* ii. N°232.—*Ib.* fol. 135. t. H.

Arct.

Arct. Zool. ii. N° 427. *Will. Angl.* p. 325. *Albin.* ii. t. 78, 79. *Edw.* t. 358. f. 1. *Lath. Syn.* v. p. 314. 3. *Lewin, Br. Birds*, vi. t. 225. *Don. Br. Birds*, i. t. 8. *Walc. Syn.* i. t. 87. *Pult. Cat. Dorſet.* p. 17.

The weight of this ſpecies of Auk is between twelve and thirteen ounces; length exceeds twelve inches. The bill is of a very ſingular form; it is about an inch and a half deep at the baſe, much compreſſed ſideways, and arched, but ends in a point, where the upper mandible is a little hooked; at the baſe of this mandible is an elevated rim, full of ſmall punctures of a light colour; next to which is a ſpace of a bluiſh-grey, at the bottom of which the noſtrils are lodged at the edge, which is a long narrow ſlit; from thence it is orange-coloured with four oblique furrows; the under mandible correſponds in colour, but has only three furrows; irides hazel, in ſome greyiſh; orbits red; above the eye is a triangular callous protuberance, beneath an oblong one; the top of the head and whole upper parts are black, paſſing round the neck in a collar; the ſides of the head and all the under parts are pure white; the chin in ſome is grey, in others white, and the cheeks are grey; quills duſky; tail ſhort, and conſiſts of ſixteen feathers; legs and feet orange; claws black, the inner one much hooked.

It is remarked that the bill of this bird varies much according to age; at firſt it has no furrows,

and is of a duſky colour, the yellow colour and furrows increaſing with age.

Theſe birds appear on many parts of our rocky coaſt about the middle of April, and begin to breed about the middle of May. On the ſtupendous cliffs of Dover, and other ſuch places, they depoſit their ſingle egg in the holes and crevices; in other places they burrow like rabbits, if the ſoil is light; but more frequently take poſſeſſion of rabbit burrows, and lay their egg many feet under ground. This is the caſe on Prieſtholm Iſle off the coaſt of Angleſea, and other ſmall iſlands off St. David's, where the ſoil is ſandy. On St. Margaret's Iſland off that place we have ſeen the fiſhermen draw them out of their holes in a ſingular manner: by introducing the hand it is ſeized by the Puffin, who ſuffers itſelf to be drawn out, as it will not quit its hold. In other places they are caught with ferrets, and the young are taken and pickled. About the latter end of Auguſt they retire from our coaſt, and are all completely migrated by the beginning of September, together with the Razorbill and Guillemot.

The egg is white, about the ſize of a Hen's.

At Dover this, as well as the Razorbill, is indiſcriminately called Willock; it is alſo variouſly called Coulternel, Lunda Bouger, Mullet, Gulderhead, Bottlenoſe, Pope, Marrot, and Sea Parrot.

The principal food of theſe birds is ſmall fiſh, particularly ſprats, with which they feed their young.

young. It is not known to what part they go when they retire from us; but they have been found in abundance in various parts of the northern as well as ſouthern hemiſphere.

PUFFIN-MANKS. Vide *Shearwater.*

PUIT. Vide *Gull-black-headed.*

PURRE.

Tringa cinclus. *Lin. Syſt.* i. p. 251. 18. *Gmel. Syſt.* ii. p. 680. *Raii Syn.* p. 100. A. 13—190. 11. *Ind. Orn.* ii. p. 735. 35. *Briſ.* v. p. 211. 10. t. 19. f. 1.—*Ib.* 8vo. ii. p. 267.

L'Alouette de Mer. *Buf.* vii. p. 548.

Sanderling. *Albin.* iii. t. 88.

Purre, *or* Stint. *Br. Zool.* ii. N° 206. t. 71—*Ib.* fol. 126. *Arct. Zool.* ii. N° 390. *Will.* p. 226.—*Ib. Angl.* p. 305. *Lath. Syn.* v. p. 182. 30. *Don. Br. Birds,* t. 32. *Pult. Cat. Dorſet.* p. 15. *Lewin, Br. Birds,* v. t. 174. *Walc. Syn.* ii. t. 150.

Tringa ruficolis. *Gmel. Syſt.* ii. p. 680? *Ind. Orn.* ii. p. 736. 36?

Red-necked Sandpiper. *Lath. Syn.* v. p. 183. 31?

PROVINCIAL.

Ox-bird. Ox-eye. Leaſt Snipe. Wagtail.

This ſpecies of Sandpiper is near eight inches in length; weight about an ounce and three quarters.

Bill black, one inch and a quarter long; irides dark. The head and hind part of the neck pale aſh-coloured brown, ſtreaked with duſky, with a whitiſh line between the bill and eye; back and wing coverts more brown, daſhed with aſh-colour; greater coverts darker, tipped with white; fore part of the neck white, mixed with brown; breaſt and belly white; quills duſky, with more or leſs white at the baſe; the ſhafts of ſome and the ſecondaries tipped with white, and edged with cinereous; the tail conſiſts of twelve feathers; the two middle ones rather longeſt, duſky, edged with aſh-colour; the reſt browniſh aſh-colour, edged with a lighter ſhade; legs duſky. In ſome birds the head is without duſky ſtreaks; others ſtreaked on the back, and rufous-brown on the rump, mixed with duſky. Dr. Latham has given another variety, ſomewhat different in the markings, and leſs in ſize, meaſuring only ſix inches and three quarters in length.

The Purre is found on all our coaſts during winter, but ſeem moſt partial to the flat ſandy ſhores and inlets, where large flocks are ſeen in company with Sanderlings. It is ſuſpected ſome few remain with us all the year, and of courſe breed here; but we do not remember to have ſeen any after the month of May till the middle of Auguſt. In the very extenſive ſandy flats of the coaſt of Caermarthenſhire we have ſeen thouſands of theſe mixed with Sanderlings during the winter; but

they all depart in April, at which time the Dunlin appears there for a ſhort time. Theſe three ſpecies are indiſcriminately called Sanderlings.

PUTTOCK. Vide *Kite* and *Buzzard-common*.

Q.

QUAIL.

Tetrao Coturnix. *Lin. Syſt.* i. p. 278. 20. *Gmel. Syſt.* ii. p. 765. *Raii Syn.* p. 58. A 6. *Will.* p. 121. t. 29. *Briſ.* i. p. 247. 14.—*Ib.* 8vo. i. p. 69.

Perdix Coturnix. *Ind. Orn.* ii. p. 651. 28.

La Caille. *Buf.* ii. p. 449. t. 16.

Quail. *Br. Zool.* i. N° 97.—*Ib.* fol. 87. t. M. 6. *Arct. Zool.* ii. p. 320. B. *Albin.* i. t. 30. *Will. Angl.* p. 169. *Lath. Syn.* iv. p. 779. 24.—*Sup.* p. 222. *Lewin, Br. Birds,* iv. t. 138. *Walc. Syn.* ii. t. 185. *Pult. Cat. Dorſet.* p. 7.

This ſpecies of Partridge is about ſeven inches and a half in length.

Bill duſky; irides hazel; in old male birds yellow. The crown of the head is black, tranſverſely marked with rufous-brown; down the middle is a yellowiſh

yellowiſh white line; above the eye, paſſing backwards, is another line of the ſame colour; on the chin and throat is a black mark which turns upwards to the ears; the reſt of thoſe parts are white; the hind part of the neck, back, ſcapulars, and tail coverts, are rufous-brown; the middle of each feather ſtreaked with yellowiſh white, ſurrounded more or leſs with black; ſides the ſame, but has not ſo much of the white ſtreaks; breaſt light ferruginous-brown; ſhafts white; belly paler; wing coverts pale rufous-brown, ſtreaked like the back, but more minutely; quills duſky, the outer webs more or leſs mottled with yellowiſh white; tail duſky, tipt with white, and conſiſts of twelve ſhort feathers hid by the coverts.

The female differs in having no black chin or throat, but only a duſky mark from the ears paſſing downwards; the breaſt is alſo ſpotted with duſky, and the coverts of the wings croſſed with yellowiſh white bars; in other reſpects the ſexes are alike. The legs of both are of a light yellowiſh brown.

The Quail is found in all parts of the old world, but not in America. It is a bold bird, and is frequently uſed in China for fighting, as we do our Game Cocks. In the flight ſeaſon, when theſe birds migrate to and from the north, they are found in prodigious quantities in all the iſlands of the Archipelago, which contains no leſs than forty-five principal ones. It is ſaid an hundred thouſand has been taken in one day on the weſt coaſt of the kingdom

kingdom of Naples. A ſmall portion only extend their flight to this country. With us they appear the beginning of May in our cultivated champaign counties, but in much leſs quantity than formerly. On their firſt arrival the males are continually making a whiſtling note, three times ſucceſſively repeated; which being imitated by a whiſtle or quail-call, they are eaſily enticed into a net: but in this way rarely any but males are taken. Great numbers were annually ſent to London from France alive before the revolution. They grow fat in confinement, where they ſeem to looſe much of their pugnacious diſpoſition. Like the Partridge it makes little or no neſt; but the female depoſits eight or ten eggs on the bare ground, of a yellowiſh colour, blotched or ſpotted with duſky, but ſubject to great variety as to the ground colour and diſpoſition of the ſpots. Green wheat is the principal place of their reſort with us, where they breed; and in the ſtubble of that grain the ſportſman moſt commonly finds them. In October they leave us, and return ſouth, leaving ſome few (probably of a later brood) behind to brave the ſeverity of our winter. Several inſtances of their having been killed in that ſeaſon have come to our knowledge.

QUEEST. Vide *Dove-ring*.

R.

RAFTER. Vide *Flycatcher-ſpotted.*

RAIL. A genus of birds, the characters of which are,

Bill ſlender, a little compreſſed, and ſlightly incurvated.
Noſtrils ſmall.
Tongue rough at the end.
Body much compreſſed.
Tail very ſhort.

RAIL, or WATER-RAIL.

Rallus aquaticus. *Lin. Syſt.* i. p. 262. 2. *Gmel. Syſt.* ii. p. 718. *Raii Syn.* p. 113. A. 2.—*Ib.* 190. 12. *Will.* p. 234. t. 56. *Ind. Orn.* ii. p. 755. 1. *Briſ.* v. p. 151. 1. t. 12. f. 2.—*Ib.* 8vo. ii. p. 251.

Gallina ſerica Geſneri. *Raii Syn.* p. 114. 4? *Will.* p. 235.

Le Rale d'eau. *Buf.* viii. p. 154. t. 13.

Velvet Runner. *Will. Angl.* p. 313?

Water Rail, Bilcock, Brook-Ouzel. *Br. Zool.* ii. N° 214. t. 75.—*Ib.* fol. 130. t. E. E. *Will. Angl.* p. 314. *Albin.* i. t. 77. *Lath. Syn.* v. p. 227.

RAI

p. 227. 1. *Lewin, Br. Birds,* v. t. 189. *Wale. Syn.* ii. t. 171. *Pult. Cat. Dorſet,* p. 15. *Don. Br. Birds,* v. t. 104.

PROVINCIAL.

Runner. Grey-ſkit. Skiddy-cock.

The length of this bird is about twelve inches; weight four ounces and a half. Bill an inch and three quarters long, of a duſky colour, reddiſh at the baſe of the upper and greater part of the lower mandible; irides red. The whole upper part of the plumage is of an olive-brown, the middle of each feather black; beneath, from chin to vent, deep aſh-colour, mixed with brown on the lower belly and vent; on the ſides, about the thighs, there are ſeveral tranſverſe bars of black and white; quills duſky; tail ſhort and black, covered by the feathers above; under tail coverts white; legs reddiſh brown; toes long and ſlender, divided to their origin. In ſome there are a few feathers at the elbow of the wing on the coverts that are barred black and white, and the tail feathers margined with brown; the middle ones wholly olivaceous brown. Others have the feathers on the fore part of the neck margined with white. But this laſt variety, as mentioned by other authors, we believe is more rare. The only difference between the ſexes is, the bill of the male being longer and more red.

The Rail is by no means a plentiful ſpecies, but

is

is ſometimes found throughout moſt parts of England in low ſituations, about water-courſes and rivulets, where it ſeeks ſhelter amongſt ſedge-ruſhes and reeds, and is ſeldom put to flight except preſſed by dogs, rather depending on its legs for ſafety. It ſwims, and even dives well occaſionally; but it delights more in wet ground, and ſhallow water it can wade through without ſwimming. Its principal food is worms, ſlugs, and inſects. When rouſed it flies only a ſmall diſtance, and that in a heavy and awkward manner, with its legs hanging down. It runs nimbly, and frequently flirts up its tail.

The neſt is rarely found; it is made of ſedge and coarſe graſs, amongſt the thickeſt aquatic plants; frequently in willow-beds. In ſuch a ſituation we found one with ſix eggs of a ſpotleſs white, and very ſmooth, rather larger than that of a Blackbird; the ſhape a ſhort oval, with both ends nearly alike.

This bird continues with us all the year, and by many is erroneouſly believed to be the Land Rail metamorphoſed; which they ſay takes place in the autumn, not knowing perhaps that bird leaves this country at that ſeaſon. The very great difference in the bills might have taught them otherwiſe, without minute inveſtigation; for that part can never change its form by ſeaſon or climate. It is found in moſt parts of Europe; is ſometimes very fat, and as well flavoured as a Land Rail.

RAIL-LAND. Vide *Gallinule-crake.*

RAIN-FOWL. Vide *Woodpecker-green.*

RAVEN.

Corvus Corax. *Lin. Syſt.* i. p. 155. 2. *Gmel. Syſt.* i. p. 364. *Ind. Orn.* i. p. 150. 1.

Corvus. *Raii Syn.* p. 39. A. 1. *Will.* p. 82. t. 8. *Briſ.* ii. p. 8. 1.—*Ib.* 8vo. i. p. 156.

Corbeau. *Buf.* iii. p. 13. t. 2.

Raven. *Br. Zool.* i. p. 218. 74.—*Ib.* fol. p. 75. *Arct. Zool.* ii. N° 134. *Albin.* ii. t. 20. *Lewin, Br. Birds,* i. t. 33. *Lath. Syn.* i. p. 367. 1. —*Sup.* p. 74. *Walc. Syn.* i. t. 32. *Pult. Cat. Dorſet.* p. 4.

Corvus auſtralis. *Gmel. Syſt.* i. p. 365? *Ind. Orn.* i. p. 151. 2?

South-ſea Raven. *Lath. Syn.* i. p. 369. 2?

This is the largeſt ſpecies of the Crow genus: there is no difference in the plumage of the ſexes. The male weighs about two pounds ſeven ounces; the female four or five ounces more; length near two feet. The bill is black, ſtrong, and thick, two inches and three quarters in length; noſtrils covered with briſtles which reach more than half way down the bill; irides duſky. The whole plumage is black, the upper parts gloſſed with blue; the under dull and duſky; tail conſiſts of twelve feathers, ſomewhat rounded; about the throat the feathers are long, looſe, and ſharp-pointed.

The

RAV

The Raven is a very hardy, crafty, and wary bird; is found in all climates.

The Corvus auſtralis of Gmelin, which Captain Cook found in the Friendly Iſles in the South Seas, is probably no other than a variety of this bird.

It is a voracious ſpecies, and ſeems to poſſeſs much of the habits of the Vulture. Like that tribe it is patient in hunger, ſcents carrion afar off, in plenty gluts itſelf, retires to a ſmall diſtance to digeſt, and then returns again. In this reſpect it is uſeful by devouring putrid matter, which would be highly obnoxious. On this account it would be treated with veneration, if its appetite was thus confined; but as it deſtroys young lambs and ſickly ſheep, which it makes a prey of by firſt picking out their eyes, the huſbandman holds it in deteſtation. Young chickens and ducks are no leſs reſpected by this carnivorous bird.

It is eaſily domeſticated, and is very miſchievous; will catch up any thing glittering, and hide it. We have been aſſured, by a gentleman of veracity, that his butler having miſſed a great many ſilver ſpoons and other articles, without being able to detect the thief for ſome time, at laſt obſerved a tame Raven with one in his mouth, and watched him to his hiding-place, where he found more than a dozen.

The Raven generally makes choice of the largeſt trees to build in. The neſt is formed of ſticks, and lined with wool, hair, and various other ſubſtances:

it

it is commonly placed in the fork of the larger branches. It lays five or ſix eggs of a bluiſh green colour, blotched, and ſpotted with brown and aſh-colour, ſomewhat larger than that of a Crow; their weight from ſix to ſeven drams.

Many breed on our rocky coaſts, where they chooſe the moſt inacceſſible places for nidification. At this time they are exceſſively bold, and will not even ſuffer the Falcon to approach their neſt unpuniſhed. The male and female pair for life, and drive their young from their haunt as ſoon as able to provide for themſelves.

It is ſometimes found quite white, or pied.

RAZOR-BILL.

Alca Torda. *Lin. Syſt.* i. p. 210. 1. *Gmel. Syſt.* ii. p. 551. *Ind. Orn.* ii. p. 793. 5. *Briſ.* vi. p. 892. t. 8. f. 1.—*Ib.* 8vo. ii. p. 383.

Alca Hoieri. *Raii Syn.* p. 119. A. 3. *Will.* p. 243. t. 64, 65.

Le Pingoin. *Buf.* ix. p. 390. t. 27.

Razor-bill, Auk, Murre. *Br. Zool.* ii. N° 230. t. 82.—*Ib.* fol. 136. *Edw.* t 358. f. 2. *Lath. Syn.* v. p. 319—*Ib. Sup.* p. 264. *Lewin, Br. Birds,* vi. t. 224. *Don. Br. Birds,* iii. t. 64. *Walc. Syn.* i. t. 84. *Pult. Cat. Dorſet.* p. 17.

This ſpecies of Auk weighs about twenty-ſeven ounces; length eighteen inches. Bill two inches long from the corner of the mouth, much com-

preſſed ſideways; three quarters of an inch deep at the largeſt part, much arched, and hooked at the end of the upper mandible; it is furniſhed with three tranſverſe furrows; the middle one is white, and croſſes the whole bill; irides light hazel.

The head, chin, throat, and whole upper parts, are duſky black, with a ſtrong tinge of a ruſty hue on the ſides of the head and throat; the quill-feathers are alſo daſhed with cinereous on the outer webs; the greater coverts and ſecondaries tipped with white, forming a narrow band acroſs the wing; from the ridge of the upper mandible is a narrow line of white feathers, which continue to the eye on each ſide; the under parts from the breaſt are white; the tail is cuneiform, conſiſting of twelve pointed feathers, the two middle of which exceed the reſt conſiderably in length; legs duſky black.

We have, in the former part of this work, expreſſed an opinion that this and the Black-billed Auk are diſtinct ſpecies, contrary to the opinion of ſo able an ornithologiſt therein mentioned; for which we have given reaſons.

The gradation in plumage and in the marks on the bill that author mentions in his *Synopſis*, which he conſiders as the different ſtages of the young of this bird before it arrives at maturity, would have ſhaken our opinion; but we have recently procured the young in the month of July, before they could fly, which differed in nothing from the old birds except in wanting the furrows on the bill.

From

From this circumſtance we can have little doubt but that they are diſtinct ſpecies. See *Auk Black-billed.*

The Razor-bill is not ſeen with us in winter, but repairs to our rocky coaſts in the ſpring, and begins to depoſit its ſingle egg the beginning of May (which is of a dirty white, blotched and ſpotted with brown and duſky), on the projecting ſhelves of the higheſt cliffs, where, in ſome ſituations, they may be ſeen hundreds in a row, but not, as we have been informed, cemented to the ſurface; for we have frequently taken them up, and laid them again in the ſame ſpot. A violent gale of wind ſometimes ſweeps away whole ranks. The egg is of a prodigious ſize in proportion to the bird, being about the ſize of that of a Turkey, of a longer ſhape. The principal food of this bird is ſmall fiſh, particularly ſprats, with which it feeds its young, taking three or four at a time in its bill, with the tails hanging out. The eggs of this and the Fooliſh Guillemot are an article of trade in ſeveral of the iſles off the coaſt of Scotland, and are uſed for refining ſugar. They are alſo eaten by the natives there, as well as in other parts; to procure which they are ſuſpended by a rope, and let down from the top of the cliffs; others climb up and down by the help of a crook faſtened to a pole: many periſh in this dangerous employment. This bird is not ſo plentiful as the Guillemot; and it is remarkable that although they

breed in the ſame cliffs they rarely are found to lay on the ſame ledges or ſhelves of the rock, but keep their breeding-places diſtinct, even where they ſwarm like bees about a hive.

REDBREAST.

Motacilla rubecula. *Lin. Syſt.* i. p. 337. 45. *Gmel. Syſt.* ii. p. 993. *Raii Syn.* p. 78 A. 3. *Will.* p. 160. t. 39. *Ind. Orn.* ii. p. 520. 42. *Briſ.* iii. p. 418. 21.—*Ib.* 8vo. i. p. 427.

Rouge-gorge. *Buf.* v. p. 196. t. 11.

Redbreaſt. *Br. Zool.* N° 147.—*Ib.* fol. 100. t. S. 2. *Artt. Zool.* ii. p. 417. D. *Albin.* i. t. 51.—*Ib. Song Birds*, t. p. 55. *Will. Angl.* p. 219. t. 39. *Hayes, Br. Birds*, t. 40. *Lath. Syn.* iv. p. 442. 38. *Lewin, Br. Birds*, iii. t. 107. *Walc. Syn.* ii. t. 238. *Don. Br. Birds*, v. t. 123. *Pult. Cat. Dorſet.* p. 9.

This well-known ſpecies of Warbler needs little deſcription. The upper parts are of a yellowiſh brown, tinged with aſh-colour; forehead, and from chin to breaſt, of a deep rufous-orange; belly and vent whitiſh. The plumage of both ſexes are alike. The neſtling-feathers of young birds are ſpotted; and they do not poſſeſs the red on the breaſt for two or three months after they leave the neſt.

It is ſaid to be a migrative ſpecies, but from no other reaſon than their more frequent and numerous appearance about our habitations in the winter, when

when the woods and fields are deſtitute of inſects; it is then they aſk the protection of man, and are ſo tame as to enter doors and windows, and pick up the crumbs fallen from the table: here they too frequently fall a ſacrifice to the watchful cat. It ſings throughout the winter, except in ſevere weather. About the beginning of April it prepares a neſt in ſome moſſy bank or outbuilding, compoſed of dead leaves, green moſs, and ſtalks of plants, lined with hair, and lays from five to ſeven whitiſh eggs, ſpotted with ruſt-colour and cinereous; their weight about twenty-ſix grains.

It is a conſtant inhabitant of the greater part of the European continent. About Bornholm it is called Tommi-liden, in Norway Peter Ronſmad, in Germany Thomas Gierdet, with us Robin-red-breaſt and Ruddock.

RED-GAME. Vide *Grous-red.*

RED-HOOP. Vide *Finch-bul.*

REDPOLE-GREATER.

Fringilla cannabina. *Lin. Syſt.* i. p. 322. 28. *Gmel. Syſt.* ii. p. 916. *Ind. Orn.* i. p. 458. 82.

Linaria rubra major. *Brif.* iii. p. 135. 30.—*Ib.* 8vo. i. p. 343. *Raii Syn.* p. 91. A. 2. *Will.* p. 191. t. 46.

La grande Linotte des vignes. *Buf.* iv. p. 58.

Greater red-headed Linnet, *or* Redpole. *Br. Zool.* i. N° 132. t. 54.—*Ib.* fol. 110. *Arct. Zool.* ii. N° 261. *Will. Angl.* p. 260. *Albin.* iii.

t. 72. 73.—*Ib. Song Birds*, t. p. 60. *Lewin, Br. Birds*, ii. t. 84. *Lath. Syn.* iii. p. 304. —*Ib. Sup.* p. 167. *Wale. Syn.* ii. t. 222. *Pult. Cat. Dorſet.* p. 12.

This ſpecies of Finch is ſubject to much variety with reſpect to the red markings which, at certain ages and ſeaſons, are found upon the head and breaſt, which has occaſioned it to be multiplied into two diſtinct ſpecies by various ornithologiſts; all of which ſeem to agree that the general colour of both are alike, but that the Redpole has none of that colour upon the breaſt.

It is probable, however, that the full plumage of this bird does not take place till the ſecond or third year, for we have ſeen them in all gradations in the breeding ſeaſon; ſome of which had ſcarcely any tinge on the head or breaſt, and yet by diſſection have proved males. The young, for ſome time after they leave their neſt, reſemble the female, and if taken into confinement in that ſtate, rarely, if ever, throw out the red ſpots, or become ſo ruſous upon the back; and even thoſe which are taken in full maturity moſt frequently loſe all the red feathers in the firſt moulting, which never return. In theſe different ſtages they are commonly known by the name of Brown Linnet, and Red Linnet

The male in full plumage has the bill bluiſh; irides hazel. The head light brown; the feathers

on

on the crown darkeſt in their middle; ſides of the neck inclining to aſh-colour; the forehead roſy red; the back, ſcapulars, and coverts of the wings, fine deep rufous-brown, lighteſt on the rump, and paleſt on the margin of each feather; the breaſt is brown, with more or leſs ſpots like that on the head; belly light rufous-brown; vent almoſt white; quill-feathers duſky black, with more or leſs white on the exterior and interior webs, which forms a conſpicuous bar of that colour on the wing; the tail is forked, the feathers, like thoſe of the quills, black, margined with white, which colour predominates on the inner webs; coverts of the tail black, edged with grey; legs brown. The weight of the male about five drams, that of the other ſex rather leſs. The plumage of the female is more duſky brown; the coverts of the wings rufous-brown; ſides of the throat plain dirty white, the middle part ſtreaked; breaſt and ſides pale brown, with duſky ſtreaks; quills and tail like the other ſex, but the former not ſo deeply margined with white, and of courſe no perceptible bar on the wing.

Theſe birds fly in flocks during winter, at which time the males have little or none of the red markings which in the return of ſpring they put forth. Furzy commons ſeem to be their chief reſort in the breeding ſeaſon, in which they well conceal their neſts; ſometimes a quickſet-hedge or a gooſebery-buſh anſwers this purpoſe.

The neſt is compoſed of moſs and bents inter-

woven with wool, and lined with wool and hair. The eggs are four or five in number, of a bluiſh white, with more or leſs purpliſh ſpecks and ſhort lines; their weight from twenty-four to thirty grains. In the month of April they repair in numbers to the abovementioned places to breed. The firſt broods are hatched in May, but if the neſt is deſtroyed will make others as late as the month of Auguſt. The Redpole ſings very prettily from the time it reſorts to the place of nidification till its young are hatched. At this ſeaſon we have caught a great many of both ſexes on their neſts, and have found the males with little tinge of red on the breaſt only; and from that every intermediate ſtage to the full red on the breaſt and head.

The quotations of authors who deſcribe the immature bird of this ſpecies will be found under the head of *Linnet-brown*.

REDPOLE-LESSER.

Fringilla linaria. *Lin. Syſt.* i. p. 322. 29. *Gmel. Syſt.* i. p. 917: *Ind. Orn.* i. p. 458. 83.

Linaria rubra minor. *Raii Syn.* p. 91. A. 3. *Will.* p. 191. t. 46. *Briſ.* iii. p. 138. 31.—*Ib.* 8vo. i. p. 344.

Le Sizerin petite Linotte des vignes. *Buf.* iv. p. 216.

Leſſer Red-headed Linnet, *or* Redpole. *Br. Zool.* N° 132. t. 54.—*Ib.* fol. 111. *Arct. Zool.* ii. N° 262. *Will. Angl.* p. 260. t. 46. *Albin.* iii. t. 75.

t. 75. *Lewin, Br. Birds*, ii. t. 85. *Lath. Syn.* iii. p. 305. 75.—*Sup.* p. 167. *Walc. Syn.* ii. t. 223. *Don. Br. Birds*, v. t. 114. *Pult. Cat. Dorſet.* p. 12.

A ſpecies of Finch much inferior in ſize to the laſt; weight about two drams and a half; length five inches.

The bill is of a light colour, inclining to dull yellow; irides hazel. The forehead is of a purpliſh red; the feathers of all the other parts above are duſky, margined with rufous-brown; chin black; throat and breaſt pink; ſides ſtreaked with duſky; belly white; quills and tail duſky, edged with pale brown; in ſome the rump is tinged with pink; legs duſky.

The female differs in being ſomewhat lighter above, and in the colour on the head, which is not ſo bright, ſometimes yellowiſh. This ſex has the black ſpot on the chin, but none of the pink on the breaſt and throat.

Young birds are deſtitute of the pink feathers; and indeed the males are ſubject to as much variety as the Greater Redpole, and ſometimes have no red at all on the breaſt.

This bird is not uncommon in the ſouthern counties of England during the winter months; at this ſeaſon is gregarious; and numbers frequently taken about London and other parts by birdcatchers: it is there called Stone Redpole. It is ſaid to breed

brown, ſpotted and blotched with black, moſt numerous at the larger end; in ſhape and appearance much like that of the Lapwing, but rather ſmaller.

We have ſeen theſe birds on Romney Marſh in the breeding ſeaſon; and when diſturbed from their neſt they fly round like the Lapwing.

The young are only obſcurely marked with white ſpots.

REDSHANK-CAMBRIDGE. Vide *Godwit-Cambridge.*

REDSHANK-SPOTTED. Vide *Snipe-ſpotted.*

REDSTART.

Motacilla Phœnicurus, *Lin. Syſt.* i. p. 335. 34. *Gmel. Syſt.* ii. p. 987.

Sylvia Phœnicurus. *Ind. Orn.* ii. p. 511. 15.

Ruticilla. *Raii Syn.* p. 78. A. 5. *Will.* p. 159. t. 39. *Briſ.* iii. p. 403. 15.—*Ib.* 8vo. i. p. 422.

Roſſignol de muraille. *Buf.* v. p. 170. t. 6. f. 2.

Redſtart. *Br. Zool.* i. N° 146.—*Ib.* fol. 99. t. S. f. 6. 7. *Arct. Zool.* ii. p. 416. B. *Will. Angl.* p. 218. *Albin.* i. t. 50.—*Ib. Song Birds,* t. p. 62. *Hayes, Br. Birds,* t. 40. *Lath. Syn.* iv. p. 421. 11. *Lewin, Br. Birds,* iii. t. 108. *Walc. Syn.* ii. t. 233. *Pult. Cat. Dorſet.* p. 8, *Don. Br. Birds,* iv. t. 82.

PROVINCIAL.

Redtail. Brantail.

This

This ſpecies of Warbler weighs about three drams and three quarters; length five inches and a half.

Bill black; irides hazel. Forehead white; the crown of the head, hind part of the neck, and back, deep bluiſh grey; cheeks and throat black; breaſt, ſides, and rump, ruſty red; tail red, except the two middle feathers, which, as well as the wings, are brown, legs black.

The Redſtart comes to us early in April, and leaves us the latter end of September. It is not uncommon in many parts of England; makes its neſt in a hole of a wall or of a tree, which is compoſed of moſs, and lined with hair and feathers; lays five or ſix eggs of a fine blue colour, rather leſs than thoſe of the Hedge Sparrow, and of a lighter ſhade. The young are at firſt ſpeckled all over, not very unlike the young of the Redbreaſt.

The female is of a light brown, with a daſh of grey on the head and back; the chin whitiſh; breaſt and ſides inclining to rufous; rump and tail like the male, but leſs bright. It is remarkable that many of the Warblers who migrate from the ſouth to breed in our climate confine themſelves to intermediate ſituations, like the Nightingale, which has never been found further north than Yorkſhire, nor further weſt than Somerſetſhire; ſo the Redſtart is rarely found in Cornwall, nor perhaps not frequently weſt of Exeter in Devonſhire.

Its ſong is ſoft and ſhort, and when perched frequently

quently vibrates its tail in a quick and ſingular manner.

REDTAIL. Vide *Redſtart.*

REDWING.

Turdus iliacus. *Lin. Syſt.* i. p. 229. 3. *Gmel. Syſt.* ii. p. 808. *Raii Syn.* p. 64. A. 4. *Will.* p. 159. *Ind. Orn.* i. p. 329. 7. *Briſ.* ii. p. 208. 3. t. 20. f. 1.—*Ib* 8vo. i. p. 216.

Le Mauvis. *Buf.* iii. p. 309.

Redwing, Swinepipe, *or* Wind-Thruſh. *Br. Zool.* N° 108.—*Ib.* fol. 91. t. P. f. 2. *Arct. Zool.* ii. 342. D. *Albin.* i. t. 35. *Lewin, Br. Birds,* ii. t. 59. *Lath. Syn.* iii. p. 22. 7. *Walc. Syn.* ii. t. 199. *Pult. Cat. Dorſet.* p. 10.

This ſpecies of Thruſh is in weight near two ounces and a half; length eight inches and a half; irides duſky. Bill duſky, yellowiſh at the baſe of the upper mandible. The whole upper parts are brown, lighter on the edges of the quill-feathers and coverts; over the eye is a whitiſh ſtreak; breaſt and ſides marked with duſky lines; body under the wings, and under wing coverts, reddiſh orange; the middle of the belly white; legs pale brown.

This bird much reſembles the Throſtle, or common Thruſh, but is rather leſs; and in that bird the ſpots on the breaſt are more diſtinct, the colour under the wings not ſo deep, and it wants the white over the eye.

The

The Redwing is a migrative ſpecies, coming to us in great flocks about the latter end of September, frequently in company with Fieldfares. It is found in greateſt abundance where white-thorn abounds, on the berries of which it feeds. When the weather is ſevere their flight is continued ſouth.

In the hard winter of 1799 vaſt abundance reſorted to the weſt of England, where a ſudden fall of ſnow, unuſually deep in that part, cut them off from all ſupply of food; and being too weak to attempt a paſſage over ſea to a warmer climate, thouſands of theſe, and their companions the Fieldfares, were ſtarved to death.

It is ſaid to breed in Norway and Sweden, and in that ſeaſon to ſing not inferior to our Throſtle; makes a neſt in ſome low buſh in the maple foreſts of the latter country; and lays ſix blue-green eggs, ſpotted with black.

One in my collection of a cream-coloured brown, with all the markings of a pale colour; bill and legs almoſt white.

REEVE. Vide *Ruff.*

RICHEL-BIRD. Vide *Tern-leſſer.*

RINGTAIL. Vide *Henharrier.*

ROBIN. Vide *Redbreaſt.*

ROCKER. Vide *Dove-rock.*

ROOK.

Corvus frugilegus. *Lin. Syſt.* i. p. 156. 4. *Gmel. Syſt.* i. p. 366. *Ind. Orn.* i. p. 152. 5.

Cornix

Cornix nigra frugilega. *Raii Syn.* p. 83. A. 3. *Will.* p. 84. t. 18.

Cornix frugilega. *Brif.* ii. p. 16. 3.—*Ib.* 8vo. i. p. 158.

Le Freux, *ou* la Frayonne: *Buf.* iii. p. 55.

Rook. *Br. Zool.* ii. N° 76.—*Ib.* fol. p. 76. *Arct. Zool.* ii. p. 250. A. *Albin.* ii. t. 22. *Will. Angl.* p. 123. *Lewin, Br. Birds,* i. t. 35. *Lath. Syn.* i. p. 372. 4.—*Ib. Sup.* p. 76. *Wale. Syn.* i. t. 34. *Pult. Cat. Dorſet.* p. 4.

This well-known ſpecies of Crow is about the ſize of the Common or Carrion Crow, and of the ſame black colour; the upper parts, like that bird, gloſſed with purple. The only mark of diſtinction in mature birds is, that this is bare of feathers about the baſe of the bill, which is whitiſh and ſcurfy. But as this is acquired by the habits of this ſpecies thruſting its bill into the ground after worms and various inſects; ſo the young of theſe two birds are not to be diſcriminated except by their note; that of the Rook is not ſo deep and hoarſe as the Crow. In their habits there is an eſſential difference: this is content with feeding on the inſect tribe, particularly what is commonly called the grub-worm, which is the *larvæ* of the Chaffer. But in rendering the huſbandman this piece of ſervice it pays itſelf by taking ſome of his corn alſo. It is gregarious at all ſeaſons, reſorting conſtantly to the ſame trees every ſpring to breed, where their neſts.

may

may be ſeen crowded one over another upon the upper branches. It lays four or five eggs, much like that of the Crow, of a greeniſh colour, ſpotted and blotched with duſky. After their young have taken wing, they all forſake their neſt-trees, but return to them again in October to rooſt; but as winter comes on they ſeek the more ſheltered places at night, ſome neighbouring wood; but generally aſſemble firſt in the uſual place, and then fly off together.

The Rook is partial to cultivated parts, as well as to the habitation of man. It has been ſaid the bill of the Rook is leſs arched than that of the Crow, and that the tail feathers are more rounded; but theſe have never appeared to us materially diſtinct.

ROLLER. A genus of birds, the characters of which are,

Bill ſtrait, bending towards the tip, edges cultrated.

Noſtrils narrow and naked.

Legs for the moſt part ſhort.

Toes, three before and one behind, divided to their origin.

ROLLER-GARRULOUS.

Coracias garrula. *Lin. Syſt.* i. p. 159. 1. *Gmel. Syſt.* i. p. 378. *Ind. Orn.* i. p. 168. 1.

Galgalus. *Briſ.* ii. p. 64. 1. t. 5. f. 2.—*Ib.* 8vo. i. p. 173.

Cornix cœrulia Geſneri. *Raii Syn.* p. 42. *Will.* p. 85.

Pica marina. *Raii Syn.* p. 41. *Will.* p. 89.

Garrulus argentoratenſis. *Raii Syn.* p. 41. 3. *Will.* p. 89. t. 20.

Rollier. *Buf.* iii. p. 135. t. 70.

Roller. *Br. Zool. App.* t. 2. *Arct. Zool.* ii. p. 235. G. *Will. Angl.* p. 131. t. 20. *Edw.* t. 109. *Lewin, Br. Birds,* ii. t. 42. *Lath. Syn.* i. p. 406. 1.—*Sup.* p. 85. *Walc. Syn.* i. t. 41. *Don. Br. Birds,* t. 33.

This is the only ſpecies that has ever been met with in England, and that rarely, probably an accidental wanderer from Germany, where they are not uncommon.

It is deſcribed by Mr. Pennant to be the ſize of a Jay; length twelve inches and a half. Bill black, ſtrait, hooked at the point; baſe beſet with briſtles, but do not cover the noſtrils; ſpace about the eyes ſomewhat bare. The head, neck, breaſt, and belly, are of a light bluiſh green; back and ſcapulars reddiſh brown; coverts on the ridge of the wing rich blue; beneath them pale green; upper part and tips of the quills duſky; the lower parts of a fine deep blue; rump of this laſt colour; tail forked, of a light blue; the outer feather ſtriped with black above, and beneath with deep blue, as is the caſe with ſuch part of the quill-feathers as are black above; the other tail-feathers are dull green; legs ſhort, and of a dirty yellow.

There

There are very few inſtances of this bird being found in England, one of which was killed in Cornwall. It is ſaid to be plentiful in Germany, Sicily, and Malta, where they are ſold in the markets and poulterers ſhops; found alſo in Sweden and Denmark. It haunts woods in the breeding ſeaſon, and makes a neſt in a hole in the ground; at other times they congregate with Rooks and other birds in tilled grounds in ſearch of food.

It is remarkably clamorous.

RUDDOCK. Vide *Redbreaſt.*

RUFF.

Tringa pugnax. *Lin. Syſt.* i. p. 147. 1. *Gmel. Syſt.* ii. p. 669. *Ind. Orn.* ii. p. 725. 1. *Raii Syn.* p. 107. A. 3. *Will.* p. 224. t. 56. *Brif.* v. p. 240. 18. t. 22. f. 1. 2.—*Ib.* 8vo. ii. p. 273.

Combattant, *ou* Paon de Mer. *Buf.* vii. p. 521. t. 29. 30.

Ruff *and* Reeve. *Br. Zool.* ii. N° 192. t. 69.—*—Ib.* fol. 123. t. E. *Arct. Zool.* ii. p. 479. A. *Will. Angl.* p. 302. t. 56. *Albin.* i. t. 72. 73. *Lath. Syn.* v. p. 159. 1. *Lewin, Br. Birds,* v. t. 106. *Walc. Syn.* ii. t. 144. *Don. Br. Birds,* t. 19.

This ſingular ſpecies of Sandpiper is in length about one foot. Bill yellowiſh, in ſome black; irides hazel. The face is covered with yellow pimples; on the back part of the neck the feathers

are very long, and ſtand out on each ſide in a remarkable manner. The colour of this, as well as of other parts, vary ſo much that ſcarce two birds are alike; in general the ruff is barred with black; in others white, or plain brown; the upper parts in general are brown, more or leſs barred with undulated lines of black; lower belly, vent, and upper tail coverts, white; the breaſt moſt times partakes of the ſame colour as the ruff; quills duſky; the four middle tail feathers are barred with black, the reſt plain cinereous-brown; legs yellow.

This is the ſummer dreſs, for the ruff is thrown out in the ſpring as well as the carunculated appearance on the face, both of which are loſt in the autumn. In young birds of the firſt year theſe are wanting, at which time they are called Stags; and, as Mr. Pennant obſerves, might be miſtaken for a different ſpecies. But that author remarks, that the coverts of the wings, which are brown inclining to aſh-colour, and colour of the tail, are invariable marks of diſtinction.

The female, or Reeve, is leſs than the male: the upper parts are brown; the middle of each feather duſky; the edges pale; the greater quills duſky; ſecondaries barred rufous-brown and black; belly, vent, and upper tail coverts, white; tail duſky; legs yellowiſh.

This ſpecies is very local with us; it ſeems to be confined to the fens of Lincolnſhire, Cambridgeſhire, the Eaſt Riding of Yorkſhire, and the Iſle of Ely.

Ely. On theſe marſhy ſituations they arrive early in ſpring, where they breed, and depart the latter end of September.

The males appear firſt, and reſort to one particular ſpot, which is trod bare by their traverſing. On theſe ſpots, well known by the fowler, great numbers are taken, chiefly males, which are fattened for the table. On the appearance of a female the other ſex begin fighting, at which time a net, which was previouſly laid, is drawn over them, and many are taken at one hawl.

The Reeve lays four white eggs, marked with large ferruginous ſpots, which are depoſited in a tuft of graſs.

RUNNER. Vide *Rail-water*.

S.

SANDERLING.

Charadrius calidris. *Lin. Syſt.* i. p. 255. 9. *Gmel. Syſt.* ii. p. 689. *Ind. Orn.* ii. p. 741. 4.

Tringa arenaria. *Lin. Syſt.* i. p. 251. 16. *Raii Syn.* p. 109. A. 11. *Will.* p. 125. *Gmel. Syſt.* ii. p. 680.

Calidris griſea minor. *Briſ.* v. p. 236. 17. t. 20. f. 2.—*Ib.* 8vo. ii. p. 272.

Le Sanderling. *Buf.* vii. p. 532.

Sanderling, *or* Curwillet. *Br. Zool.* ii. N° 212. t. 73.—*Ib.* fol. 129. *Artt. Zool.* ii. N° 403. *Will. Angl.* p. 303. *Albin.* ii. t. 74. *Lewin, Br. Birds,* v. t. 183. *Lath. Syn.* v. p. 197. —*Sup.* p. 253. *Walc. Syn.* ii. t. 160. *Pult. Cat. Dorſet.* p. 16.

A ſmall ſpecies of Plover; weight about two ounces; length eight inches. Bill black, one inch in length; irides duſky. Fore part of the head and ſides beneath the eyes, as well as the whole under parts from chin to vent, white; crown of the head, back of the neck, and ſcapulars, grey, with ſlight duſky ſtreaks down the ſhafts of each feather; wing coverts the ſame colour, but nearly plain; greater quills duſky; ſecondaries grey, tipped with white; tail grey, the exterior feathers lighteſt; legs black. This is the general winter plumage.

Other ſpecimens in my collection, which were killed on the coaſt of Cornwall in the latter end of July and in Auguſt; one has the head, neck, and ſides of the breaſt, ſtreaked with black, and tinged with ferruginous; back and ſcapulars marked with large ſpots of black, and ſome of the feathers edged with light ferruginous. In another the head and neck are dark aſh-colour; back and ſcapulars grey, diſtinctly and prettily marked with large angular ſpots of black. Theſe ſeem to be young birds of that ſeaſon in their gradation of plumage.

In

In ſome the wing coverts are dark brown; others aſh-colour, with more or leſs duſky ſtreaks; and all the inner webs of the quills, and part of the outer webs of the ſecondaries, white; and the middle feathers of the tail duſky.

Whether this bird breeds with us is not yet thoroughly known. We have ſeen them on many parts of the coaſt in the month of April, and in July, but never in the intermediate months. Mr. Boys of Sandwich informs us he thinks they breed on that coaſt, as well as the Purre; and we received ſeveral eggs from that gentleman found on that ſandy ſhore, which are unknown to us, and may poſſibly belong to one of theſe two birds. They are not much unlike that of the Black Tern, but ſmaller.

The Sanderling is found on many of our ſhores, where it flocks together with the Purres, but is not ſo plentiful a ſpecies; and both are indiſcriminately called Oxbirds by ſome perſons. This is alſo called Curwillet and Towwilly.

SANDPIPER. A genus of birds of the order Grallæ, the characters of which are,

Bill ſtrait, ſlender, not exceeding an inch and a half in length.

Noſtrils ſmall.

Tongue ſlender.

Toes divided, or very ſlightly connected at the baſe by a ſmall membrane; hind toe ſmall.

There is a very great degree of connection between this genus and that of the Snipe and Plover. In the former the length of the bill in that genus, and in the latter the want of a back-toe, are the only characteriſtic marks of diſtinction; and theſe in ſome ſpecies run ſo nearly into one another as ſcarcely to be divided. In the young of the Red-ſhank the bill ſcarcely exceeds an inch and a half long; and in the Grey Sandpiper there is no back-toe, but only a ſort of ſpur, very ſmall.

SANDPIPER-ABERDEEN. Vide *Sandpiper-red.*

SANDPIPER-ASH-COLOURED.

Tringa cinerea. *Gmel. Syſt.* ii. p. 673. *Ind. Orn.* ii. p. 733. 25. *Br. Zool.* ii. N° 194.—*Ib.* fol. 124. t. E. f. 1. *Arct. Zool.* ii. N° 236. *Lath. Syn.* v. p. 177. 22. *Lewin, Br. Birds,* v. t. 171. *Walc. Syn.* ii. t. 147.

The length of this ſpecies is about ten inches; weight from four ounces and a quarter to five and three quarters. The bill rather exceeds an inch and a quarter in length, ſwells a little towards the point, ſtrait and black; irides duſky. The top of the head, back of the neck, back, and ſcapulars, aſh-coloured; the firſt a little ſpotted or ſtreaked with duſky; the feathers of the two latter bordered with dirty white, and the ſhafts duſky; rump and upper tail coverts white, with undulated bars of black; the three firſt quill-feathers duſky, the

outer

outer webs of the other primores more or leſs edged with white; ſecondaries dark aſh-colour, bordered with white; the ſhafts of all white; greater wing coverts duſky, more or leſs margined with white; leſſer coverts aſh-colour, with white margins; the throat, lower part of the breaſt, and belly, white; fore part of the neck, upper breaſt, and ſides, ſtreaked and waved with duſky; under tail coverts white, with a few ſpots; tail aſh-colour, edged with white, and ſhafts white; legs duſky; toes bordered with a narrow finely-ſcallopped membrane.

We have ſeen theſe birds in prodigious flocks on the ſouth coaſt of Wales; and Mr. Pennant obſerves the ſame on the ſhores of Flintſhire in the winter.

They depart from us the latter end of March or beginning of April.

Like the moſt of this tribe it is ſubject to much variety. A ſpecimen we received from Penzance in Cornwall the 19th of Auguſt had all the feathers of the back, ſcapulars, and wing coverts, marked with ſemicircular black and whitiſh lines: this we take to be a young bird of that ſeaſon.

We have long entertained doubts reſpecting the diſtinction between this and the Knot. The plumage of that bird, as deſcribed by ſome authors, ſcarcely differs from this in the ſeveral varieties of its change. Mr. Pennant, however, remarks, that the Knot, as a ſpecial mark of diſtinction, has

its

its toes divided to the very bottom; but in this alſo there is ſcarcely any appearance of a membrane that connects the toes together.

Notwithſtanding theſe doubts in our own mind, we have given the Knot, as deſcribed by others, a diſtinct place in this work.

On the coaſt of Caermarthenſhire this is known by the name of Howſter.

Is ſuppoſed to breed in Denmark.

It alſo inhabits America; continues the whole ſummer at Hudſon's Bay, where it breeds, and is called Saſqua piſqua niſhiſh.

SANDPIPER-BLACK.

Tringa Lincolnienſis. *Ind. Orn.* ii. p. 734. 27.
Black Sandpiper. *Br. Zool.* ii. N° 197. *Lath. Syn. Sup.* p. 251.

This ſpecies was ſhot in Lincolnſhire, and communicated to the world by Mr. Pennant, whoſe deſcription we take the liberty to copy. "It was the ſize of a Thruſh; the noſtrils black; the irides yellow; the beak ſhort, blunt at the point, and duſky; the head ſmall, and flatted at top; the colour white, moſt elegantly ſpotted with grey; the neck, ſhoulders, and back, mottled in the ſame manner, but darker, being tinged with brown; in ſome lights theſe parts appear of a perfect black, and gloſſy; the wings were long; the quill-feathers black, croſſed near their baſe with a white line; the

the throat, breaſt, and belly, white, with faint brown and black ſpots of a longiſh form, irregularly diſperſed, but on the belly become larger and more round; the tail ſhort, entirely white, except the two middle feathers, which are black; legs long and ſlender, and of a reddiſh brown colour."

It is not impoſſible this may be the young or variety of the Purple Sandpiper of this work, the bill not having obtained the reddiſh colour at the baſe, and the upper parts of the plumage not the full dark colours. The tail, however, is eſſentially different, and the bill appears to be ſhorter.

SANDPIPER-BROWN.

Tringa fuſca. *Ind. Orn.* ii. p. 733. 26. *Br. Zool.* ii. N° 195. *Lath. Syn. Sup.* p. 250.

This ſpecies Mr. Pennant obſerves to be in the collection of Mr. Tunſtal. Its ſize was that of a Jack Snipe. The bill black; the head, upper parts of the neck, and back, are of a pale brown, ſpotted with black; coverts of the wings duſky, edged with dirty white; under ſide of the neck white, ſtreaked with black; the belly white; tail cinereous; legs black. To this deſcription the ſame author adds, that it was bought in the London market.

We are in ſome doubts with reſpect to this bird being a diſtinct ſpecies, eſpecially as we know the Sandpiper claſs varies ſo much in plumage in the young and adult ſtates.

A young

A young ſpecimen of the Dunlin now before us, which was killed in July on the coaſt of Cornwall, exactly correſponds with the above, except that the head and back of the neck are of a ferruginous brown.

SANDPIPER-COMMON.

Tringa hypoleucos. *Lin. Syſt.* i. p. 250. 14. *Gmel. Syſt.* ii. p. 678. *Ind. Orn.* ii. p. 734. 28.

Tringa minor. *Raii Syn.* p. 108. A. 6. *Will.* p. 223. t. 55.

Guinetta. *Briſ.* v. p. 183. 2. t. 16. f. 2.—*Ib.* 8vo. ii. p. 260.

La Guignette. *Buf.* vii. p. 540.

Common Sandpiper. *Br. Zool.* ii. N° 204. t. 17. *Ib.* fol. 125. *Arct. Zool.* ii. N° 388. 23. *Will. Angl.* p. 301. t. 55. *Lath. Syn.* v. p. 178. 23. *Lewin, Br. Birds*, v. t. 172. *Walc. Syn.* ii. t. 148. *Pult. Cat. Dorſet.* p. 15.

The weight of this ſpecies is about two ounces; length ſeven inches and a half; irides duſky. The bill is brown, an inch and a quarter long, and ſlender. The head, back of the neck, and whole upper parts, as well as the ſides of the breaſt, light brown, of a gloſſy greeniſh hue on the back and ſcapulars, ſtreaked down the ſhaft of each feather with a narrow duſky line; the wing coverts with minute undulated lines; the cheeks and throat of a light colour; over the eye a whitiſh ſtreak; breaſt and

and belly white; thighs brown; quills brown, with a white ſpot on their inner webs, except the firſt; the four middle feathers of the tail like the back, with fine tranſverſe lines of duſky, outer one white, with brown bars, the intermediate ones either tipped or ſpotted white on their margins; legs brown.

The Common Sandpiper viſits this country in the ſpring, and chiefly frequents our lakes and rivers, on the borders of which it makes a neſt compoſed of moſs and dry leaves, which is moſt commonly placed in a hole in the bank. It lays four or five eggs of a dirty white, marked with duſky and cinereous ſpots, moſt at the larger end. When diſturbed it makes a piping noiſe as it flies; and when running on the ground the tail is conſtantly in motion. Great numbers breed on the banks of the lakes in Scotland. In the autumn theſe birds are very much infeſted with the *Hippoboſca birundines*.

It is probable many of the Sandpipers are capable of ſwimming, if by accident they wade out of their depth. Having ſhot and winged one of this ſpecies as it was flying acroſs a piece of water, it fell, and floated towards the verge; and as we reached to take it up, the bird inſtantly dived, and we never ſaw it again riſe to the ſurface; poſſibly it got entangled in the weeds, and was drowned. Other ſpecies we have known, when wounded and fallen into the water, make way on the ſurface with their legs,

legs, and do not drown like land-birds in general. It is known in ſome places by the name of Summer Snipe. Is found in moſt parts of Europe, even as far north as Siberia.

SANDPIPER-GREEN.

Tringa Ochropus. *Lin. Syſt* i. p. 250. 13. *Gmel. Syſt.* ii. p. 676. *Ind. Orn.* ii. p. 729. 12.

Tringa Aldrov. *Raii Syn.* p. 108. A. 7. 8. *Will.* p. 222. 223. t. 55.—*Ib. Angl.* p. 300, 301. t. 55. *Brif.* v. p. 177. 1. t. 16. f. 1.—*Ib.* 8vo. ii. p. 259.

Becaſſeau, *ou* Cul-blanc. *Buf.* vii. p. 534.

Green Sandpiper. *Br. Zool.* i. N° 201.—*Ib.* fol. 125. t. F. 2. f. 3. *Arct. Zool.* ii N° 389. *Lath. Syn.* v. p. 170. 12. *Lewin, Br. Birds,* v. t. 170. *Pult. Cat. Dorſet.* p. 15.

Wood Sandpiper. *Lin. Tranſ.* i. p. 130. t. 2.

This elegant ſpecies weighs about three ounces and a quarter; length full ten inches. The bill is an inch and a half in length, very ſlender, and duſky; irides hazel. Head, neck, and breaſt, are marked with numerous ſtreaks of duſky and aſh-colour, largeſt on the latter; over the eye a whitiſh ſtreak; back, ſcapulars, and wing coverts, brown, gloſſed with green; the two firſt marked with roundiſh ſmall white ſpots; lower part of the breaſt, belly, upper tail coverts, and chin, white; the feathers under the wings duſky, with ſmall bars of

white ſhaped like the letter V; quills duſky; tail white; the two middle feathers marked with four duſky bars; the two next on each ſide with three, the fourth with two, and the two outer with one bar; legs duſky green; toes united at the baſe by a ſmall membrane.

The Green Sandpiper is by no means plentiful in England; it moſtly frequents pools and ſmall ſhallow ſtreams; is a ſolitary bird; comes to us about the middle of September, and continues as late as the end of April, when it departs northward to breed. Is ſometimes ſeen in pairs before it leaves us. Is ſaid to be found in Siberia and Iceland, and alſo in America. When diſturbed it makes a very ſhrill whiſtling note as it flies. We have ſeen this bird as early as the ſecond of Auguſt.

SANDPIPER-GREENWICH.

Tringa Grenovicenſis. *Ind. Orn.* ii. p. 731. 16.

Greenwich Sandpiper. *Lath. Syn. Sup* p. 249. 38. *Lewin, Br. Birds*, v. t. 180. *Walc. Syn.* ii. t. 154.

This bird was killed at Greenwich, and is now in the collection of Dr. Latham, who has given it to the world as a new ſpecies: his deſcription therefore we take the liberty of borrowing.

It is the ſize of the Redſhank; weight near eight ounces; length twelve inches and a half.

Bill an inch and a half long, black; crown of the head

head reddiſh brown, ſtreaked with black; nape, cheeks, and neck, aſh-colour; the middle of the feathers duſky down the ſhaft; lower part of the neck and back black; the feathers margined on the ſides with pale ferruginous, and ſome of thoſe of the back at the tips alſo; chin nearly white; fore part of the neck very pale aſh-colour, as far as the breaſt, which is of a duſky white; belly, ſides, vent, and upper tail coverts on each ſide, and whole of the under ones, white; leſſer wing coverts aſh-colour; the greater the ſame, obſcurely margined with pale ferruginous; greateſt tipt with white; under wing coverts pure white; prime quills duſky, the ſhafts more or leſs white; ſecondaries and primores very little differing in length; the lower part of the back, rump, and middle of the tail coverts, aſh-colour; tail a little rounded at the end, browniſh aſh-colour, ſomewhat mottled with browniſh near the tips, and fringed near the end with pale ferruginous; legs duſky olive-green, bare an inch above the knee; the outer and middle toe connected at the baſe.

SANDPIPER-GREY.

Tringa ſquatarola. *Lin. Syſt.* i. p. 252. 23. *Gmel. Syſt.* ii. p. 682. *Ind. Orn.* ii. p. 729. 11.

Pluvialis cinerea. *Raii Syn.* p. 111. A. 3. *Will.* p. 229. t. 57.

Vanellus griſeus. *Briſ.* v. p. 100. 2. t. 9. f. 1. —*Ib.* 8vo. ii. p. 237.

Vanneau pluvier. *Buf.* viii. p. 68.

Grey Plover. *Will. Angl.* p. 309. t. 57. *Albin.* i. t. 76.

Grey Sandpiper. *Br. Zool.* ii. N° 191.—*Ib.* fol. 122. *Artt. Zool.* ii. N° 393. *Lath. Syn.* v. p. 168. 11.—*Sup.* p. 248. *Lewin, Br. Birds*, v. t. 169. *Walc. Syn.* ii. t. 146. *Pult. Cat. Dorſet.* p. 15.

This ſpecies rather exceeds the Golden Plover in ſize; the weight about ſeven ounces; length twelve inches.

Bill black, one inch and a quarter in length; irides duſky. The upper part of the head, the neck, back, ſcapulars, and wing coverts, duſky; the feathers more or leſs margined and tipped with aſh-colour, leaſt ſo on the head and neck; and on the coverts the grey predominates, and is almoſt of a white; the cheeks and throat white, with a few duſky lines; belly and under tail coverts white; quill-feathers black; the inner webs more or leſs white, as well as the ſhafts; from the fifth ſome white begins to appear on the outer web down the ſhaft, which increaſes in the next, and from the ſeventh to the tenth the whole of the outer web is white, except at the point; the tail is white, elegantly barred with black; on the middle feathers are ſix or ſeven; the inner webs of the lateral ones barred only near the end; the outermoſt has only one faint bar at the tip; legs duſky; back-toe

extremely ſmall; the claw almoſt adhering to the leg. The under ſcapulars, or thoſe long feathers underneath the wing at the baſe, in the ſeveral ſpecimens we have examined, were black; and the rump and upper tail coverts white, barred with black.

A variety is ſaid to have the forehead, throat, and rump, white; and the upper tail coverts white, edged with grey and pale yellow.

The Grey Sandpiper does not appear to be a plentiful ſpecies in England. We have bought it in the market at Briſtol, and have received it from our friend Mr. Boys of Sandwich, killed on that coaſt. It is not found here in the ſummer months, and of courſe retires northward to breed; frequents the ſea-ſhores only, and ſeldom more than ſix or ſeven are ſeen in a flock. Said to be found in Siberia and Carolina in large flocks.

SANDPIPER-HEBRIDAL.

Mr. Pennant has deſcribed a bird in the *Britiſh Zoology* under this title. Dr. Latham, in his *Synopſis*, has conſidered it as only a variety of the Turnſtone, with whom we do not heſitate to accord in opinion. *Vide* Turnſtone.

SANDPIPER-LITTLE.

Tringa puſilla. *Lin. Syſt.* i. p. 252. 20. *Gmel. Syſt.* ii. p. 681. *Ind. Orn.* ii. p. 737. 38.

Cinclus dominicenſis minor. *Brif.* v. p. 222. 13. t. 25. f. 2.—*Ib.* 8vo. ii. p. 269.

Little Sandpiper. *Br. Zool.* ii. N° 207. *Gen. Birds*, p. 65. t. 12. *Arct. Zool.* ii. N° 397. *Lath. Syn.* v. p. 184. 32. *Lewin, Br. Birds*, v. t. 175.

This is about the ſize of a Hedge Sparrow; the length between five and ſix inches. Bill brown, with the tip black; the head and all the upper parts brown, edged with black and pale rufous-brown; greater coverts and quills duſky, tipped with white; breaſt and belly white; tail duſky; legs black.

Mr. Pennant is the only perſon who has mentioned this ſpecies to be found in England, one having been killed near Cambridge in September.

Said to be found in northern Europe and in America, as well as St. Domingo.

SANDPIPER-PURPLE.

Tringa nigricans. *Lin. Tranſ.* iv. p. 40. t. 2.
Purple Sandpiper. *Walc. Syn.* ii. t. 155.
Sea Sandpiper. *Lin. Tranſ.* iv. p. 22. t. 1?
Tringa Maritima. *Gmel. Syſt.* ii. p. 678. 36?
Selninger Sandpiper. *Lath. Syn.* v. p. 173. 15? *Ind. Orn.* ii. p. 731. 18? *Arct. Zool.* ii. p. 480. C.

This ſpecies rather exceeds the Purre in ſize; the length eight inches and a half. The bill is ſlender, an inch and a quarter long, tapering towards

the point, a very little curved, and of a dull red-colour, except at the apex and ſides, which are duſky; irides hazel. The head and neck duſky black; eyelids whitiſh; the throat white; back and ſcapulars black, gloſſed with purple and edged with aſh-colour; the wing coverts black, tipped with white; the larger ones above the primores deeply ſo; quills black, ſlightly edged with white on the exterior webs, except three of the ſecondaries, which are almoſt wholly white; theſe, with the white-tipped coverts, form a ſlight oblique bar on the wing when extended; the ſhafts white; breaſt and all beneath white, prettily ſpotted with black, except the middle of the belly and vent; the rump, coverts of the tail, and four middle tail feathers, black, gloſſed like the back; the other tail feathers light cinereous; in all twelve; the legs and toes dull red; claws black and blunt; toes nearly divided to their origin; middle toe an inch long.

This bird was killed at Lougharne on the coaſt of Caermarthenſhire in January, in company with the Purre; two others were ſhot there the ſame winter, and were called by the fiſhermen Redlegs; but theſe did not come under our inſpection. The one from which the above deſcription is taken is now in my collection.

I have ſince been favoured with the ſkin of one from Mr. Boys of Sandwich, which was killed on that coaſt in the winter of 1799. It correſponded exactly

exactly with the above, except that the bill was rather longer and ſtrait, and the breaſt more duſky.

SANDPIPER-RED.

Tringa iſlandica. *Lin. Syſt.* i. Addend. *Gmel. Syſt.* ii. p. 682. 24. *Ind. Orn.* ii. p. 737. 39.
Scolopax ſubarquata. *Gmel. Syſt.* ii. p. 658. 25?
Red Sandpiper. *Br. Zool.* ii. N° 202. t. 72. *Arct. Zool.* ii. N° 393. *Lath. Syn.* v. p. 186. 34.
Aberdeen Sandpiper. *Br. Zool.* ii. N° 203.
Red Sandpiper. *Lewin, Br. Birds,* v. t. 177. *Pult. Cat. Dorſet.* p. 15.

This ſpecies is from eight to ten inches in length. The bill is brown, one inch and a half long, and a little bent downwards; head, hind part of the neck, and beginning of the back, duſky, marked with red; fore part of the neck and breaſt cinereous, mixed with ruſt-colour, and obſcurely ſpotted with black; leſſer wing coverts cinereous; quills duſky; ſecondaries tipt with white; the two middle feathers of the tail duſky, the others cinereous; legs long and black.

That given by Linnæus is on the upper parts like the Woodcock; the under parts ruſo-ferruginous; rump whitiſh, undulated with black; the ſhafts of the quills and tail feathers white.

Dr. Latham, in his *Synopſis*, has made the Aberdeen Sandpiper of the *Britiſh Zoology* a variety only

of this bird, and probably with great truth, as it differs only in the breaſt being of a reddiſh brown mixed with duſky, and the belly and vent white.

We cannot too often repeat the caution neceſſary in identifying the various ſpecies of this genus, eſpecially thoſe which rarely viſit us, when we know they are ſo ſubject to vary their plumage by age or ſeaſon; and it is more than probable they are multiplied already beyond their real extent.

The Red Sandpiper, it is ſaid, has appeared in great flocks on the coaſt of Eſſex; found alſo on the coaſts of New York, Labrador, and Nootka Sound; Iceland, and about the Caſpian Sea.

SANDPIPER-RED-NECKED.

We conceive this bird to be the young of the Purre not arrived at the full plumage, or perhaps the Dunlin in the ſame ſtate; both of which we have frequently ſeen with the head and neck of a ferruginous colour, ſoon after their arrival in the month of Auguſt. The deſcription given in Latham's *Synopſis*, Vol. V. p. 183, is as follows:

Size of the Purre. The bill is ſhorter than the head; the crown and hind part of the neck ſtriated ferruginous and black; fore part of the neck as far as the breaſt deep ferruginous; the reſt not unlike the Purre; legs black. *Vide* Purre.

SANDPIPER-SCALLOP-TOED. Vide *Phalarope.*

SANDPIPER-SEA.

We find this ſpecies in Latham's *Synopſis* under the title of Selninger Sandpiper, where it is deſcribed thus:

Size of a Starling. Upper parts varied with grey and black; middle of the back tinged with violet; fore part of the neck duſky; under parts of the body white; tail duſky; the four outer feathers ſhorteſt, and edged with white; legs yellow.

In Mr. Markwick's Catalogue of Birds of Suſſex, given in the *Linnæan Tranſactions*, we find the following deſcription:

Length, from the tip of the bill to the end of the tail, eight inches and an half; breadth, from tip to tip of the wings when extended, fifteen inches and an half. Bill an inch and a quarter long, yellow from the baſe half way, and the tips black; weight two ounces and an half. The head, neck, ſhoulders, and back, are of a dark duſky aſh-colour, with the edges of each feather on the back ſomewhat lighter; prime quill-feathers of the wings duſky, their ſhafts white, particularly the firſt; the ſecondary quills are lighter, with white tips, and the hindmoſt are almoſt white, having only one duſky ſpot. The tail is ſhort, and conſiſts of twelve feathers, of which the four middlemoſt are of a dark duſky colour, and the four on each ſide gradually ſhorter, and of a pale aſh-colour, edged with white. The under

ſide of the neck and breaſt duſky, with the feathers on the breaſt fringed with white. The chin, lower part of the breaſt, belly, thighs, and vent, white, except a few duſky ſpots on the ſides, thighs, and vent. The legs and feet are yellow, naked above the knees, and the toes entirely divided, without the leaſt connecting membrane between any of them. The claws are blackiſh.

This gentleman adds, that a ſmall flock, conſiſting of ten or twelve, was ſeen on the coaſt near Bexhill on the 8th of December, two of which came under his inſpection.

Upon compariſon of the above deſcription with that of our Purple Sandpiper, which has been alſo given in the ſame volume of the *Linnæan Tranſactions*, we can have but very little doubt but theſe two are one and the ſame ſpecies, allowing for the pencil of the different draftſmen, and the different manner in which naturaliſts are found to deſcribe the plumage of birds.

Whoever will be at the pains to compare the ſeveral deſcriptions together will find the leading characters agree, particularly the white feathers of the ſecondary quills and the tail; and much allowance is to be made for the variation in the colour of the bill and legs, which in twenty-four hours after death change materially, eſpecially from an orange-yellow to a dull red or rufous-brown. We think it proper to pen theſe doubts, although we have given each a diſtinct place in this work.

For the ſynonyms, ſee *Purple Sandpiper*.

SANDPIPER.

SANDPIPER-SPOTTED.

Tringa macularia. *Lin. Syſt.* i. p. 249. 7. *Gmel. Syſt.* ii. p. 672. *Ind. Orn.* ii. p. 734. 29.

Turdus aquaticus. *Briſ.* v. p. 255. 20.—*Ib.* 8vo. ii. p. 275.

La Grive d'eau. *Buf.* viii. p. 140.

Spotted Tringa. *Edw.* t. 277. f. 2.

Spotted Sandpiper. *Br. Zool.* ii. N° 196.—*Ib.* fol. 124. *Arct. Zool.* ii. N° 385. *Lath. Syn.* v. p. 179. 24. *Lewin, Br. Birds,* v. t. 173. *Walc. Syn.* ii. t. 149.

This is about the ſize of a Thruſh; length eight inches. Bill duſky, reddiſh at the baſe; the upper parts of the birds are of a greeniſh brown, marked with duſky ſpots on the head, of a longiſh form; theſe increaſe on the neck to the back, where they are much larger; the rump plain; the ſhoulders and wings marked with the ſame colour, but the ſpots are tranſverſe; the under ſide of the body is white, marked with duſky ſpots; the two middle tail-feathers greeniſh brown, the others white, croſſed with duſky lines; legs dull fleſh-colour.

The female has none of the ſpots underneath, except on the throat.

In the *Britiſh Zoology* it is ſaid that the ſpots on the upper parts are of a triangular form and black.

This ſpecies does not appear to be noticed in this country ſince the time of Edwards; that which he took his deſcription from was ſhot in Eſſex.

Said

Said to inhabit North America, and to breed in Pennſylvania and Hudſon's Bay.

SANDPIPER-WOOD.

Tringa Glareola. *Lin. Syſt.* i. p. 250. 13. β. *Faun. Suec.* N° 184. *Gmel. Syſt.* i. p. 677. *Ind. Orn.* ii. p. 730. N° 13.

Wood Sandpiper. *Arct. Zool.* ii. p. 482. G. *Lath. Syn.* v. p. 172. 13.

This ſpecies is about the ſize of a Jack Snipe, but of a more ſlender form.

Length, from the apex of the bill to the end of the tail, nine inches; to the end of the toes eleven inches and a half; weight two ounces and a quarter; bill not quite an inch and a quarter long, the baſe half duſky green, the other black, ſlender, a trifle bending downward at the point, upper mandible rather the longeſt, tapering to a blunt point; irides duſky.

From the bill to the eye a duſky ſtreak, above which, on each ſide, is white paſſing over the eye; the middle of the forehead and crown duſky black, ſtreaked with dirty white; coverts of the ears duſky; chin and throat white; neck duſky, cloſely and finely ſtreaked with dirty white, which gives it a cinereous hue, fore part lighteſt; breaſt, belly, ſides, vent, and under tail coverts, ſpotleſs white; the feathers on the back duſky black, with a purpliſh gloſs, marked with a dull yellowiſh ſpot

on

on each ſide the webs near the tip; ſcapulars the ſame, with ſeveral ſpots on the margins; the coverts and tertials of the wings the ſame, but without the purpliſh gloſs, and the ſpots inclining more to white on the coverts; the ſmaller coverts on the ridge of the wing plain duſky black; primary and ſecondary quills, and firſt row of greater coverts immediately impending them, black, ſlightly tipped with white, except three or four of the firſt feathers; the ſhaft of the firſt quill is white; upper part of the rump black, with a few fine ſtreaks of white; lower rump and upper tail coverts white, thoſe next the tail ſpotted with black; the tail conſiſts of twelve feathers, the middlemoſt rather the longeſt; theſe are barred with black and white alternately, a little oblique; on the outer webs are eight black bars, on the inner webs ſix; the next feather has ſix bars on the outer and four on the inner web; the third has five and three bars in the ſame manner; the fourth feather has five and one; the fifth and outer feathers are only ſpotted on the margin of the outer web, with one ſpot on the inner web of the former; the latter is plain white on the interior web; the black bars on the middle feathers do not exactly correſpond, thoſe on the inner webs riſe higher at the ſhaft, and often run into the ſuperior bar on the outer web; the legs are of an olive-green, long and ſlender, meaſuring three inches from the knee to the end of the middle toe, and bare of feathers one inch above the knee; the outer

toe

toe connected by a membrane as far as the firſt joint.

There is little doubt but this is the *Tringa Glareola* of Linnæus. It cannot be confounded with the *Tringa Ochropus*, or Green Sandpiper, by thoſe who have had an opportunity of comparing them. It differs materially from that bird by the ſuperior length of the legs; the plumage too is very different when compared; nor has it any of thoſe ſingular white marks under the wings, as in the Green Sandpiper, repreſenting the letter V. The tail alſo in that bird is nearly even at the tip, and is only partly barred; whereas this is barred quite to the baſe, is rather cuneiform, and the feathers more pointed than in that bird.

In the ſpecimen now before us, ſhot on the coaſt of ſouth Devon early in the month of Auguſt, the outer feather of the tail on each ſide is longer than the two ſucceeding ones, and equal in length to the fourth, from which they gradually increaſe in length to the middle ones, which exceed the outer by a quarter of an inch. Whether this ſingular form of the tail is to be depended on as permanent, future experience muſt determine, as at this ſeaſon, when birds are moulting, ſuch a circumſtance cannot be fully relied on, being well known that birds always loſe the correſponding feathers of the tail and wings nearly at the ſame time.

Mr. Markwick, in the firſt volume of the *Linnæan Tranſactions*, has given a figure of the Green Sandpiper,

piper, *Tringa Ochropus*, of Linnæus, which he conſidered as the *Glareola*, or Wood Sandpiper. That gentleman, however, in the ſecond volume of the ſame work, acknowledges the poſſibility of his bird being the Green Sandpiper, but ſuſpects the Wood Sandpiper to be a mere variety of that bird.

This bird, however, is perfectly diſtinct from that or any other ſpecies that has come within our knowledge, ſhould it not prove to be the Glareola.

The few authors who have deſcribed this ſpecies are ſilent with reſpect to the length of the legs, which in the bird here given are ſingularly long in proportion to the body, and by far ſuperior in length to any ſpecies of Sandpiper of equal ſize we are acquainted with. This circumſtance, however, may have been overlooked in a ſkin or ill-preſerved ſpecimen. In the markings our bird ſeems to correſpond with that little the few authors ſtate who have deſcribed the *Glareola*, except that they all make the back to be brown; but as we are aware of the difficulty in conveying the idea of colour by deſcription, and the near approach of ſome kinds of brown to duſky, allowance may be made. The ſex and ſeaſon of the year in which a bird is killed muſt alſo be conſidered. This was a male.

The Wood Sandpiper is ſaid to inhabit the moiſt woods of Sweden.

SCOOPER. Vide *Avoſet.*

SCOTER.

SCOTER.

Anas nigra. *Lin. Syſt.* i. p. 196. 7. *Gmel. Syſt.* ii. p. 508. *Ind. Orn.* ii. p. 848. 43. *Briſ.* vi. p. 420. 28. t. 38. f. 2.—*Ib.* 8vo. ii. p. 471.

Anas niger minor. *Raii Syn.* p. 141. A. 5. *Will.* p. 280. t. 74.

La Macreuſe. *Buf.* ix. p. 234. t. 16.

Scoter, *or* Black Diver. *Br. Zool.* ii. N° 273. —*Ib.* fol. 253. *Artt. Zool.* ii. N° 484.—*Sup.* p. 76. *Will. Angl.* p. 366. t. 74. *Lath. Syn.* vi. p. 480. 36. *Lewin, Br. Birds,* vii. t. 249. *Walc. Syn.* i. t. 59.

PROVINCIAL.

Black Duck.

This ſpecies of Duck weighs about three pounds; length twenty-one inches. The baſe of the bill is furniſhed with a knob, which, with the hill, is black, divided down the middle with a line of orange, which ſpreads over half the bill after quitting the knob, but does not reach the tip by half an inch; darkeſt on the knob; irides duſky; eyelids yellow. The plumage is wholly black, gloſſy about the head and neck; the legs and feet duſky; the tail is cuneiform, and conſiſts of ſixteen pointed feathers; feet long and broad. It is ſaid the knob is in ſome of a red colour, in others green.

The female has no knob at the baſe of the bill; the

the plumage more dull, with ſometimes a mixture of grey on the chin and breaſt; both ſexes want the nail at the end of the bill, uſual in this genus.

Theſe birds are great divers, and moſtly reſide at ſea diſtant from ſhore; but are only ſeen with us in the winter ſeaſon, at which time they are plentiful on ſome parts of the coaſt of France, where they are taken by nets as they are diving after ſhell-fiſh, which ſeems to be their principal food. We received two taken by accident in the ſame manner off the coaſt of South Wales. Is never obſerved to viſit our rivers or freſh-water lakes.

It is found in great plenty in moſt of the northern parts of the world, and particularly in the great lakes and rivers of Siberia, where it probably breeds.

The male of this ſpecies has nothing remarkable in the *trachea* or windpipe.

SCOUT. Vide *Guillemot-fooliſh.*
SCRABER. Vide *Guillemot-black.*
SCREECH. Vide *Thruſh-miſſel.*
SCREECH-MARTIN. Vide *Swift.*
SEA DOTTEREL. Vide *Turnſtone.*
SEA-HEN. Vide *Guillemot-fooliſh.*
SEA-MALL. Vide *Gull-common.*
SEA-MEW. Vide *Gull-common.*
SEA-TURTLE. Vide *Guillemot-black.*
SEA-PARROT. Vide *Puffin.*
SEA-PIE. Vide *Oyſtercatcher.*

SEDGE-BIRD. Vide *Warbler-ſedge.*

SERULA. Vide *Merganſer-red.*

SHAG.

Pelecanus Graculus. *Lin. Syſt.* i. p. 217. 4. *Gmel. Syſt.* ii. p. 574. *Ind. Orn.* ii. p. 887. 15.

Corvus aquaticus minor. *Raii Syn.* p. 123. A. 4. *Will.* p. 249. t. 63.

Phalacrocorax. *Briſ.* vi. p. 516. 2.—*Ib.* 8vo. ii. p. 496.

Petit Cormoran, *ou* Nigaud. *Buf.* viii. p. 319.

Shag, *or* Crane. *Will. Angl.* p. 330. t. 63. *Lath. Syn.* vi. p. 598. 14. *Lewin, Br. Birds,* vii. t. 264. *Walc. Syn.* i. t. 93. *Pult. Cat. Dorſet.* p. 21.

This ſpecies of Pelican is in weight about four pounds; length twenty-nine inches; breadth three feet ten inches. The bill duſky, near four inches long; ſides of the mouth and chin bare yellow ſkin, minutely ſpeckled with black on the latter. The whole bird appears, at a little diſtance, to be black, but on nearer inſpection the head and neck, upper breaſt and rump, are found to be gloſſed with green; the feathers are ſomewhat pointed on the upper part of the back, ſcapulars, and wing coverts, and beautifully gloſſed with purple, violet, and green, each feather regularly bordered with velvet-black; the under parts of the body leſs gloſſed with green; legs duſky black; middle claw ſerrated; the feathers next to the bare ſkin on the chin are uſually white.

The

The female weighs about three pounds and a quarter; length twenty-ſeven inches. The feathers on the upper parts are not ſo dark and gloſſed with the colours of the male; but the margin of the feathers on the ſcapulars and coverts are black; the under parts are duſky and grey mixed; the legs and toes of a duſky colour, lighteſt on the fore part; the irides of both ſexes green, and the tail is compoſed of twelve ſtiff feathers, duſky, daſhed with cinereous.

The above deſcription is taken from the birds ſhot from their neſt; but we have ſeen many others of a lighter colour, both above and below; and in ſome the belly is of a dirty white, mixed with browniſh aſh-colour; all of which had twelve feathers in the tail, and their weight did not exceed four pounds two ounces, which is the material diſtinction between this and the Corvorant, which weighs ſix or ſeven pounds, and the tail invariably compoſed of fourteen feathers. The habits of theſe two ſpecies ſomewhat differ. This is never known to viſit our freſh-water rivers, which the Corvorant frequently will, and in ſome places make their neſt in trees, on which they often perch by the ſides of rivers. The Shag keeps wholly to the ſalt-water, and breeds on our rocky coaſts, where it makes a neſt of ſticks and ſea-weed, and lays two or three white eggs of a long oval; their weight about an ounce and three quarters.

We have counted no leſs than thirty Shags neſts

on a ſmall rock a little detached from the ſhore, from which place we took ſome eggs and young birds.

Theſe birds, by reaſon of their weight in proportion to their feathers, ſwim deep in the water, ſhewing only their head, neck, and back; are expert divers, and devour a prodigious quantity of fiſh.

Linnæus ſeems to ſuſpect this bird may prove to be the young of the Corvorant; but the reverſe is now proved beyond doubt. Mr. Pennant mentions what we conſider as a variety with a creſt on the back of the head two inches long; in every other reſpect it exactly correſponds with the above deſcription of the male bird.

SHAG-CRESTED.

Pelecanus criſtatus. *Gmel. Syſt.* ii. p. 575. 21. *Ind. Orn.* ii. p. 888. 16.
Shag. *Br. Zool.* ii. N° 292. t. 102.—*Ib.* fol. 159.
Creſted Corvorant. *Arct. Zool.* p. 583. A.
Creſted Shag. *Lath. Syn.* vi. p. 606. 15.

This correſponds in every reſpect with the Common Shag, except that it has the feathers on the back of the head elongated into a ſort of pendant creſt. We are inclined to believe it only a variety occaſioned by age, although Mr. Pennant and Dr. Latham, as well as ſome other authors, make it a diſtinct ſpecies.

SHEARWATER.

SHEARWATER.

Procellaria Puffinus. *Lin. Syſt.* i. p. 213. 6. *Gmel. Syſt.* ii. p. 566. *Ind. Orn.* ii. p. 824. 11. *Briſ.* vi. p. 131. 1.—*Ib.* 8vo. ii. p. 395.

Puffinus Anglorum. *Raii Syn.* p. 134. A. 4. *Will.* p. 252.

Avis Deomedia, Shearwater. *Raii Syn.* p. 133. 1. & A. 2. *Will.* p. 251.—*Ib. Angl.* p. 332. 334.

Le Puffin. *Buf.* ix. p. 321.

Manks Puffin. *Edw.* t. 379. *Will. Angl.* p. 333.

Shearwater Petrel. *Br. Zool.* ii. N° 258.—*Ib.* fol. 146. t. M. *Arct. Zool.* ii. N° 462. *Lath. Syn.* vi. p. 406. 11.—*Sup.* p. 269. *Lewin, Br. Birds*, vi. t. 218. *Walc. Syn.* i. t. 90. *Pult. Cat. Dorſet.* p. 19.

This ſpecies of Petrel weighs ſeventeen ounces; length fifteen inches.

The bill is an inch and three quarters long, yellow, tipped with black. The head and whole upper ſide of the body, wings, tail, and thighs, black; the under parts, from chin to vent, white; the legs are weak and compreſſed; whitiſh before, and duſky behind.

This bird does not make its appearance often in the ſouth, but is found in the Orkney Iſles and in the Calf of Man in the breeding ſeaſon, where they take poſſeſſion of rabbit-burrows or other holes, and lay one white egg, blunt at each end. The young are fit to take in Auguſt, when great

numbers are killed and barrelled with ſalt, which the inhabitants boil and eat with potatoes. Are ſaid to make their appearance firſt in February, but not to ſettle at the breeding place till April, and depart by the beginning of September. Are called in the Iſle of Man Manks Puffin, in the Orknies Lyre.

SHELL, or SHELD-APPLE. Vide *Croſsbill.*

SHIELDRAKE.

Anas Tadorna. *Lin. Syſt.* i. p. 195. 4. *Gmel. Syſt.* ii. p. 506. *Ind. Orn.* ii. p. 854. 56. *Raii Syn.* p. 140. A. 1. *Will.* p. 278. t. 70, 71. *Briſ.* vi. p. 344. 9. t. 33. f. 2.—*Ib.* 8vo. ii. p. 477.

Tadorne. *Buf.* ix. p. 205. t. 14.

Shieldrake. *Br. Zool.* ii. N° 278.—*Ib.* fol. 154. t. Q. *Arct. Zool.* ii. p. 972. D. *Will. Angl.* p. 363. t. 70. 71. *Albin.* i. t. 94. *Hayes, Br. Birds,* t. 28. *Lath. Syn.* vi. p. 504. 51. —*Sup.* p. 275. *Don. Br. Birds,* t. 71. *Oſt. Men.* t. 51. *Lewin, Br. Birds,* vii. t. 248. *Walc. Syn.* i. t. 57. *Pult. Cat. Dorſet.* p. 20.

PROVINCIAL.

Bargander. St. George's Duck. Burrow Duck. Pirennet. Sly-gooſe.

This elegant ſpecies of Duck weighs about two pounds and a half, or rather more; length two feet three inches. The bill is red, and turns upwards, furniſhed

furniſhed with a ſmall knob at the baſe; the nail at the tip black; irides duſky. The head and part of the neck greeniſh-black; the reſt of the neck, back, rump, and upper tail coverts, white; on the breaſt is a broad band of a bay-colour, growing narrower as it paſſes backwards under the wings, and encloſes the lower part of the neck behind; through the middle of this band on the breaſt runs a black liſt, which ſpreads very broad at and near the vent; the ſcapulars black; coverts of the wings white, except ſome of the outer ones, which are black; the greater quills black; the four next the body bay on their exterior webs; the ten next gloſſy-green, tinged with copper, forming a ſpeculum on the wing; the under coverts of the tail ferruginous; tail conſiſts of fourteen white feathers, tipt with black; legs reddiſh fleſh-colour. Both ſexes are nearly alike.

The Shieldrake is not uncommon on many parts of our coaſt, and remains with us all the year. The female makes choice of a rabbit-burrow to depoſit her eggs in, which are numerous, ſometimes as many as ſixteen, and which ſhe covers with down from her own body. The neſt is generally near the water where ſhe leads her young ſoon after they are hatched. It is rarely met with remote from ſalt water; but if the eggs are taken and hatched under a hen, the young become tame, and may be kept in ponds; but it rarely breeds in confinement. A nobleman informed me he had

make a neſt of ruſhes, in which they lay ten or twelve rufous-coloured eggs.

Is found in Germany, Ruſſia, and America.

SHOVELER-BLUE-WINGED. Vide *Shoveler.*

SHOVELER-RED-BREASTED.

Anas rubens. *Gmel. Syſt.* ii. p. 519. 82. *Ind. Orn.* ii. p. 857. 62.

Red-breaſted Shoveler. *Br. Zool.* ii. N° 281. *Lath. Syn.* vi. p. 512. 57. *Pult. Cat. Dorſet.* p. 21.

This ſpecies of Duck is about the ſize of the laſt. The bill is alſo like that bird, but the colour is of a browniſh yellow; irides yellow. Breaſt and throat of a reddiſh brown; back brown, paleſt towards the ſides; the tips and pinions of the wings grey; quills brown; the reſt of a greyiſh brown; the ſpeculum or ſpot in the wing purple, edged with white; vent brown, with darker ſpots; tail ſhort and white; legs ſhort and ſlender; feet ſmall, and of a reddiſh brown-colour.

The colours of the female are more faint, and the ſpeculum of the wings blue.

This bird is ſaid to be ſometimes taken in the decoys in Lincolnſhire; but we believe it to be a rare ſpecies in England.

SHREITCH. Vide *Thruſh-miſſel.*

SHRIKE. A genus of birds, the characters of which are,

Bill ſtrait at the baſe, the end hooked, and furniſhed with a notch near the tip of the upper mandible.

The baſe not furniſhed with a cere, but with briſtles on the ſides of the mouth.

Tongue jagged at the end.

Toes, the outer one connected to the middle one as far as the firſt joint.

The Butcher-bird, or Shrike genus, ſeems to have puzzled many of the beſt naturaliſts in reſpect to the place theſe birds ſhould hold in the ſyſtem of Ornithology; and the great Linnæus had changed his opinion more than once, and at laſt left it in the Accipitrine order.

Mr. Pennant, in his laſt edition of *Britiſh Zoology*, brought theſe birds into the order of Pies; and Dr. Latham has judiciouſly followed the example in his *General Synopſis*.

SHRIKE-CINEREOUS.

Lanius excubitor. *Lin. Syſt.* i. p. 135. 11. *Gmel. Syſt.* i. p. 300. *Ind. Orn.* i. p. 67. 6.

Lanius, ſeu Collurio cinereus major. *Raii Syn.* p. 18. A. 3. *Will.* p. 53. t. 10. *Briſ.* ii. p. 141. 1.—*Ib.* 8vo. p. 197.

La Pie-grieſche griſe. *Buf.* i. p. 296. t. 20.

Greater

Greater Butcher-bird. *Will. Angl.* p. 87. t. 13.

Great Ginerepus Shrike. *Br. Zool.* N° 71. t. 33. — *Ib.* fol. p. 73. t. C. *Arct. Zool.* ii. N° 127. *Lewin, Br. Birds*, i. t. 30. *Lath. Syn.* i. p. 160. 4. *Walc. Syn.* i. t. 29. *Pult. Cat. Dorſet.* p. 4. *Don. Br. Birds*, iv. t. 87.

PROVINCIAL.

Mattages. Wierangle. Murdering-bird. Skreek, or Skrike. Night-jar. Mountain Magpie. French-pie.

The weight of this ſpecies rather exceeds two ounces; length ten inches; breadth fourteen. The bill is black, ſtrong, and much hooked at the end; irides duſky; the mouth beſet with ſtrong briſtles. From the baſe of the upper mandible through the eye a black ſtripe; the plumage of the whole upper part is pale aſh-colour, except the ſcapulars, which are white; the coverts of the wings black; quill-feathers black, with a white bar acroſs their middle, and many of them tipped with white; the under parts, from chin to vent, white; the tail conſiſts of twelve feathers of unequal length, which gives it a cuneiform ſhape; the two middle ones are black, the next ſlightly tipped with white, on the reſt the white gradually increaſes obliquely to the outer feather, which is only black at the baſe; legs black.

The

The female differs chiefly in the under parts, which are of a dirty white, marked with numerous ſemicircular brown lines.

A variety is ſpoken of which has the leſſer wing coverts and ſcapulars ſomewhat of a rufous-brown.

The Great Cinereous Shrike, or Butcher-bird, is rather a ſcarce bird in England. It is ſaid to breed in ſome of our mountainous ſituations, coming in May, and departing in September. However this may be in general, the only two which came under our inſpection were both males; one was killed on the fifteenth, and the other on the twenty-ſecond of November, in Wiltſhire.

In the ſtomach of one of theſe birds was a Shrew (*Sorex araneus, Lin. Syſt.*)

It is ſaid to make a neſt of heath and moſs, lined with wool and goſſamer, and to lay ſix eggs of a dull olive-green, ſpotted with black at the larger end.

This bird is ſometimes trained for catching ſmall birds in Ruſſia. Is ſaid to kill rats and mice, and valued in ſome countries for that property. When it has killed its prey it does not tear it like the Hawk, but fixes it to a thorn for the purpoſe of pulling it to pieces. Is ſaid to imitate the notes of ſome other birds by way of decoying them to their deſtruction.

SHRIKE-RED-BACK.

Lanius Collurio. *Lin. Syſt.* i. p. 136. 12. *Gmel. Syſt.* i. p. 300. *Ind. Orn.* i. p. 69. 11. *Briſ.* ii. p. 151. 4—*Ib.* 8vo. p. 202.

Lanius

Lanius minor ruffus. *Raii Syn.* p. 18. A. 4. *Will.* p. 54.

Merulæ congener alia. *Raii Syn.* p. 67. 13? *Will.* p. 144. 3.

L'Ecorcheur. *Buf.* i. p. 304. t. 21.

Leſſer Butcher-bird, *or* Fluſher. *Will. Angl.* p. 88. 2. 89. 3. *Albin.* ii. t. 14.

Red-backed Shrike. *Br. Zool.* i. N° 72.—*Ib.* fol. p. 74. t. C. 1. *Arct. Zool.* ii. N° 131. *Lewin, Br. Birds,* i. t. 30. *Lath. Syn.* i. p. 167. 25—*Sup.* p. 52. *Walc. Syn.* i. t. 30. *Pult. Cat. Dorſet.* p. 4.

This ſpecies weighs eight drams; length ſeven inches. Bill black, like the preceding; irides hazel. The upper part of the head and neck, and the rump are of a fine light grey; the forehead is black, which extends in a ſtreak through the eyes; the back, ſcapulars, and coverts of the wings, ferruginous; quills brown, ſlightly edged on the outer webs lighter; from the throat downwards of a bloſſom colour, paleſt about the vent; the tail is compoſed of twelve feathers, longeſt in the middle; theſe are more or leſs white at the baſe, black towards the end, and ſlightly tipped with white, except thoſe in the middle, which are wholly black, and the ſhafts are black throughout; the legs are black.

The female weighs two drams more. The whole upper parts are of a ferruginous-brown; the back of the neck daſhed with grey; the baſe of the upper mandible

mandible whitiſh; beneath the eye a brown ſtreak; breaſt and ſides dirty white, marked with numerous ſemicircular duſky lines; middle of the belly and vent white; the quills and tail-feathers brown; the outer web of the exterior feather of the latter white.

The Red-back Shrike, or Butcher-bird, viſits us in May, and departs in September; chiefly haunts incloſed moiſt ſituations, and makes its neſt in ſome thick hedge, compoſed of moſs and fibrous roots put together with wool, and lined with hair. It lays five or ſix eggs of a bluiſh-coloured white, with cinereous-brown ſpots, moſt at the larger end; ſometimes the eggs are white, with duſky ſpots; their weight from forty to fifty grains. When it has young, and you approach the neſt, the birds are clamorous, making a chattering noiſe. Its principal food is inſects, with which it feeds the young, particularly the Chaffer, or Dorbeetle. Theſe it transfixes on a thorn, tears off the body, leaving the clytra, wings, and head, behind. The male has alſo a chirping note, not very unlike the Houſe Sparrow; and we have heard it make a ſort of a ſong.

It appears to be a local ſpecies; is not uncommon in the north of Wiltſhire, and part of Glouceſterſhire and Somerſetſhire, particularly about Briſtol, where we have taken its young, and kept them for ſome time. Theſe lived in amity for about two months, when violent battles enſued, and two out of four were killed. The other two were chained

chained in the manner Goldfinches frequently are; they were extremely docile, would come to the call for the ſake of a fly, of which they were extremely fond; when raw meat was given them would endeavour to faſten it to ſome part of their open cage in order to tear it; would eat mice and ſmall birds cut in pieces, feather, fur, and bones, diſgorging the refuſe like the Hawk tribe.

One was killed by ſwallowing too large a quantity of mouſe-fur, which it could not eject, and was ſtrangled; the other became ſo fat that it expired in a fit in the hand while it was feeding on inſects.

The young reſemble the female till the following ſpring.

Is found in Ruſſia and France, and in Italy is common.

SHRITE. Vide *Thruſh-miſſel.*

SILK-TAIL. Vide *Chatterer.*

SISKIN. Vide *Aberdevine.*

SKRIKE, or SKREEK. Vide *Shrike-cinereous.*

SKUA. Vide *Gull-ſkua.*

SMEW.

Mergus albellus. *Lin. Syſt.* i. p. 209. 5. *Gmel. Syſt.* ii. p. 547. *Ind. Orn.* ii. p. 831. 6. *Briſ.* vi. p. 245. 3. t. 24. f. 1.—*Ib.* 8vo. ii. p. 425.

Mergus major cirrhatus. *Raii Syn.* p. 135. A. 3. *Will.* p. 254. t. 64.

——— Rheni. *Raii Syn.* p. 135. 5. *Will.* p. 255.

Smew, *or* White Nun. *Albin.* i. t. 89. *Br. Zool.*

Zool. ii. N° 262.—*Ib.* fol. 148. t. N. 1: *Lath. Syn.* vi. p. 428. 5.—*Ib. Sup.* p. 271. *Lewin, Br. Birds,* vi. t. 234. *Pult. Cat. Dorſet.* p. 19. *Walc. Syn.* i. t. 82.

This ſpecies of Merganſer rather exceeds the Teal in ſize: length about ſeventeen inches; weight twenty-four ounces. Bill near two inches long, and black. The head, neck, and whole under parts of the body, pure white; on each ſide the head an oval black ſpot, gloſſed with green; the feathers on the back of the head are long, forming a pendant creſt, black underneath; on each ſide the lower part of the neck is a curved black ſtreak, pointing forwards; the inner ſcapulars black; coverts on the ſides of the wings and greater quills are black; the middle coverts white; the leſſer quills, and the coverts immediately over them, black, tipped with white; tail cinereous; legs bluiſh grey.

The female weighs fifteen ounces; length ſixteen inches. The bill is lead-colour; head ferruginous, ſlightly creſted; the cheek, chin, throat, and all beneath, white; the breaſt clouded with grey; on the ſide of the head the ſame oval ſpot as in the male; back duſky, daſhed with cinereous; wings like the male; legs duſky, in ſome inclining to aſh-colour.

The markings of this ſex, as well as its ſize, correſponds ſo nearly to that of the Minute Merganſer, that we have long been inclined to believe that bird to be no other than this ſpecies, the young

males not attaining their full ſize and plumage for ſeveral years, but greatly reſembling that of the female.

Dr. Latham ſeems to have decided this matter without doubt, which we have mentioned under the head of Merganſer-Minute, to which we refer our readers.

We have, notwithſtanding this good authority, been inclined to keep them ſeparate in this work, giving to each the various ſynonyms of different authors, who have conſidered them as diſtinct ſpecies.

The *labyrinth* of this bird may be ſeen in the fourth volume of the *Linnæan Tranſactions*.

Theſe birds frequent freſh waters as well as the ſea in winter, but are not known to breed with us. It moſtly inhabits the northern lakes, and is driven to the ſouth only by ſevere weather.

SMEW-MINUTE. Vide *Merganſer-minute*.

SMEW-RED-HEADED. Vide *Smew*.

SNIPE. A genus of birds, the characters of which are,

Bill more than one inch and a half, ſlender, ſtrait, weak.

Noſtrils linear, lodged in a furrow.

Tongue ſlender, pointed.

Toes divided to their origin, or ſlightly connected; back-toe ſmall.

SNIPE-BROWN.

Scolopax griſea. *Gmel. Syſt.* i. p. 658. *Ind. Orn.* ii. p. 724. 33.

Brown Sniſe. *Arct. Zool.* ii. N° 369. *Lath. Syn.* v. p. 154. 28.

The weight of this ſpecies is three ounces and a quarter; length, from the apex of the bill to the end of the tail, eleven inches; to the end of the middle toe thirteen inches and a half.

Bill two inches and a half long, rather inclining downwards at the point, not ſo ſlender as that of the common Snipe, and ſpreads a little broad and compreſſed near the end, of a duſky colour, lighteſt at the baſe; upper mandible rather the longeſt, ſerrated within along the middle of the roof; both mandibles punctured or rough near the tip; irides duſky.

From the bill to the eye a duſky ſtroke, above that, paſſing over the eye, a white one; cheeks and throat white, with a few brown ſtreaks on the former; upper and under eyelids white; the crown of the head and neck cinereous brown, lighteſt on the fore part of the latter, and on the former the feathers are duſky in the middle; back and ſcapulars dark brown, margined with cinereous and rufous-brown; greater quills duſky, the interior ones, and the largeſt coverts immediately impending them, ſlightly tipped with white: ſhaft of the firſt quill white, the ſecond grey; the ſmaller coverts above

is a duſky line; the throat white; cheeks, neck, and upper breaſt, mottled with black and light ferruginous; the back and ſcapulars are black, barred with ferruginous-brown, and ſtriped with yellowiſh buff-colour in longitudinal lines; quills are black, the firſt edged with white; the ſecondaries tipped with the ſame; thoſe next the body are with their coverts ſtriated and barred with light ferruginous; lower breaſt and belly white; vent brown; upper tail coverts brown, barred with black; the tail conſiſts of fourteen black feathers, barred and ſpotted with dull orange-red towards the end, with a narrow bar of black near the tip, where it is pale rufous; legs vary, in ſome duſky or lead colour, others green.

This is a plentiful ſpecies in moſt parts of England, and is found in all ſituations, in high as well as low lands, depending much on the weather. In very wet times it reſorts to the hills; at other times frequents marſhes, where it can penetrate its bill into the earth after worms, which is its principal food.

Some few remain with us the whole year, and breed in the more extenſive marſhes and mountainous bogs. We have frequently taken the young before they could fly in the north of England and in Scotland. Near Penryn in Cornwall there is a marſh where ſeveral breed annually, and where we have taken their eggs, which are four in number, of an olivaceous colour, blotched and ſpotted with rufous-

rufous-brown; ſome with duſky blotches at the larger end, and ſome few elſewhere.

The neſt is made of the materials around it, coarſe graſs, and ſometimes heath. It is placed on a tump or dry ſpot, near a ſplaſh or ſwampy place; the eggs, like thoſe of the Lapwing, placed invariably with their ſmaller ends inwards, being much pointed; their weight three drams and a half. In the breeding ſeaſon the Snipe changes its note entirely from that it makes in the winter. The male will keep on wing for an hour together, mounting like a Lark, uttering a ſhrill piping noiſe; then deſcend with great velocity, making a bleating ſound not unlike an old Goat, which is repeated alternately round the ſpot poſſeſſed by the female, eſpecially while ſhe is ſitting on her neſt.

This bird has been met with in almoſt every part of the world.

SNIPE-GREAT.

Scolopax major. *Gmel. Syſt.* ii. p. 661. *Ind. Orn.* ii. p. 714. 4.

Great Snipe. *Br. Zool.* ii. N° 188. *Arct. Zool.* ii. p. 470. B. *Lath. Syn.* v. p. 133. 4. *Lewin, Br. Birds*, iv. t. 157. *Walc. Syn.* ii. t. 137. *Pult. Cat. Dorſet.* p. 14.

This bird weighs about eight ounces; length ſixteen inches. The bill is four inches long, like that of the Woodcock. Crown of the head black,

divided down the middle by a pale ſtripe; above and beneath each eye is another ſtripe of the ſame; the upper parts of the body very like the Common Snipe; the under parts white; on the neck, breaſt, and ſides, the feathers are edged with duſky; quills duſky; tail reddiſh brown or ruſt-colour, barred with black; the two middle feathers plain; legs black.

This ſpecies is rarely met with in England; it was firſt deſcribed by Mr. Pennant from a ſpecimen ſhot in Lancaſhire, now in the Leverian Muſeum. Is ſaid to have been met with in Kent; and we are informed one was killed in Wiltſhire, which weighed ſeven ounces and a half.

It is poſſible this bird is ſometimes killed together with the common ſpecies, and not diſcriminated by the ſportſman, who only takes it for a fine large Snipe.

Dr. Latham aſſures us he ſaved one from the hands of the cook at a friend's houſe, where, without diſcrimination, it was intended for the table with ſeveral of the common kind, and which is now in his muſeum.

SNIPE-HALF. Vide *Snipe-jack.*

SNIPE-JACK.

Scolopax Gallinula. *Lin. Syſt.* i. p. 244. 8. *Gmel. Syſt.* ii. p. 662. *Ind. Ornt.* ii. p. 715. 8.

Gallinago minima. *Raii Syn.* p. 105. A. 3. *Will.* p. 214.

Gallinago

Gallinago minor. *Briſ.* v. p. 303. 3. t. 26. f. 2.
—*Ib.* 8vo. ii. p. 287.
La petite Becaſſine. *Buf.* vii. p. 490.
Gid, Judcock, Jack Snipe. *Will. Angl.* p. 291. *Br. Zool.* ii. N° 189. t. 68.—*Ib.* fol. 121. *Arct. Zool.* ii. N° 367. *Albin.* iii. t. 86. *Lath. Syn.* v. p. 136. 8. *Lewin, Br. Birds,* iv. t. 159. *Walc. Syn.* ii. t. 139. *Pult. Cat. Dorſet.* p. 14.

This is about half the ſize of the Common Snipe, and weighs about two ounces: length eight inches and a half.

The bill is nearly two inches long, of a lead-colour, black at the point, and the ridge of the upper mandible light horn-colour; irides duſky. The crown of the head black, ſlightly edged with ruſt-colour, bordered on each ſide with a yellowiſh ſtreak, beneath which is a duſky one; and cloſe above the eye is another ſtreak of a light colour; from the bill to the eye is a duſky ſtroke; the neck is varied with ferruginous-brown and duſky, daſhed with cinereous above; the back, rump, and ſcapulars of a fine gloſſy changeable green and purple; the exterior webs of the latter deep buff-colour, forming two very conſpicuous lines from the ſhoulders to the tail; quills duſky; wing coverts duſky, bordered with cinereous and brown; the lower breast and all beneath white; the tail cuneiform, conſiſting of twelve pointed duſky feathers, daſhed more or leſs with ferruginous; legs greeniſh.

The Jack Snipe is not near ſo numerous as the other ſpecies, but is frequently found in the ſame places; is a more ſolitary bird, and never known to be gregarious. Amongſt ruſhes, or other thick covert, it will lie till in danger of being trod on, and when rouſed ſeldom flies far.

It comes to us later than the Common Snipe, and is never known to remain in this country during the breeding ſeaſon.

Is found in moſt parts of Europe, and in North America.

SNIPE-JADREKA.

Scolopax limoſa. *Lin. Syſt.* i. p. 245. 13. *Gmel. Syſt.* ii. p. 666. *Ind. Orn.* ii. p. 719. 18. *Briſ.* v. p. 262. 1.—*Ib.* 8vo. ii. p. 277.

Fedoa noſtra ſecunda. *Raii Syn.* p. 105. A. 5. *Will.* p. 216.—*Ib. Angl.* p. 293. II.

La Barge. *Buf.* vii. p. 500. t. 27.

Leſſer Godwit. *Br. Zool.* ii. N° 182.—*Ib.* fol. 120.

Jadreka Snipe. *Lath. Syn.* v. p. 146. 17. *Arct. Zool.* ii. N° 375. *Lewin, Br. Birds*, iv. t. 162.

PROVINCIAL.

Stone Plover.

This bird is ſeventeen inches in length; weight about nine ounces.

Bill four inches long, duſky, the baſe yellowiſh; irides white. The head and neck cinereous; cheeks, chin,

chin, rump, and vent, white; back brown; on the wings a line of white; two middle tail feathers black, the others white at the ends, which increaſes on the outer feathers, making the outer one almoſt white; legs duſky.

This ſpecies is ſaid to have been killed in England, but we believe very rarely. Is found in Iceland, Greenland, Sweden, and Ruſſia.

SNIPE-LEAST. Vide *Purre.*

SNIPE-POOL. Vide *Redſhank.*

SNIPE-SPOTTED.

Scolopax Totanus. *Lin. Syſt.* i. p. 245. 12? *Gmel. Syſt.* ii. p. 665. *Ind. Orn.* ii. p. 721. 24.

Totanus alter. *Raii Syn.* p. 186. 11. *Will.* p. 221.—*Ib. Angl.* p. 299.

Barker. *Albin.* ii. t. 71.

Spotted Redſhank. *Br. Zool.* ii. N° 186.

Spotted Snipe. *Lath. Syn.* ii. p. 148. 19. *Lewin, Br. Birds,* iv. t. 164.

This ſpecies is about the ſize of the Greenſhank. The head is of a pale aſh-colour, marked with oblong ſtreaks of black; back duſky, varied with triangular white ſpots; wing coverts aſh-coloured, ſpotted in the ſame manner; quills duſky; breaſt, belly, and thighs, white; the firſt thinly ſpotted with black; the middle feathers of the tail aſh-coloured, the ſide feathers whitiſh, barred with black; legs long, and of a bright red.

In

In Latham's *Synopſis* a variety of this bird is deſcribed, the length of which is ſixteen inches; the bill more than two inches long, and brown; orbits, chin, and throat, white, with a white line from the bill to the eye; cheeks and fore part of the neck white, with ſhort duſky ſtreaks; crown and upper part of the neck brown, ſtreaked with white; wings black, marked with elegant triangular ſpots of white; breaſt and belly white; legs long and of a rich yellow; ſometimes red.

We do not know whether this variety has been met with in England; the other has been killed with us, though rarely.

One lately came under our inſpection killed in Auguſt on the ſouth coaſt of Devonſhire. It weighed about ſix ounces; length thirteen inches; the bill two inches and a half long, and very ſlender; the upper mandible longeſt, and black; the baſe of the under mandible red. The head and back of the neck duſky, daſhed with cinereous, lighteſt on the latter; from the bill to the eye a duſky ſtreak; chin and throat white; back, ſcapulars, and wing coverts, duſky, ſpotted with grey; the fore part of the neck, breaſt, and all the under parts, mottled brown and white; lower part of the back and rump white; the coverts of the tail barred with black and white; the ſix firſt quill-feathers duſky black, the reſt more or leſs ſcallopped or barred with white on the edges of both webs; the firſt feather has a white ſhaft; the tail conſiſts of twelve

twelve feathers; the two middle ones rather longeſt; all obſcurely marked with numerous bars of black and brown; the edges ſcallopped with white; legs orange-red, four inches long from the end of the toe to the knee, and one inch bare above the knee.

SNIPE-SUMMER. Vide *Sandpiper-common.*

SNITE. Vide *Snipe-common.*

SNORTER. Vide *Wheatear.*

SNOW-FLEAK, *or* SNOW-BIRD. Vide *Bunting-Snow.*

SNOW-HAMMER. Vide *Bunting-tawny.*

SPARLING-FOWL. Vide *Dundiver.*

SPARROW-HOUSE.

Fringilla domeſtica. *Lin. Syſt.* i. p. 323. 36. *Gmel. Syſt.* ii. p. 925. *Ind. Orn.* i. p. 432. 1.

Paſſer domeſticus. *Raii Syn.* p. 86. A. *Will.* p. 182. *Briſ.* iii. p. 72. 1.—*Ib.* 8vo. i. p. 327.

Le Moineau. *Buf.* iii. p. 474. t. 29. f. 1.

Houſe Sparrow. *Br. Zool.* i. N° 127. t. 15.—*Ib.* fol. 107. *Arct. Zool* ii. p. 382. G. *Will. Angl.* p. 249. *Albin.* i. t. 62. *Lewin, Br. Birds,* ii. t. 77. *Lath. Syn.* iii. p. 248. 1.—*Ib. Sup.* p. 163. *Walc. Syn.* ii. t. 215. *Pult. Cat. Dorſet.* p. 12.

This well-known ſpecies of Finch weighs near ſeven drams; length about ſix inches. The bill is duſky; irides hazel. The crown of the head aſh-colour; round the eye, and between that and the bill,

bill, is black; behind the eyes, ſurrounding the back part of the head, bay; cheeks white; chin and under part of the neck black, mixed with grey; belly dirty white; the coverts of the wings are cheſnut and black mixed, with a whitiſh bar acroſs them; the back a mixture of black and rufous; quills duſky, with rufous edges; tail duſky, edged with grey; legs brown.

The bill of the female is lighter; behind the eye a line of white; the head and whole upper parts are brown, the under dirty white, daſhed with aſh-colour; no black on the chin or neck.

The Sparrow is too well known in every part of England; it inhabits the dwellings of the rich and poor, taking poſſeſſion of the humble thatched cottage in preference to the ſumptuous palace. Is rarely ſeen far from the habitation of man, as it delights in the ſpoil of his labour. The higheſt cultivated parts produce the greateſt quantity. It might be ſaid of this bird, as of ſome ſpecies of water-fowl, which remaining always within ſoundings, warn the mariner of his approach to land; ſo on the extenſive and dreary mountains not a ſparrow is ever to be ſeen; and the ſight of one beſpeaks ſome habitation near. It makes a neſt conformable to the place it chooſes for incubation, whether in a hole of a wall, in thatch, or under the tiles of a houſe, or in a Martin's neſt, it muſt conform to the ſize of the place; but when the neſt is made in a tree it is of large ſize, and covered at the

the top, compoſed of hay and ſtraw, lined warmly with feathers and fragments of thread or worſted, bits of cloth, or any refuſe material of that ſort found about houſes. It lays ſix eggs of a whitiſh colour, ſpotted with duſky and cinereous, weighing from forty-three to forty-eight grains.

Accidental varieties, ſuch as white, black, and yellowiſh ſpecimens, are ſometimes ſhot.

SPARROW-MOUNTAIN. Vide *Sparrow-tree.*

SPARROW-REED. Vide *Bunting-reed.*

SPARROW-REED-LESSER. Vide *Warbler-ſedge.*

SPARROW-TREE.

Fringilla montana. *Lin. Syſt.* i. p. 324. 37. *Gmel. Syſt.* ii. p. 925. *Ind. Orn.* i. p. 433. 2.

Paſſer montanus. *Raii Syn.* p. 87. 15. *Will.* p. 185. *Briſ.* iii. p. 79. 2.—*Ib.* 8vo. i. p. 329.

Loxia hamburgia. *Gmel. Syſt.* i. p. 854.

Pyrrhula hambergenſis. *Briſ.* iii. p. 314. 2.—*Ib.* 8vo. i. p. 396.

Friquet. *Buf.* iv. p. 389. t. 29. f. 2.

Le Hambouvreux. *Buf.* iv. p. 398.

Hamburg Tree Creeper. *Albin.* iii. t. 24.

Hamburg Groſbeak. *Lath. Syn.* iii. p. 149. 64.

Tree *or* Mountain Finch. *Br. Zool.* N° 128.—*Ib.* fol. 109. *Arct. Zool.* ii. N° 246. *Will. Angl.* p. 252. t. 25. *Albin.* iii. t. 66. *Edw.* t. 269. *Lewin, Br. Birds,* ii. t. 78. *Lath. Syn.* iii. p. 252. 2.—*Sup.* p. 163. *Walc. Syn.* ii. t. 216. *Don. Br. Birds,* iv. t. 88.

This

This ſpecies is rather leſs than the Houſe Sparrow; length five inches and a half. Bill black; irides greyiſh hazel. The head and nape cheſnut; chin black; a ſpot of the ſame colour behind the eye; the upper parts of the body rufous-brown, ſpotted with black, inclining to greeniſh towards the rump; ſides of the neck, the breaſt, and under parts, duſky white; wing coverts rufous, edged with black, and croſſed with two bars of white; the greater coverts black, with ferruginous edges; quills blackiſh, with rufous edges; tail even at the end; colour rufous-brown; legs pale yellow.

The female wants the black marks on the throat and ears, and the colours are duller.

The Tree Sparrow is much more local than the common ſpecies; it ſeems to be moſtly confined to Lancaſhire, Yorkſhire, and Lincolnſhire, and ſaid to be very plentiful in the laſt county, flying in large flocks in the winter ſeaſon, but has never been obſerved farther north.

It always makes its neſt in trees, which is ſaid to be compoſed of bents, and lined with feathers, and lays five eggs of a pale brown colour, with ſpots of a darker ſhade. Whether this bird is an inhabitant of the more ſouthern counties we have not been able to aſcertain.

SPARROW-WATER. Vide *Bunting-reed.*

SPARROW-WATER-LESSER. Vide *Warbler-ſedge.*

SPOONBILL. A genus of birds, the characters of which are,

Bill long, broad, flat, and thin, the end widening into a roundiſh form, not unlike a ſpoon.
Noſtrils ſmall, placed near the baſe.
Tongue ſmall and pointed.
Feet ſemipalmated.

SPOONBILL-WHITE.

Platalea leucorodia. *Lin. Syſt.* i. p. 231. 1. *Gmel. Syſt.* ii. p. 613. *Ind. Orn.* ii. p. 667. 1.
Platea, ſive Pelecanus. *Raii Syn.* p. 102. 1. *Will.* p. 212. t. 52. *Briſ.* v. p. 352. 1.—*Ib.* 8vo. ii. p. 300.
La Spatule. *Buf.* vii. p. 448. t. 24.
Spoonbill, *or* Pelican. *Albin.* ii. t. 66. *Will. Angl.* p. 288. t. 5.
White Spoonbill. *Br. Zool. App.* t. 9. *Arct. Zool.* ii. p. 441. A.—*Sup.* p. 66. *Lath. Syn.* v. p. 13. 1. *Oſt. Menag.* t. p. 61. *Lewin, Br. Birds*, iv. t. 142. *Walc. Syn.* ii. t. 123. *Pult. Cat. Dorſet.* p. 13.

Weight about three pounds and a half; length two feet eight inches.

The bill is near ſeven inches long, and three quarters of an inch broad in the narroweſt part; two inches towards the point in the largeſt part of

the ſpoon; colour black, ſometimes brown, with an orange-coloured ſpot near the tip of the upper mandible; it is alſo croſſed with ſeveral indentations and dotted protuberances; the irides in ſome grey, others reddiſh; the lore, and round the eyes and throat, the ſkin is bare and black. The whole plumage is white; ſometimes the quills are tipped with black; the legs are black, ſix inches long; thighs bare about half way; toes connected by a ſmall web, extending as far as the ſecond joint of the outer and firſt joint of the inner toe.

The Spoonbill is rarely met with in England. Mr. Pennant mentions that a flock of theſe birds migrated into the marches near Yarmouth in Norfolk in April 1774. We have alſo been aſſured it is ſometimes ſeen on the coaſt of Devonſhire in the winter. Is ſaid to build its neſt in high trees near to the ſea, and to lay three or four white eggs, powdered with a few pale red ſpots, about the ſize of that of a Hen.

SPRAT-LOON. Vide *Diver-ſpeckled.*

STANNEL. Vide *Keſtril.*

STARE. A genus of birds, the characters of which are,

Bill ſtrait, depreſſed.

Noſtrils guarded above with a prominent rim.

Tongue hard and cloven.

Toes, middle one united to the baſe of the outmoſt as far as the firſt joint.

STARE,

STARE, or STARLING.

Sturnus vulgaris. *Lin. Syſt.* i. p. 290. 1. *Gmel. Syſt.* ii. p. 801. *Ind. Orn.* i. p. 321. 1. *Raii Syn.* p. 67. A. 1. *Will.* p. 144. t. 37. *Briſ.* ii. p. 439. 1.—*Ib.* 8vo. i. p. 280.

L'Etourneau. *Buf.* iii. p. 176. t. 15.

Stare, *or* Starling. *Br. Zool.* i. N° 104. t. 46.—*Ib.* fol. t. P. 2. f. 1. *Arct. Zool.* ii. p. 331. A. *Will. Angl.* p. 196. t. 37. *Albin.* i. t. 40.—*Ib. Song Birds*, t. 3. p. 11. *Hayes, Br. Birds*, t. 32. *Lewin, Br. Birds*, ii. t. 56. *Lath. Syn.* iii. p. 2.—*Sup.* p. 137. *Walc. Syn.* ii. t. 195. *Pult. Cat. Dorſet*, p. 8.

Weight about three ounces; length eight inches and three quarters.

The bill is not quite an inch and a half long, bluiſh at the baſe, yellow at the point, and when opened the gape extends far back in the head; the noſtrils are ſurrounded by an oval prominent rim; irides hazel. The head, neck, and upper parts of the back are black, gloſſed with purple and green as viewed in different lights; the feathers on the neck are long, narrow, and pointed; the breaſt, lower part of the back, coverts of the wings, and rump, are black, ſhaded with dark green; the whole plumage, throat excepted, elegantly marked with ſmall ſpots, white on the breaſt, and yellowiſh brown on the head and back; the larger quill-feathers are duſky on the inner webs, and, except

the outer one, black on their exterior webs; the other quills are of a dark aſh-colour, bordered with green, and the whole edged with reddiſh brown; the tail is ſhort, compoſed of twelve feathers, much like thoſe of the wing; legs reddiſh brown.

This is a plentiful ſpecies; is found in almoſt every part of the old continent; many ſtay with us the whole year; but the vaſt flocks that are ſeen in ſevere winters probably migrate to this country in ſearch of food, and return northward in the ſpring. We have obſerved continued flights of theſe birds going weſtward into Devonſhire and Cornwall in hard weather, and return eaſtward as ſoon as the froſt breaks up. Their food is chiefly inſects, but in defect of theſe will eat grain. Great numbers are frequently taken in winter in our pigeon-houſes, where they rooſt for the ſake of warmth; but we believe rarely if ever ſuck the eggs of that bird, which has been attributed to them. It will, indeed, ſometimes build in ſuch places, but more frequently in the hole of a tree, and ſometimes in an old building. The neſt is made of dry graſs, on which it lays four or five light-blue eggs, about one dram three quarters in weight. The natural notes of this bird are a ſhrill whiſtle and a chattering noiſe; but in confinement, where it becomes very docile, is taught to imitate the human voice, and to whiſtle tunes.

The flight of the Stare is not undulated, but ſmooth and even, and it walks or runs upon the ground

ground like the Wagtail or Lark genus, and is rarely obſerved to hop like the Thruſh.

In the north of England it is called Chepſter, or Chep-Starling.

STEINGAL, or STONEGALL. Vide *Keſtril.*

STINT. Vide *Purre.*

STONE-CHAT, or STONE-CHATTER.

Motacilla rubicola. *Lin. Syſt.* i. p. 332. 17. *Gmel. Syſt.* ii. p. 969.

Rubetra. *Briſ.* iii. p. 428. 25. t. 23. f. 1.—*Ib.* 8vo. i. p. 429.

Sylvia Rubicola. *Ind. Orn.* ii. p. 523. 49.

Œnanthe, noſtra tertia. *Raii Syn.* p. 76. A. 4. *Will.* p. 169. t. 41.

Traquet. *Buf.* v. p. 215. t. 13.

Stone-chatter, Stone Smich, Moor-titling. *Br. Zool.* i. N° 159.—*Ib.* fol. 103. t. S. 2. f. 5. 6. *Will. Angl.* p. 235. t. 41. *Albin.* i. t. 52. *Hayes, Br. Birds,* t. 39. *Lath. Syn.* iv. p. 448. 46. *Lewin, Br. Birds,* iii. t. 108. *Walc. Syn.* ii. t. 239. *Pult. Cat. Dorſet.* p. 9. *Don. Br. Birds,* iv. t. 92.

PROVINCIAL.

Stone-ſmith. Blacky-top.

This ſpecies of Warbler weighs about five drams; length five inches and a quarter.

The bill is black, broad at the baſe, and beſet

with briſtles; irides duſky. The head and throat black; on each ſide the neck is a large white ſpot; the feathers on the back are black, edged with tawny; the lower part of the rump and tail coverts white, in ſome ſpotted; the breaſt is of a deep ruſt-colour; belly lighter; quill-feathers duſky; the leſſer ones edged on their exterior webs with bright ruſt-colour; the coverts of the wings black, ſlightly tipped and edged with ruſty brown, except ſome of the larger ſeries next the body, which are of a pure white, making a large conſpicuous ſpot on the wing; the tail is black, ſlightly tipped and edged with pale brown; legs black.

Male birds of the firſt year have the black feathers on the head mixed with rufous.

The female has no black on the head, but is of the ſame colour as the back, which is lighter than that part in the male; on the throat is an obſcure duſky ſpot; the white on the rump is alſo wanting, but it poſſeſſes that mark on the wings.

The Stone-chat has much the habits of the Whinchat, frequenting commons and furzy places, where they ſit upon the uppermoſt ſprays, and dart at every fly that paſſes, frequently returning to the ſame place again, like the Flycatcher, to which they ſeem as nearly allied as the Gold Finch or Pied Flycatcher, inſects and worms being their only food. This bird ſeems to be more general than the other ſpecies; it remains with us the whole year, and is one of the firſt birds that makes its neſt in the

ſpring.

ſpring. We have frequently found it with eggs before the middle of April.

The neſt is placed on or very near the ground, at the bottom of a furze, or ſome other buſh; is compoſed of moſs and bents, lined with hair, and ſometimes mixed with ſmall feathers. The eggs are five in number, of a blue colour, with ſmall rufous ſpots at the larger end, which in ſome are faint, in which caſe it is ſcarcely to be known from thoſe of the Whinchat; their weight is about thirty grains.

It is probable this ſpecies quits its uſual place of abode, the dreary moors and furzy commons, in very ſevere weather, as we have obſerved in ſome of the moſt temperate parts of England, Devonſhire and Cornwall; for, on the fall of a conſiderable quantity of ſnow, not one was to be ſeen where many were the day before; nor did they return for ſome time after the ſnow was melted. It is hardly poſſible they ſhould quit the kingdom in the middle of winter; and yet none were to be ſeen any where about that country.

It is likely a partial migration takes place in autumn, as we do not ſee ſo many in winter as in ſummer. In the early part of the ſpring it ſings very prettily, ſpringing into the air, and ſuſpending itſelf ſome time on wing. But as it breeds ſo early its ſong is of ſhort duration, as few birds ſing after their young are hatched.

When the young leave the neſt the old birds are

extremely clamorous and bold, and are as artful in enticing any one from their young as they are in concealing their neſt.

STONE-RUNNER. Many of the Sandpipers ſo called.

STONE-SMITH, or SMICH. Vide *Stone-chat.*

STORK.

Ardea Ciconia. *Lin. Syſt.* i. p. 235. 7. *Gmel. Syſt.* ii. p. 622. *Raii Syn.* p. 97. A. 1. *Will.* p. 210. t. 52. *Ind. Orn.* ii. p. 676. 9. *Briſ.* v. p. 365. 2. t. 32.—*Ib.* 8vo. ii. p. 305.

Cicogne blanche. *Buf.* vii. p. 253. t. 12.

White Stork. *Arct. Zool.* p. 455. C. *Will. Angl.* p. 286. t. 52. *Albin.* ii. t. 64. *Lath. Syn.* v. p. 47.—*Ib. Sup.* p. 234. *Lewin, Br. Birds,* iv. t. 144. *Walc. Syn.* ii. t. 125.

A ſpecies of Heron, about the ſize of a Turkey; length three feet three inches. The bill is ſeven inches three quarters long, of a fine red colour. The plumage is wholly white, except ſome of the ſcapulars, the greater coverts, and quill-feathers, which are black; the orbits of the eyes are bare and blackiſh; the ſkin, legs, and bare part of the thighs, are red.

The ſexes are alike.

This bird is rarely met with in England. Several inſtances, however, are on record; one was killed at Saliſbury in February 1790.

Vaſt

Vaſt numbers reſort to ſome parts of Holland to breed, and even as far north as Ruſſia, and depart in autumn ſouthward to winter in Egypt and other warmer climates. In moſt countries the inhabitants hold them in great veneration, probably from their property of deſtroying reptiles, on which they feed. They provide boxes for them on the tops of their houſes. In theſe places the Stork is ſo bold as to walk the ſtreets without fear. It makes a large neſt of ſticks, and lays from two to four eggs, about the ſize of that of a Gooſe, of a dirty yellowiſh white.

STORM-COCK. Vide *Petrel-ſtormy.*

STORM-FINCH. Vide *Petrel-ſtormy.*

SULA. Vide *Gannet.*

SWALLOW. A genus of birds, the characters of which are,

Bill ſhort, broad at the baſe, the point ſmall, and a little bending.
Noſtrils open.
Tongue ſhort, broad, cloven.
Tail forked; wings long.
Toes, three forward, one behind.

SWALLOW-CAR. Vide *Tern-black.*

SWALLOW-CHIMNEY.

Hirundo ruſtica. *Lin. Syſt.* i. p. 343. 1. *Gmel. Syſt.* ii. p. 1015. *Ind. Orn.* ii. p. 572. 1.

Hirundo domeſtica. *Raii Syn.* p. 71. A. 1.

Will. p. 155. t. 39. *Brif.* ii. p. 486. 1.—*Ib.* 8vo. i. p. 294.

Hirondelle de cheminée. *Buf.* vi. p. 591. t. 25. f. 1.

Chimney, *or* Common Swallow. *Br. Zool.* i. N° 168. t. 58.—*Ib.* fol. 96. *Will. Angl.* p. 212. t. 39. *Albin.* i. t. 45. *Arct. Zool.* ii. N° 330. *Lath. Syn.* iv. p. 561.—*Ib. Sup.* p. 192. *Lewin, Br. Birds,* iii. t. 123. *Walc. Syn.* ii. t. 251. *Pult. Cat. Dorfet.* p. 13.

This ſpecies weighs between five and ſix drams; length ſix inches and a half. Bill black; irides hazel. Forehead and chin ferruginous red; crown of the head and the whole upper parts black, gloſſed with purpliſh blue; breaſt and belly duſky white; the tail is much forked; the two middle feathers plain, the reſt marked on their inner webs, near the end, with an oval white ſpot; legs duſky.

The female has not the exterior feathers of the tail ſo long as in the other ſex.

The Swallow ſeems to be known in moſt parts of the world. In England it is a very common bird, coming to us in the ſpring, and departing in September to more mild climates, and is ſaid to winter in Senegal, and probably many other warm countries. It has taken the name of Chimney Swallow with us on account of breeding in our chimnies. In ſome countries it makes its neſt againſt rocks; with us it is not unuſual to find the neſt in outhouſes, upon the beams or rafters.

The

The neſt is made of mud plaſtered together and lined with feathers, and is open at top. The eggs are four or five in number, white, ſpeckled with ruſty red, weighing about thirty grains.

The food of this bird, as of the whole genus, is winged inſects, in catching which it is extremely dexterous; and, conſidering the velocity of its flight, the ſight muſt be incomparably quick. It makes its firſt appearance with us in April, ſometimes as early as the firſt week, if the weather is mild; and it ſometimes happens that after their arrival a long eaſterly wind prevails, which ſo benumbs the inſect tribe, that thouſands die for want of food. We recollect as late as the ninth of May the Swallows on a ſudden diſappeared from all the neighbouring villages around. The thermometer was at 42, and we were at a loſs to conceive what was become of theſe birds, which a day or two before were ſeen in abundance. But by chance we diſcovered hundreds collected together in a valley cloſe to the ſea ſide, at a large pool which was well ſheltered. Here they ſeem to have found ſome ſpecies of fly, though ſcarce ſufficient to ſupport life; for many were ſo exhauſted that after a ſhort time on wing were obliged to pitch on the ſandy ſhore.

Why it ſhould be neceſſary to account for the loſs of this tribe of birds in the winter, by making them to immerſe during that ſeaſon, is extraordinary, when at the ſame time no doubts have been entertained of the migration of other birds, whoſe

powers

powers on wing are far inferior. And yet there have not been wanting perſons who have declared they have ſeen them drawn up in nets, and reſtored from their benumbed ſtate. Others are ſaid to lay torpid in cliffs, hollow trees, and ſuch places: but even this more probable account is to be doubted, except perhaps with reſpect to a few of the latter broods, which had not ſtrength to undertake ſo long a flight. If we calculate the velocity of this bird on wing, and that it can and does ſuſpend itſelf in the air for fourteen or ſixteen hours together in ſearch of food, it cannot fly over a leſs ſpace than between two and three hundred miles in that time. We have frequently obſerved upon the downs Swallows follow, and repeatedly fly round with great eaſe, a horſe in a full trot, at a rate not leſs than ten miles an hour, in order to pick up the flies rouſed from the graſs by the motion of his feet.

It is certain, however, ſome few are ſeen in the winter months before Chriſtmas, although they had all diſappeared long before.

For further remarks we refer our readers to the *Hiſtory of the Martin.*

SWALLOW-SEA. Vide *Tern-common.*

SWALLOW-SEA-LESSER. Vide *Tern-leſſer.*

SWAN-WHISTLING, or WILD SWAN.

Anas Cygnus (ferus). *Lin. Syſt.* i. p. 194. *Gmel. Syſt.* ii. p. 501. *Ind. Orn.* ii. p. 833. 1.

Cygnus ferus. *Briſ.* vi. p. 292. 12. t. 28.—*Ib.* 8vo.

8vo. ii. p. 439. *Raii Syn.* p. 136. A. 2. *Will.* p. 272. t. 69.

Cygne ſauvage. *Buf.* ix. p. 3.

Wild Swan. *Br. Zool.* ii. N° 264.—*Ib.* fol. 149. t. Addend. *Will. Angl.* p. 356. t. 69. *Edw.* t. 150.

Whiſtling Swan. *Arct. Zool.* ii. N° 469.—*Ib. Sup.* p. 75. *Lath. Syn.* vi. p. 433.—*Sup.* p. 272. *Lewin, Br. Birds*, vii. t. 236. *Walc. Syn.* i. t. 55. *Pult. Cat. Dorſet.* p. 19.

PROVINCIAL.

Elk, *or* Hooper.

This is a very diſtinct ſpecies from the Common or Mute Swan, of which we ſhall take notice in this place merely to ſhew the difference; for as that bird is confined to a half-domeſticated ſtate, and is real property, it cannot claim a place amongſt the wild or native birds of this country.

The Whiſtling Swan is inferior in ſize to the other ſpecies; the weight from fifteen to twenty-five pounds; length four feet ten inches; breadth ſeven feet. The bill is between four and five inches in length, two-thirds of which is yellow from the baſe, and bare of feathers up to the eyes; the end part is black, running to an angle up the middle; irides of a very light yellow. The whole plumage is of a pure white; in ſome there are a few marks

of

of faint ruſt-colour on the head; the tail conſiſts of twenty feathers; legs black.

The tame Swan is not only much ſuperior in weight, but is at once diſtinguiſhed by a large black callous knob on the baſe of the bill; but the more eſſential diſtinction is in the extraordinary convolution of the windpipe, peculiar to the Whiſtling Swan. This enters a large cavity in the breaſt-bone, to the depth of more than two inches; then returns and forms a junction with the lungs.

Dr. Latham has given a very excellent figure of this wonderful formation in an eſſay on the *tracheæ* of birds in the fourth volume of the *Linnæan Tranſactions*. This, which both ſexes poſſeſs, makes a material diſtinction in the two ſpecies. What peculiar uſe this is of, we are at a loſs to determine. The common Swan is ſeldom heard to make any noiſe, and that a very faint one; whereas this bird has a prodigious loud call that may be heard to a great diſtance, which it frequently utters as it is flying, and reſembles greatly that of the Cuckow.

Having killed one of this ſpecies out of a flock of ten or twelve, its companions flew round ſeveral times, making a moſt melancholy cry before they flew off. This put us in mind of the ſolemn dirge of the dying Swan deſcribed by the ancient poets, and may poſſibly have given riſe to thoſe accounts; only by them it is made to be ſung by the dying bird.

This ſpecies viſits the lakes of Scotland every

winter, but comes more ſouthward only in ſevere weather; ſome few are ſaid to breed in the Orkney iſles. It is found in all the northern countries, Iceland, Lapland, the deſerts of Tartary and Siberia, as far as Kamtſchatka; alſo in America; and not uncommon at Hudſon's Bay.

Is ſaid to lay only four eggs.

SWIFT.

Hirundo Apus. *Lin. Syſt.* i. p. 344. 6. *Gmel. Syſt.* ii. p. 1020. *Raii Syn.* p. 72. A. 4. *Will.* p. 156. t. 39. *Ind. Orn.* ii. p. 582. 32. *Briſ.* ii. p. 512. 15.—*Ib.* 8vo. i. p. 301.

Le grand Martinet. *Buf.* vi. p. 643.

Swift. *Br. Zool.* ii. N° 171. t. 57.—*Ib.* fol. 97. *Arct. Zool.* ii. N° 334. *Will. Angl.* p. 214. t. 39. *Albin.* ii. t. 55. *Lath. Syn.* iv. p. 584. 34. *Lewin, Br. Birds,* iii. t. 126. *Walc. Syn.* t. 254. *Pult. Cat. Dorſet.* p. 13.

PROVINCIAL.

Screech, or Screech Martin. Black Martin.

This ſpecies of Swallow is nearly an ounce in weight; length near eight inches; breadth about eighteen.

The bill is black; irides duſky. The whole plumage is black, except the chin, which is whitiſh; the wings are extremely long in proportion, and the legs ſo ſhort that it riſes from the ground with difficulty; the tail is forked; legs and toes black.

It

Sarcelle d'été. *Buf.* ix. p. 268.
Anas Balbul. *Gmel. Syſt.* ii. p. 443. 124.
Summer Teal. *Will. Angl.* p. 373. t. 76. *Albin* iii. t. 103, 104. *Lath. Syn.* vi. p. 552. 89.

This ſpecies of Duck weighs about twelve ounces; length fourteen inches and a half. The bill is black; irides light hazel. Head and neck bay; on the ſide of the head a green patch paſſing backwards, bordered beneath with a whitiſh line; the lower part of the hind neck, upper part of the back, and part of the ſcapulars, as well as the ſides of the body, a mixture of black and white in fine undulated lines; lower part of the neck before and breaſt whitiſh, marked with roundiſh ſpots of black; belly of the ſame colour, without ſpots; vent black, bounded with buff-colour; wing coverts brown; quills duſky; ſome of the ſecondaries wholly black, and others gloſſy green, on their outer webs, forming a ſpeculum on the wing; the coverts immediately over theſe are tipt with white; the tail is cuneiform, conſiſting of ſixteen brown feathers, edged with whitiſh; legs duſky brown.

The female has the head, neck, back, and ſides of the body, brown, the feathers more or leſs edged with whitiſh; belly and vent white; ſpeculum in the wing like the male.

The male of this ſpecies has a bony labyrinth in the lower part of the windpipe. The Teal is the ſmalleſt of the Duck tribe. It comes to us in winter, and

and frequents our freſh waters in ſmall flocks. Many are caught in the decoys; ſome few breed with us.

Mr. White, in his *Natural Hiſtory of Selborne*, mentions young Teal having been taken on the verge of a pond in Wolmer Foreſt.

It is alſo ſaid to breed in the moſſes about Carliſle; and we are informed they have been known to breed in confinement.

The neſt is not uncommon in France. It is made of ruſhes lined with down, placed on the ſide of a pond ſo as to riſe or fall with the water. The eggs are about the ſize of thoſe of a Pigeon, of a dirty white, ſpotted with brown. This bird, like moſt of the Duck tribe, is ſubject to variety, which has occaſioned it to be made into two or three ſpecies, to one of which has been given the name of Summer Teal.

TEAL-SUMMER. Vide *Teal* and *Garganey*.

TERN. A genus of birds, the characters of which are,

Bill ſtrait, ſlender, ſharp-pointed.
Noſtrils linear.
Tongue ſlender, ſharp.
Wings very long.
Tail forked.
Feet ſmall, webbed.
Back toe ſmall.

TERN-BLACK.

Sterna fiſſipes. *Lin. Syſt.* i. p. 228. 7. *Gmel. Syſt.* ii. p. 610. *Ind. Orn.* ii. p. 810. 23.

Sterna nigra. *Briſ.* vi. p. 211. 4.—*Ib.* 8vo. ii. p. 417.

Larus niger Geſneri. *Raii Syn.* p. 131. A. 3. *Will.* p. 269.

Larus niger fidipes alis longioribus Aldr. *Raii Syn.* p. 131. 4. *Will.* p. 270. v. t. 68.

Larus minor, fidipes noſtras. *Raii Syn.* p. 132. A. 6. *Will.* p. 270. iv.

Hirondelle de mer noire, *ou* l'Epouventail. *Buf.* viii. p. 341.

Black Tern. *Br. Zool.* ii. N° 256.—*Ib.* fol. 145. t. L*. 1. f. 1. *Arct. Zool.* ii. N° 450. *Lath. Syn.* vi. p. 366. 22.—*Sup.* p. 267. *Lewin, Br. Birds*, vi. t. 206. *Walc. Syn.* i. t. 122. *Don. Br. Birds*, iv. t. 74.

PROVINCIAL.

Cloven-footed Gull. Scare-Crow.

This ſpecies is leſs than the Common Tern; weight about two ounces and three quarters; length ten inches.

Bill black; irides duſky. Forehead, ſides of the head beneath the eyes, throat, and fore part of the neck, white; the reſt of the head, back of the neck, and under part of the body, black; back, wings, and tail, deep aſh-colour; vent and under tail co-

verts

verts white; the tail is leſs forked than in either of the other ſpecies; the outer feathers edged with white; legs dull red, the webs much indented or ſemipalmated.

In ſome the forehead and fore part of the neck are mottled with black; and as moſt authors have omitted, or at leaſt have not made mention of any white on the forehead, ſuch is probably another variety.

The female has no white about the head.

The Black Tern has all the actions and manners of the other ſpecies, but ſeems to prefer freſh-water fiſh and inſects to marine. It breeds on the verge of pools in ſwampy places, and never, that we could trace, on the ſea-ſhore, but frequently remote from the ſea. It is found on the fenny parts of Lincolnſhire and Cambridgeſhire, and is called at this laſt place Car-Swallow. Is plentiful about the reedy pools on Romney Marſh in Kent, where they firſt appear the latter end of April or beginning of May, and breed in the ſedgy places on the verge of the pools; and though very near the ſea is rarely ſeen on the ſhores till after the breeding ſeaſon, and then not commonly. It lays three or four light olive-brown eggs, blotched and ſpotted with brown and black, about the ſize of thoſe of the Magpie.

On the ſea-ſhore that bounds the above extenſive marſh all the other ſpecies are found during the incubating ſeaſon.

TERN-BROWN.

Sterna obſcura. *Gmel. Syſt.* i. p.608. *Ind. Orn.* ii. p. 810. 25.

Sterna fuſca. *Raii Syn.* p. 131. A. 15. *Will.* p. 268. VIII.

Brown Tern. *Br. Zool.* ii. N° 253.—*Ib.* fol. 143. *Will. Angl.* p. 352. *Lath. Syn.* vi. p. 368. 23.

This we believe to be no other than the young of the Common Tern, having ſeen them in the plumage deſcribed by Ray and Willughby, and copied by various authors ſince. The head is black; wings partly brown and aſh-colour; the reſt of the plumage above brown, beneath white; the tail not forked.

Mr. Pennant has placed this amongſt his Gulls, no doubt from its not having a forked tail; but he expreſſes a ſuſpicion that it is the young of the Greater Tern. Dr. Latham has done the ſame.

TERN-COMMON.

Sterna Hirundo. *Lin. Syſt.* i. p. 227. 2. *Gmel. Syſt.* ii. p. 606. *Ind. Orn.* ii. p. 807. 15.

Hirundo marina. *Raii Syn.* p. 131. A. 1. *Will.* p. 268. t. 68.

Sterna major. *Briſ.* vi. p. 203. 1. t. 19. f. 1.—
—*Ib.* 8vo. ii. p. 415.

La grande Hirondelle de mer. *Buf.* viii. p. 331. t. 27.

Sea Swallow. *Albin.* ii. t. 88. *Will. Angl.* p. 352.

Common,

TER

Common, *or* Greater Tern. *Br. Zool.* ii. N° 254. t. 90.—*Ib.* fol. 144. t. L*. *Lath. Syn.* vi. p. 361. 14. *Lewin, Br. Birds*, vi. t. 204. *Walc. Syn.* i. t. 119. *Don. Br. Birds*, i. t. 23. *Pult. Cat. Dorſet.* p. 18.

PROVINCIAL.

Gull-Teazer.

The weight of this ſpecies is about four ounces and a quarter; length fourteen inches. Bill two inches and a half long, of a crimſon colour, black at the end, and very ſharp-pointed; irides duſky. The top of the head, taking in the eyes and nape, black; beneath the eyes, and from thence to the noſtrils, the neck, and all the under parts, white; the back and wings are aſh-colour; quills grey; the outer ones darkeſt, ſhafts white; tail much forked, and white, except the outer web of the exterior feather, which is black; legs crimſon; claws black.

Of the four ſpecies of Terns which viſit this country in ſpring, this is the moſt plentiful. It frequents our flat, ſandy, or ſhingly ſhores, preferring the latter; amongſt which it lays three or four eggs, without any neſt. Theſe are about the ſize of a Pigeon's, of an olivaceous brown, blotched and ſpotted with duſky.

It is found in great abundance on part of the

Suſſex and Kentiſh coaſts, particularly about Winchelſea, and from thence to Dungeneſs; is a noiſy, reſtleſs bird, conſtantly on wing in ſearch of inſects and ſmall fiſh, after which it darts into the water with great force, ſeizes, and inſtantly returns; for though it is web-footed is never ſeen to ſwim or dive. It is commonly known by the name of Sea Swallow, from its actions on wing being ſimilar to that bird. On the ſouth coaſt of Devonſhire is called Gull-teazer, as it is frequently ſeen to purſue and perſecute the leſſer Gulls till they diſgorge, which it dexterouſly catches before it reaches the water. Whether this property is common to the other ſpecies we have not been able to aſcertain; but in general their manners and habits are very ſimilar; none of which have ever been obſerved to ſettle on the water. This is ſometimes, on its firſt arrival, found ſkimming over freſh-water rivers and lakes for a few days; but we do not find they breed in ſuch places in England. It is found in moſt of the northern parts of Europe during the breeding ſeaſon, even as far as Greenland and Spitzbergen, as well as North America, where in New England it is called Mackarel Gull; and at Hudſon's Bay is known by the name of Black-head. Mr. Pennant calls it the Greater Tern. The young birds are mottled with brown and white, and are moſt probably the Brown Tern deſcribed by Ray and other ornithologiſts.

TERN-GREATER. Vide *Tern-common.*

TERN-LESSER.

Sterna minuta. *Lin. Syſt.* i. p. 228. 4. *Gmel. Syſt.* ii. p. 608. *Ind. Orn.* ii. p. 809. 19.

Sterna minor. *Briſ.* vi. p. 206. 2. t. 19. f. 2. —*Ib.* 8vo. ii. p. 416.

Larus piſcator. *Raii Syn.* p. 131. A. 2. *Will.* p. 269. II.

La petite Hirondelle de mer. *Buf.* viii. p. 337.

Leſſer Sea Swallow. *Albin.* ii. t. 90. *Will. Angl.* p. 353. t. 68.

Leſſer Tern. *Br. Zool.* ii. N° 255. t. 90.—*Ib.* fol. 144. t. L. 2. *Arct. Zool.* ii. N° 449. *Lath. Syn.* vi. p. 364. 18. *Lewin, Br. Birds,* vi. t. 205. *Walc. Syn.* i. t. 121. *Don. Br. Birds,* iv. t. 96. *Pult. Cat. Dorſet.* p. 18.

PROVINCIAL.

Richel-bird.

This is the ſmalleſt of the tribe, meaſuring about eight inches and a half in length; weight about two ounces. Bill yellow, tipped with black; irides duſky. The forehead is white; the reſt of the head above and the nape black; from the bill to the eye a black ſtreak; the ſides of the head beneath the eyes, the neck, and all the under parts, pure white; the back, ſcapulars, and wings, pale grey, darkeſt on the quills; tail white; legs yellow.

This elegant little ſpecies has all the habits of the

Common Tern, and breeds in the ſame places, but is far leſs numerous. We have with no ſmall difficulty found their eggs amongſt the ſhingle, which were always two in number, placed in a ſmall depreſſion, without any neſt; they are about the ſize of thoſe of the Black Tern, of a very pale brown, ſpotted all over with cinereous and duſky, but not ſo pointed as the eggs of that bird.

What has been ſaid of the manners and habits of the Common Tern will equally anſwer for this; to that ſpecies therefore we refer our reader. It is not, however, always found with that bird, as this is ſometimes found to breed where the other is not.

TERN-SANDWICH.

Sterna cantiaca. *Gmel. Syſt.* ii. p. 606.

Sterna Boyſii. *Ind. Orn.* ii. p. 806. 10.

Sandwich Tern. *Lath. Syn.* vi. p. 356. 9.—*Sup.* p. 266. *Boys, Sandwich*, t. p. 851. *Lewin, Br. Birds*, vi. t. 203. *Walc. Syn.* i. t. 120. *Don. Br. Birds*, v. t. 120.

VAR.

La Guiſette. *Buf.* viii. p. 339.

Cloven-footed Gull. *Albin.* ii. t. 82?

Sterna nævia. *Lin. Syſt.* i. p. 228. 5. *Gmel. Syſt.* ii. p. 609. *Briſ.* vi. p. 216. 6. t. 20. f. 1.—*Ib.* 8vo. ii. p. 418.

Kamtſchatkan Tern. *Arct. Zool.* ii. p. 525. A. *Lath. Syn.* vi. p. 358. 9. var. A.

This

This is the largeſt of the Britiſh ſpecies; length about eighteen inches.

The bill is black; irides duſky. The upper part of the head and nape, taking in the eyes, is black; the reſt of the head, neck, under parts of the body, and tail, white; back, ſcapulars, and wings, cinereous-grey; the firſt five or ſix prime quills are tipped with black; the inner webs more or leſs white towards the baſe; tail conſiderably forked; the outer feathers daſhed with cinereous on the exterior webs; legs duſky, with a tinge of red.

Young birds not maturely feathered are more or leſs clouded with brown on the upper parts of the body and wings, and the head ſpotted with white.

This bird was firſt noticed by Mr. Boys of Sandwich, where it is not uncommon, and communicated to Dr. Latham, who firſt gave it to the world. It does not appear to be ſo plentiful as either of the other ſpecies, for in a whole ſummer's reſidence on the coaſt of Suſſex and Kent, where the others are in plenty, we were not able to procure more than two ſpecimens; nor could we ever find where it bred. It comes to us and retires about the ſame time as the others, and has no doubt been confounded for the Common Tern, to which it bears very great reſemblance, except in ſize, and in the colour of the bill and legs; the bill of this is alſo much ſtronger, the legs much longer, and the tail not ſo much forked. It has, however, all the manners and habits of the common ſpecies, as far

as we have been able to collect. We are informed it breeds on the Sandwich coaſt, but have never with certainty heard of its eggs being found. We were favoured with an egg ſaid to belong to this bird from Mr. Lewin, and have ſeen another from the ſame gentleman in the cabinet of Dr. Latham; both of which appear in ſize and colour to be that of the Common Tern.

THROSTLE.

Turdus Muſicus. *Lin. Syſt.* i. p. 292. 2. *Gmel. Syſt.* i. p. 809. *Raii Syn.* p. 64. A. 2. *Will.* p. 138. t. 37. *Ind. Orn.* i. p. 327.

Turdus minor. *Briſ.* ii. p. 205. 2.—*Ib.* 8vo. i. p. 214.

La Grive. *Buf.* iii. p. 280.

Throſtle, Mavis, *or* Song Thruſh. *Will. Angl.* p. 188. *Br. Zool.* i. N° 107.—*Ib.* fol. 91. t. P. f. 2. *Arct. Zool.* ii. p. 342. C. *Albin.* i. t. 34.—*Ib. Song Birds*, t. 2. *Lewin, Br. Birds*, ii. t. 58. *Lath. Syn.* iii. p. 18. 2.—*Sup.* p. 139. *Walc. Syn.* ii. t. 198. *Pult. Cat. Dorſet.* p. 10.

This common ſpecies of Thruſh weighs about three ounces; length nine inches. The bill is nearly an inch long, duſky, the under mandible yellowiſh at the baſe; irides hazel. The head and whole upper parts are of a yellowiſh brown, with a few obſcure duſky lines on the former; the throat, neck, and ſides, yellowiſh; breaſt and belly white,

elegantly

elegantly ſpotted on the former with triangular duſky ſpots, ſomewhat reſembling arrow-heads pointing upwards; the under coverts of the wings dull orange-yellow; legs light brown.

This well-known bird is admired by every one for its ſong. Every wood and grove re-echoes with its melodious notes in the ſpring, frequently beginning its tuneful lays as early as February if the weather is mild, contending with the Miſſel in its love-ſtrained notes. As the ſong is a prelude to incubation, ſo this ſpecies makes its neſt in March, compoſed of dried graſs and green moſs externally, and plaſtered within with rotten wood mixed with cowdung or clay, which is ſo compact as to hold water, and ſometimes proves fatal to their eggs, for in a rainy ſeaſon we have often found it full. It lays four or five blue eggs, ſpotted with black at the larger end; their weight from eighty to ninety grains. The neſt is placed ſometimes on a ſtool or ſtump of a tree, very near the ground, or againſt the ſide of a tree, and frequently in a hedge or ſolitary buſh.

The Throſtle remains in England the whole year, but is ſuppoſed to quit the more northern parts in winter. It is not, however, gregarious with us at any time, although it has been obſerved to paſs through Livonia, Courland, and Proſſia, together with the Miſſel and Fieldfares, in prodigious quantities about Michaelmas, in their flight to the Alps.

Is ſaid to be migratory in France, viſiting Burgundy when the grapes are ripe, and doing great damage to the vineyards.

In various parts of England is known by the names of Song Thruſh, Mavis, and Greybird.

A ſpecies of Thruſh has been mentioned under the denomination of Heath Throſtle; is ſaid to have the breaſt darker and the tail ſhorter than this bird; but we have not ſufficient grounds to believe it is diſtinct from this.

The food of the Throſtle is inſects and berries of various kinds; but it is particularly fond of ſhelled ſnails, eſpecially the *Helix nemoralis*, which it breaks by reiterated ſtrokes againſt ſome ſtone. It is not uncommon to find a great quantity of fragment ſhells together, as if brought to one particular ſtone for that purpoſe.

THROSTLE-COCK. Vide *Thruſh-miſſel.*

THRUSH. A genus of birds, the characters of which are,

Bill nearly ſtrait, but bending a little towards the point, and ſlightly notched near the end of the upper mandible.

Noſtrils oval, naked.

Tongue ſlightly jagged at the end.

Mouth furniſhed with a few ſlender hairs at the corners.

Toes, the middle one connected to the outer one as far as the firſt joint.

THRUSH-GOLDEN. Vide *Oriole.*

THRUSH-HOLM. Vide *Thruſh-miſſel.*

THRUSH-MISSEL.

Turdus viſcivorus. *Lin. Syſt.* i. p. 291. 1. *Gmel. Syſt.* ii. p. 806. *Raii Syn.* p. 64. A. 1. *Will.* p. 137. t. 36. *Ind. Orn.* i. p. 326. 1.

Turdus major. *Briſ.* ii. p. 200. 1.—*Ib.* 8vo. i. p. 213.

La Draine. *Buf.* iii. p. 295. t. 19. f. 1.

Miſſel Thruſh. *Br. Zool.* i. N° 105.—*Ib.* fol. 90. t. P. f. 1. *Arct. Zool.* ii. p. 341. B. *Will. Angl.* p. 187. t. 36. *Albin.* i. t. 33. *Lewin, Br. Birds,* ii. t. 57. *Lath. Syn.* iii. p. 16. 1. *Walc. Syn.* ii. t. 197. *Pult. Cat. Dorſet.* p. 10.

PROVINCIAL.

Throſtle-cock. Screech, or Shrietch Thruſh. Holm Thruſh. Miſſeltoe Thruſh.

This is the largeſt ſpecies of Thruſh; weight near five ounces; length eleven inches.

The bill is duſky; the baſe of the lower mandible yellowiſh; irides hazel. The whole upper parts of the bird are of a light brown, a little inclined to rufous on the rump; ſides of the head and throat yellowiſh white, ſpotted with brown; from thence to the vent white; the breaſt marked with triangular ſpots, belly and ſides with roundiſh ones of a duſky colour; the two largeſt ſeries of wing coverts

covert are tipped with white; quills brown, edged with dusky on the outer web; tail the same; the three or four outer feathers tipped with white, and the inner web of the exterior one almost white; legs are of a light colour, inclining to yellow.

There is very little difference in the plumage of the sexes, but the female is not quite so bright in colour.

The Missel Thrush is by no means plentiful in England, and seems to be less so in winter. It begins to sing in January, if the weather is mild, but ceases to sing as the thermometer sinks below 40 degrees. About the middle of March it makes a nest in the fork of some tree, especially such as are covered with white moss, particularly apple-trees; frequenting orchards more than any other place in the spring; and never builds in a bush. The nest is invariably made of white moss and coarse grass, wove together with wool, and lined with fine dry grass. The eggs are four or five, and rarely six, in number, of a flesh-colour, marked with deep and light rust-coloured spots; their weight something more than two drams.

The song of this bird is much louder and superior to that of the Throstle; frequently perching upon the uppermost branch of a tall tree, it sings while the female is making her nest, and during incubation; but becomes silent as soon as the young are hatched, and is no more heard till the beginning of the new year. If the young are taken,

its

its ſong continues as before; and if the female is deſtroyed, it continues in ſong the whole ſummer. This experiment we have tried upon this and ſeveral other ſong birds, and always found it invariable. The Miſſel is a very bold bird during the breeding ſeaſon, drives all others from the neighbourhood of its neſt, and will even attack the Magpie and Jay. Its food, like the other ſpecies, is inſects and berries, particularly that of the miſſeltoe, which has been ſuppoſed neceſſary to paſs through the body of this bird to make it vegetate. That the ſeed of the berry will propagate after paſſing the organs of digeſtion, is no more wonderful than that corn ſhould grow when voided whole by a horſe. But ſuch a preparation is no more neceſſary in the one than in the other, but may be conſidered as one of the methods Nature takes to diſperſe the ſeeds of various plants.

THRUSH-SAND. Vide *Owzel-water.*
THRUSH-SONG. Vide *Throſtle.*
THRUSH-WATER. Vide *Owzel-water.*
THRUSH-WIND. Vide *Redwing.*
TIDLEY. Vide *Wren.*
TIDLEY-GOLDFINCH. Vide *Wren-gold-creſted.*
TINKERSHIRE. Vide *Guillemot-fooliſh.*
TITLING. Vide *Warbler-hedge.*

TITMOUSE. A genus of birds, the characters of which are,

Bill ſtrait, a little compreſſed, ſtrong, hard, ſharp-pointed.

Noſtrils round, and covered with reflected briſtles.

Tongue truncated, the end terminated by three or four briſtles.

Toes divided to their origin; back-toe long and ſtrong.

TITMOUSE-BEARDED.

Parus biarmicus. *Lin. Syſt.* i. p. 342. 12. *Gmel. Syſt.* ii. p. 1011. *Ind. Orn.* ii. p. 570. 23.

Parus barbatus. *Briſ.* iii. p. 567. 12.—*Ib.* 8vo. i. p. 468.

Le Meſange barbue, *ou* la Mouſtache. *Buf.* v. p. 518. t. 18.

Leaſt Butcher-bird. *Edw.* t. 55.

Bearded Titmouſe. *Br. Zool.* i. N° 167.—*Ib.* fol. 74. t. C. 2. *Arct. Zool.* ii. p. 428. H. *Albin.* i. t. 48. *Lath. Syn.* iv. p. 552. 20. —*Ib. Sup.* p. 190. *Don. Br. Birds*, i. t. 1. *Oſt. Menag.* p. 78. t. *Lewin, Br. Birds*, iii. t. 122. *Walc. Syn.* ii. t. 148.

This very elegant ſpecies is about the ſize of the Blue Titmouſe, but, on account of the length of tail, is much longer, meaſuring about ſix inches and

and a quarter. The bill is near half an inch long, of an orange colour, differing ſomewhat from the reſt of the genus, being a little arcuated; the upper mandible longeſt; irides yellow. The head is pale aſh-colour; beneath the eye is a tuft of looſe black feathers, ending in a point downwards; the hind part of the neck and back light rufous-orange; ſcapulars whitiſh; throat white; breaſt cinereous fleſh-colour; belly, ſides, and thighs, like the back, but paler; vent black; quill-feathers duſky; the inner webs of the primores white; the ſecondaries edged, and thoſe next the body tipped with the ſame colour as the back; the tail is about three inches long, and very cuneiform, nearly the colour of the back; the three outer feathers more or leſs tipped with white; legs black.

The female differs in having no black mark under the eye or at the vent; the head is light ferruginous, ſpotted with black; between the bill and eye a duſky ſpot.

The hiſtory of this ſpecies is very little known, although it breeds with us, and continues the whole year. It is found in the marſhes amongſt the reeds between Erith and London, in Glouceſterſhire, and amongſt the great reedy tracks near Cowbit in Lancaſhire. We have alſo killed it near Winchelſea in Suſſex, amongſt the reeds cloſe to the ſea-ſhore, in the month of June; there were five together, doubtleſs the brood of that year. One of the young which we procured had its neſtling

feathers, much the colour of the female; but the feathers were of a looſer texture, as in all young birds. We took much pains to find the neſt, but without ſucceſs, unleſs it is ſo like that of the Reed Wren's as not to be diſtinguiſhed, as that bird bred in the ſame place, and many ſuch neſts were taken. Authors have differed with reſpect to the ſhape and compoſition of the neſt, as well as in the place of nidification; one making it the ſhape of a purſe ſuſpended to a branch of a willow; another gives it placed on the ground amongſt ſedge of a looſe texture, compoſed of the down of the reed intermixed with narrow leaves; and that it lays four eggs of a reddiſh white, ſpotted with brown; others have undoubtedly taken the neſt of the Reed Wren for it; ſo that no certain concluſion is to be drawn from theſe various accounts.

The Bearded Titmouſe does not appear to inhabit any other places but ſuch where reeds grow in abundance, where it no doubt makes its neſt; and on the ſeeds of which, as well as inſects, it feeds; and we may attribute this deſideratum in the natural hiſtory of this bird to the inacceſſible ſwampy places they frequent.

It has been by various authors ranked with the Butcher-bird or Shrike, and was called Leaſt Butcher-bird in a former edition of the *Britiſh Zoology*, but afterwards removed by that celebrated author Mr. Pennant into this genus.

TITMOUSE-BLUE.

Parus cæruleus. *Lin. Syſt.* i. p. 341. 5. *Gmel. Syſt.* ii. p. 1008. *Raii Syn.* p. 74. A. 4. *Will.* p. 175. t. 43. *Ind. Orn.* ii. p. 566. 12. *Briſ.* iii. p. 544. 2.—*Ib.* 8vo. i. p. 462.

La Meſange bleue. *Buf.* v. p. 413.

Blue Titmouſe. *Br. Zool.* i. N° 163. t. 57. f. 2. —*Ib.* fol. 114. t. W. f. 5. *Arct. Zool.* ii. p. 427. D. *Will. Angl.* p. 242. t. 43. *Albin.* i. t. 47. *Hayes, Br. Birds,* t. 38. *Lath. Syn.* iv. p. 543. 10. *Don. Br. Birds,* t. 57. *Lewin, Br. Birds,* iii. t. 120. *Walc. Syn.* ii. t. 245. *Pult. Cat. Dorſet.* p. 10.

PROVINCIAL.

Tomtit. Nun. Hickmall.

Length about four inches and a half; weight three drams.

The bill is duſky; irides dark hazel. Forehead and cheeks white; that on the former inclines backwards, and forms a line round the crown of the head, which is of a fine blue; behind the circle of white is another of a deep blue, ſurrounding the head entirely, and joining the baſe of the under mandible, where it is nearly black; from the bill through the eye is a ſmall black line; the back is of a yellowiſh green; wings and tail blue; breaſt and belly yellow; legs lead-colour.

This bird would be much more admired for its beauty if it was leſs common. In winter it frequents houſes for the ſake of plunder; will devour fleſh greedily, whether freſh or putrid: and indeed is omnivorous. Is a conſtant attendant where horſe-fleſh is kept for hounds, as well as the farm-yard, being partial to oats, which it plucks out, and retiring to a neighbouring buſh fixes the grain between its claws, hammers it with the bill to break the huſk. In the ſummer, inſects are their chief food, in ſearch of which it plucks off a number of young buds from fruit and other trees. The neſt is always made in ſome hole either of a tree or wall, compoſed of moſs, lined with feathers and hair. The eggs are ſix or ſeven in number, rarely eight, white, ſpeckled with ruſt-colour at the larger end; their weight ſeventeen grains. It has been ſaid this bird will ſometimes lay as many as twenty eggs in the ſame neſt; but this is certainly an error, for in the great abundance of neſts we have ſeen with eggs and young, never more than eight were found. The female is tenacious of her neſt, and will often ſuffer herſelf to be taken rather than quit it, and will frequently return again after being taken out. Upon ſuch an occaſion it menaces the invader in a ſingular manner, hiſſing like a ſnake, erecting all its feathers, and uttering a noiſe like the ſpitting of a cat; and if handled bites ſeverely. It has no ſong, but makes a ſhrill note quickly repeated. It is found in every part of Europe.

TITMOUSE-COLE.

Parus ater. *Lin. Syſt.* i. p. 341. 7. *Gmel. Syſt.* ii. p. 1009. *Raii Syn.* p. 73. A. 2. *Will.* p. 175. t. 43. *Ind. Orn.* ii. p. 564. 8.

Parus atricapillus. *Briſ.* iii. p. 551. 5.—*Ib.* 8vo. i. p. 464.

La petite Charbonniere. *Buf.* v. p. 400.

Colemouſe. *Br. Zool.* i. N° 164. t. 57. f. 3.—*Ib.* fol. 114. *Arct. Zool.* ii. N° 327. *Will. Angl.* p. 241. t. 43. *Lath. Syn.* iv. p. 540. 7. *Lewin, Br. Birds,* iii. t. 180. *Walc. Syn.* ii. t. 246. *Don. Br. Birds,* iv. t. 79. *Pult. Cat. Dorſet.* p. 10.

This ſpecies weighs about two drams and a quarter; length four inches and a quarter. The bill is duſky; irides hazel. Crown of the head gloſſy black, divided on the hind part with a white ſpot; the throat and underſide of the neck black; cheeks white; back, rump, and tail, of a bluiſh grey, inclining on the rump to a buff-colour; breaſt and belly of a yellowiſh white; the ſides more yellow; quill-feathers like the tail; the coverts of the ſecondaries, and the ſmaller coverts immediately above them, are tipped with white; legs lead colour.

As this bird has been conſidered by ſome naturaliſts to be the ſame as the Marſh Titmouſe, we ſhall remark that the head of this is invariably of a gloſſy black, that of the other is of a dull ſooty

black; the black under the chin extends much lower down in this, in both ſexes, than in the other ſpecies; the white mark on the head in this is never to be found in either ſex of the other; and the tail of this is nearly a quarter of an inch ſhorter.

It muſt alſo be obſerved, that in the numerous ſpecimens we have examined no white was ever to be found on the coverts of the wings in the Marſh Titmouſe, which is conſtantly met with in this. But in the young of this ſpecies, before the white ſpot is thrown out on the head, ſome white in the wings is viſible; in which ſtate a figure is given in the folio edition of the *Britiſh Zoology* for the Marſh Titmouſe.

Mr. Willughby has well defined the diſtinction of theſe birds. Dr. Latham ſeems to have doubted the diſtinction, and appears to be the more confirmed in the opinion from an error in Sepp. who has given a figure of each as male and female.

We can, however, with certainty refute this opinion from various opportunities of attending to the neſts of both ſpecies.

The neſt of this bird is placed in ſome hole, either in a wall or a tree; is compoſed of moſs and wool, lined with hair; the eggs are ſix or ſeven in number, leſs than thoſe of the Marſh Titmouſe, of the ſame colour, white ſpotted with ruſty red; but the ſpots are ſmaller and more numerous; their weight fourteen or fifteen grains.

The Colemouſe is not ſo plentiful a ſpecies as the Marſh, keeps more in woods, and ſeems to live entirely on inſects, as we have never been able to diſcover it partaking of fleſh or grain with the other ſpecies; its note is alſo different.

TITMOUSE-CRESTED.

Parus criſtatus. *Lin. Syſt.* i. p. 340. 2. *Gmel. Syſt.* ii. p. 1005. *Raii Syn.* p. 74. 6. *Will.* p. 175. t. 43. *Ind. Orn.* ii. p. 567. 14. *Briſ.* iii. p. 558. 8.—*Ib.* 8vo. i. p. 466.

La Meſange huppé. *Buf.* v. p. 447.

Creſted Titmouſe. *Arct. Zool.* ii. p. 427. F. *Will. Angl.* p. 242. t. 43. *Albin.* ii. t. 57. *Lath. Syn.* iv. p. 545. 12. *Don. Br. Birds*, ii. t. 26. *Walc. Syn.* ii. t. 250.

The weight of this ſpecies is about two drams and a half; length four inches three quarters. The bill is duſky; irides hazel. Forehead and crown white, prettily undulated with black lines; on the hind-head the feathers are very long and black, which the bird erects in form of a conic creſt; cheeks dirty white, ſpotted with black; a black ſtroke croſſes the cheek under the eye, which turns back in an acute angle under the ear, forming the letter V; behind that is a bed of white ſurrounding the cheeks, and bounded externally by a black edge, which forms a peak on the hind-head, and alſo joins the black on the throat and chin; the

back is greeniſh brown; breaſt and belly pale buff; wings and tail much like the back, but deeper coloured; legs bluiſh lead-colour.

The Creſted Titmouſe is a ſolitary retired ſpecies, inhabits only the gloomy foreſt, particularly thoſe which abound with evergreens It has not been found in South Britain, but is not uncommon amongſt the large tracts of pines in the north of Scotland, particularly in the foreſt of Glenmoor, the property of the Duke of Gordon, from whence we have ſeen it. The eggs are ſaid to be white, with ſmall ſpots of red; the neſt we do not find to be deſcribed by any one.

Is ſaid to be found in Normandy, and in many parts between that and Sweden.

TITMOUSE-GREAT.

Parus major. *Lin. Syſt.* i. p. 341. 3. *Gmel. Syſt.* ii. p. 1006. *Raii Syn.* p. 73. A. 1. *Will.* p. 174. t. 43. *Ind. Orn.* ii. p. 562. 1. *Brif.* iii. p. 539. 1.—*Ib.* 8vo. i. p. 461.

La groſſe Meſange, *ou* Charbonniere. *Buf.* v. p. 392. t. 17.

Great Titmouſe, *or* Ox-eye. *Br. Zool.* i. N° 162. —*Ib.* fol. 113. t. W. f. 4. *Arct. Zool.* ii. p. 425. A. *Will. Angl.* p. 240. t. 43. *Albin.* i. t. 46. *Hayes, Br. Birds*, t. 18. *Lath. Syn.* iv. p. 536. 1. *Don. Br. Birds*, t. 69. *Lewin, Br. Birds*, iii. t. 117. *Walc. Syn.* ii. t. 244. *Pult. Cat. Dorſet.* p. 10.

PROVINCIAL.

Great Black-headed Tomtit. Blackcap.

This ſpecies weighs about ten drams; length five inches and three quarters. The bill is black; irides duſky. The head and throat black; cheeks white; back olive-green; the rump bluiſh grey; belly yellow, tinged with green, divided down the middle to the vent by a broad black liſt, moſt conſpicuous in the male; quill-feathers duſky; coverts bluiſh, the larger tipped with white; tail duſky; the outer feathers white on the exterior webs, the others margined with bluiſh grey; legs lead-colour.

The Great Titmouſe has all the habits of the Blue Titmouſe. The neſt is made of moſs, lined with hair, placed in the hole of a wall or tree. We once found it in the barrel of a garden pump. It lays ſometimes as many as eight eggs, but more commonly ſix, white, ſpotted with ruſt-colour, which are ſo exactly like thoſe of the Nuthatch as not to be diſtinguiſhed; their weight about thirty grains.

We have known this bird depoſit its eggs in the hole of a decayed tree upon the rotten wood, without the leaſt appearance of a neſt: probably the neſt had been deſtroyed juſt at the time the bird was compelled to lay, and had not time to make another.

The common note of this bird is a ſort of chatter,

ter, but in the ſpring it aſſumes a greater variety, a ſhrill whiſtle, and a very ſingular noiſe, ſomething like the whetting of a ſaw; but theſe ceaſe with incubation.

It is a common ſpecies in almoſt every part of England, as well as throughout Europe; ſaid alſo to inhabit the Cape of Good Hope.

TITMOUSE-LONG-TAILED.

Parus caudatus. *Lin. Syſt.* i. p. 342. 11. *Gmel. Syſt.* ii. p. 1010. *Raii Syn.* p. 74. A. 5. *Will.* p. 176. t. 43. *Ind. Orn.* ii p. 569. 20.

Parus longicaudatus. *Briſ.* iii. p. 570. 13.—*Ib.* 8vo. i. p. 469.

La Meſange à longue queue. *Buf.* v. p. 437. t 19.

Long-tailed Titmouſe. *Br. Zool.* i. N° 166. *Arct. Zool.* ii. p. 428. G. *Will. Angl.* p. 242. t. 43. *Albin.* ii. t. 57. f. 1. *Hayes, Br. Birds,* t. 33. *Lath. Syn.* iv. p. 550.—*Ib. Sup.* p. 190. *Don. Br. Birds,* t. 16. *Lewin, Br. Birds,* iii. t. 121. *Walc. Syn.* ii. t. 249. *Pult. Cat. Dorſet.* p. 10.

PROVINCIAL.

Huckmuck. Bottle-tom. Longtail-mag. Long-tail-capon. Longtail-pie. Mumruffin.

This is the ſmalleſt of the tribe; the weight about two drams; length five inches and a quarter.

The

The bill is very ſhort and black; irides hazel, edges of the eyelids yellow. On the crown of the head is a white ſtreak ſurrounded by black, which, riſing at the baſe of the bill, paſſes over each eye, and joining behind the head forms a broad liſt down the back to the rump; the reſt of the plumage above is of a purpliſh hue; the ſides of the head, throat, and under part of the neck, white, mixed with grey; from that to the vent dull purpliſh; quill-feathers black, thoſe next the body edged with grey; coverts black; the tail conſiſts of twelve feathers of unequal length; the four middle ones are wholly black, the next has a ſmall white mark on the outer web near the point, the others tipped and obliquely marked with white on the exterior webs; the ſecond feathers from the middle are the longeſt, meaſuring three inches and a half; the outer feathers on each ſide only one inch and three quarters; the legs black. In ſome the whole upper part of the neck is black, the under parts greyiſh white, except on the ſides and vent, which are of a pale vinaceous colour; and acroſs the breaſt is an obſcure duſky band.

This very elegant and ſingular ſpecies is confined chiefly to the woods and thickets, where it makes a curious oval neſt in the fork of ſome buſh or branch of a tree. In this particular it deviates from the reſt of the claſs, which invariably build in ſome hole, the Bearded and Creſted Titmice perhaps excepted; but which remains to be diſcovered. The

neſt

neſt of this bird, however, is equally well ſecured, being made of white moſs and liverwort curiouſly and firmly wove together with wool, covered at the top, with only a ſmall hole on the ſide, and lined with a prodigious quantity of feathers. This ſingular fabric is a work of time, taking four or five weeks to complete it. Low ſituations ſeem to be its delight, eſpecially about ſuch trees and hedges as are covered with white moſs and lichen, amongſt which it moſt commonly places its neſt.

The egg is leſs than any Britiſh bird, except the Golden-creſted Wren, weighing about twelve grains; colour white, ſparingly marked with ſmall ruſt-coloured ſpots towards the larger end. We are frequently told this little creature will lay upwards of twenty eggs before it ſits; but we have never been able to find more than twelve, and more frequently only nine or ten. Even this is a ſurpriſing quantity of prolific matter to be produced from ſo ſmall a body in ſo ſhort a ſpace of time as ten days, equal to the weight of the bird. To ſupply this great expenditure of animal matter, as well as the ordinary excretion, a ſupply of food conſiderably more than its own weight in that time is abſolutely neceſſary.

Its food principally conſiſts of inſects and their *larvæ*, in ſearch of which it picks off the buds from the trees. After the young are fled they keep with the parent birds till the returning ſpring; are continually in motion in ſearch of food, ſhifting their quarters

quarters daily. When in motion it is always uttering a ſmall note, by which means they keep together.

TITMOUSE-MARSH.

Parus paluſtris. *Lin. Syſt.* i. p. 341. 8. *Gmel. Syſt.* ii. p. 1009. *Raii Syn.* p. 73. A. 3. *Will.* p. 175. t. 43. *Ind. Orn.* ii. p. 563. 9. *Briſ.* iii. p. 555. 7.—*Ib.* 8vo. i. p. 465.

La Meſange de marais. *Buf.* v. p. 403.

Marſh Titmouſe, *or* Black-cap. *Er. Zool.* ii. N° 165. t. 57. f. 4.—*Ib.* fol. 114. t. W. f. 3. *Arct. Zool.* ii. p. 427. E. *Will. Angl.* p. 241. t. 43. *Albin.* iii. t. 58. f. 1. *Lath. Syn.* iv. p. 541. 8.—*Ib. Sup.* p. 189. *Lewin, Br. Birds*, iii. t. 119. *Walc. Syn.* ii. t. 247. *Pult. Cat. Dorſet.* p. 10.

PROVINCIAL.

Little-black-headed Tomtit.

The length of this ſpecies is about four inches and a half; weight two drams and a half.

The bill is duſky; irides dark hazel. Crown of the head black, but not gloſſy; cheeks of a dirty white; chin black, in ſome ſpotted with white; the back of a ruſty grey; breaſt and belly of a light browniſh buff-colour; quills and tail bluiſh grey, lighteſt on their outer margins; legs lead-colour.

We muſt here again remark, that this ſpecies is

not

not to be confounded with the Cole Titmouſe. If no other marks of diſtinction were wanting than the white on the back of the head and wing coverts, it would be ſufficient; but this is a larger bird, the colour on the back is not of that bluiſh caſt, and the tail is longer.

We ſhall, however, refer our reader to the hiſtory of that ſpecies, where we have more copiouſly defined the diſtinction.

The Marſh Titmouſe has more the habits of the Blue ſpecies, partaking with it of fleſh, and attending the oat-ricks. It appears partial to low wet ground, where old willow trees abound, in the holes of which it frequently makes its neſt. We have ſeen it artfully excavating the decayed part of that tree, carrying the chips in its bill to ſome diſtance, always working downwards, making the bottom for the reception of the neſt larger than the entrance. The neſt is compoſed of moſs and thiſtle-down, ſometimes a little wool, and lined with the down of the thiſtle. It lays five or ſix white eggs, ſpotted with ruſty red, moſtly at the larger end; their weight from nineteen to twenty-one grains.

We ſhall here remark, that all the ſpecies of Titmice, whoſe eggs are known, are ſimilar in colour, and only to be diſtinguiſhed from each other by ſize and weight. Thoſe of the Nuthatch, Creeper, Wren, Yellow Wren, Wood Wren, and Leſſer Pettychaps, all agree in their markings, and ſo like thoſe of the Titmice that it is ſcarce poſſible

to

to ſeparate them with certainty if once mixed together; and it is ſomewhat remarkable, all theſe birds breed in holes, or make a covered neſt.

This bird is not ſo common as the Great or Blue ſpecies, but more plentiful than the Cole-Mouſe. It is alſo found in many other parts of Europe, eſpecially in Sweden and Italy.

The notes of the three ſpecies with black heads are much alike, eſpecially that which is like the whetting of a ſaw, and the whiſtle made uſe of only in the ſpring; but the chatter is to be diſtinguiſhed.

TITMOUSE-WOOD. Vide *Wren-golden-creſted.*

TOMTIT. Vide *Titmouſe-blue.*

TOMTIT-BLACK-HEADED. Vide *Titmouſe-great* and *Marſh.*

TONEY-HOOP. Vide *Finch-bul.*

TOWWILLE. Vide *Sanderling.*

TREE-CLIMBER or CREEPER. Vide *Creeper* and *Sparrow-tree.*

TULIAC. Vide *Gull-ſkua.*

TURNSTONE.

Tringa Interpres. *Lin. Syſt.* i. p. 248. 4. *Gmel. Syſt.* ii. p. 671. *Ind. Orn.* ii. p. 738. 45.

Tringa Morinella. *Lin. Syſt.* i. p. 249. 6.

Morinellus marinus. *Raii Syn.* p. 112. A. 5. *Will.* p. 231. t. 58.

Arenaria. *Briſ.* v. p. 132. 1.—*Ib.* 8vo. ii. p. 246.

Arenaria cinerea. *Brif.* v. p. 137. 2. t. 11. f. 2. —*Ib.* 8vo. ii. p. 247.

Le Tournepierre. *Buf.* viii. p. 110. t. 10.

Hebridal Sandpiper. *Br. Zool.* ii. N° 200. *Arct. Zool.* ii. N° 382.

Turnſtone, *or* Sea Dotterel. *Br. Zool.* ii. N° 199. —*Ib.* fol. 125. t. E. 2. f. 2. *Will. Angl.* p. 311. *Edw.* t. 141. *Lath. Syn.* v. p. 188. 37. —*Sup.* p. 249. *Lewin, Br. Birds*, v. t. 179. *Walc. Syn.* ii. t. 153. *Pult. Cat. Dorſet.* p. 15.

This ſpecies of Sandpiper is about the ſize of a Throſtle; length nine inches and a half; weight rather more than four ounces.

The bill is black and ſtrong, about an inch in length, and turns a little upwards; irides hazel. Forehead and chin white; acroſs the breaſt a broad band of black; the fore part of the neck black, joining to that on the breaſt, and encircling the upper part of the neck; a black ſtreak from the eyes, and another from the bill, meet in an angle at the collar on the ſide of the neck; above the eye a ſtreak of white, on the ears a ſpot of the ſame; the back part of the neck is white, mixed with brown, which paſſes down each ſide of the breaſt; the crown of the head, upper part of the back, and ſcapulars, is a mixture of black and ferruginous; the latter whitiſh on their exterior edges; leſſer wing coverts nearly the ſame, but lighter; lower part of the back under the ſcapulars white; quills duſky,

duſky, the ſecondaries tipped with white; the greater coverts edged with the ſame; the ſhafts of the primary quills white; belly, vent, and under tail coverts, white; upper tail coverts white, croſſed with a bar of black; tail black, tipped with white, except the middle feathers; legs orange.

The bird from which the above deſcription is taken was killed in September on the coaſt of South Wales; another in my collection, killed in Cornwall the beginning of Auguſt, has a faint collar round the neck, which, with the band on the breaſt, is duſky brown; the chin is white, but the whole head is brown, with duſky ſtreaks; the back and ſcapulars black, each feather deeply margined with light ferruginous; the exterior feather of the tail white on the outer web; legs light yellowiſh brown. This is undoubtedly a young bird.

The Turnſtone is ſubject to great variety in reſpect to the markings about the head and neck; but the black on the breaſt, and more or leſs round the neck, at once diſtinguiſh it from any other ſpecies.

In ſome the baſe half of the tail is white, and the quills next the body the ſame, as well as the baſe of the reſt, except the four firſt. In others the lower part of the back, rump, and upper tail coverts, are wholly white.

This bird is ſometimes met with on the coaſt in ſmall flocks of five or ſix; probably the brood, which in moſt, if not in all, of this claſs conſiſts of

four young. It is not known to breed with us, but viſits ſome of our ſhores in Auguſt, and departs in the ſpring. It is ſaid to breed in Hudſon's Bay, makes a ſlight neſt on the dry ground, and lays four olive-coloured eggs, ſpotted with black. Departs from thence in September.

The name has been given it from its manner of turning up the ſtones in ſearch of marine inſects.

TURTLE. Vide *Dove-Turtle.*

TWINK. Vide *Finch-chaf.*

TWITE.

Fringilla montium. *Gmel. Syſt.* ii. p. 917. 68. *Ind. Orn.* i. p. 459. 84.

Linaria montana. *Briſ.* iii. p. 145. 33.—*Ib.* 8vo. i. p. 346. *Raii Syn.* p. 91. A. 4. *Will.* p. 191.

Mountain Linnet. *Br. Zool.* N° 133. t. 53.—*Ib.* fol. 111. *Will. Angl.* p. 261. *Arct. Zool.* ii. p. 380. C. *Lewin, Br. Birds,* ii. t. 86. *Lath. Syn.* iii. p. 307. 76. *Pult. Cat. Dorſet.* p. 13.

This ſpecies of Finch is rather larger than the Linnet or Greater Redpole; length about ſix inches. The bill is yellowiſh, brown at the tip; irides hazel. The top of the head and rump red; upper parts duſky black, edged with rufous; beneath rufous white, with blackiſh ſpots on the throat; belly white.

In ſome the whole head is light rufous, ſtreaked

on

on the top with duſky; the back, ſcapulars, and wing coverts, dark rufous-brown, with paler edges; rump tinged with red; greater quills and tail duſky black, more or leſs edged with white on the exterior webs; upper part of the breaſt and ſides rufous-white, ſpotted and ſtreaked with dark rufous-brown; lower part of the breaſt, belly, and under tail coverts, white; tail forked; legs duſky. Like the Redpole, this bird is ſubject to much variety in reſpect to the red markings. Some are deſcribed to have the top of the head, breaſt, and rump, red. In its general appearance it is much like the female of that bird, but darker on the upper parts, as well as more rufous on the cheeks and throat; the red on the rump is never found in that bird.

The Twite is gregarious, accompanying the Linnet in large flights, and is taken about London together by the birdcatchers. A variety has been given as the Mountain Linnet; it has a twittering note, but has not been obſerved to ſing.

Dr. Latham favoured me with the neſt and eggs, which he received out of Yorkſhire.

The neſt is formed of moſs and roots of plants mixed with heath, and lined with finer heath and fibrous roots. The egg is the ſize of that of the Linnet, of a blue-white, faintly ſpotted with purpliſh-red at the larger end.

The female is ſaid to want the red mark on the rump, and may therefore be frequently miſtaken

for the common Linnet before it has thrown out the other red markings.

It is poſſibly found in many other parts of England in the winter ſeaſon, but not generally diſtinguiſhed from the Linnet.

V.

VELVET-RUNNER. Vide *Rail-water.*

W.

WAGEL.

Larus nævius. *Lin. Syſt.* i. p. 225. 5. *Gmel. Syſt.* ii. p. 598. *Briſ.* vi. p. 167. 5. t. 15.—*Ib* 8vo. ii. p. 406.

Wagellus Cornubienſium. *Raii Syn.* p. 130. A. 13. *Will.* p. 266. t. 66.

Larus marinus. *Ind. Orn.* ii. p. 814. N° 6. *γ.*

Le Goéland varié, Griſard. *Buf.* viii. p. 413. t. 33.

Wagel Gull. *Br. Zool.* ii. N° 247. A. t. 88.—*Ib.* fol. ii. 422. *Arct. Zool.* ii. N° 453.—*Ib. Sup.* p. 70. *Will. Angl.* p. 349. t. 66. *Lath. Syn.* vi. p. 375. 6. *Lewin, Br. Birds,* vi. t. 209. *Walc. Syn.* i. t. 111. *Pult. Cat. Dorſet.* p. 18.

There

There is no bird which ſeems to have puzzled ornithologiſts more than this. Indeed all the Gull tribe are difficult to diſcriminate until they arrive at maturity, by reaſon of the various plumage they aſſume, and the length of time arriving to their full plumage; ſome not under three years. We have at this time a live Gull mottled brown and white, the ſize of the Herring Gull, which we have had near two years, and which is certainly that bird; and yet all the change perceived is, that it has now leſs brown on the head and neck. From theſe circumſtances no doubt the ſpecies have been multiplied; and as the Winter Gull has been proved to be no other than the Common Gull in its firſt feathers, and the Red-legged Gull the young of the Black-headed Gull; ſo we have no doubt but that the Wagel is no other than the Great Black-backed Gull. For a deſcription, therefore, we refer our readers to that bird, where the plumage of the young is deſcribed. Thoſe who wiſh to conſult other authors upon this head we refer to ſuch as we have here quoted.

WAGTAIL. A genus of birds, whoſe characters are,

Bill weak and ſlender, ſlightly notched at the end.
Tongue lacerated at the tip.
Legs ſlender.

Their tail is long, and conſtantly in motion; their pace is running, not hopping; their flight undulated.

WAGTAIL. Vide *Parre.*

WAGTAIL-GREY.

Motacilla boarula. *Lin. Mant.* 1771. p. 527. *Gmel. Syſt.* ii. p. 997. *Ind. Orn.* i. p. 502. 4.

Motacilla cinerea, an flava altera Aldr. *Raii Syn.* p. 75. 3. *Will.* p. 172. 3.

Motacilla flava. *Briſ.* iii. p. 471. 41. t. 23. f. 3. —*Ib.* 8vo. i. p. 439.

Bergeronette jaune. *Buf.* v. p. 268.

Yellow Wagtail. *Albin.* ii. t. 58.

Grey Wagtail. *Br. Zool.* i. N° 144.—*Ib.* fol. 105. *Will. Angl.* p. 238. *Edw.* t. 259. *Lath. Syn.* iv. p. 398. 4.—*Sup.* p. 178. *Don. Br. Birds*, t. 40. *Lewin, Br. Birds*, iii. t. 95. *Walc. Syn.* ii. t. 227. *Pult. Cat. Dorſet.* p. 8.

PROVINCIAL.

Winter Wagtail.

This very elegant ſpecies weighs about five drams; length ſeven inches and three quarters. Bill duſky; irides dark hazel. Crown of the head, cheeks, back of the neck, back, and ſcapulars, dark cinereous; greeniſh yellow on the rump; the eyelids, chin, throat, and breaſt, buff-coloured yellow; behind the eye a ſtroke of the ſame; belly pale yellow;

low; the wing coverts and quills black, the former bordered with light brown, the latter ſlightly edged on the exterior webs of the primores, and the three next the body deeply margined on their outer webs with yellowiſh white; the inner webs of all, except the three or four largeſt, are white at the baſe; the vent, ſides of the upper and the under tail coverts, bright yellow; the tail is four inches long, the feathers of which bend a little downwards; the outer feather is entirely white; the ſecond white, except on the outer web, which is black to within half an inch of the end; the third like the laſt, with the addition of a ſmall ſtreak of black on the edge of the interior web; the fourth and fifth black; middle ones duſky, tinged with aſh-colour, and edged with yellow towards the baſe; legs duſky brown.

This is the winter plumage. About the beginning of March ſome ſpots of black are ſeen on the throat and chin, which increaſes till thoſe parts are wholly black, except the tips of the feathers, which are ſlightly edged with white.

All authors ſeem to agree in the miſtake that the female does not poſſeſs the black mark on the throat, and that ſuch is the diſtinguiſhing mark of the male ſex; but obſerving that none had that mark till after the month of February, we were led to diſſect ſeveral before the black appeared, and afterwards, and found both ſexes with and without that mark.

The Grey Wagtail viſits us about the latter end

of September, and departs in April. At firſt not one is to be ſeen with the black throat, and in the month of March none are to be found without more or leſs black on that part; but in the female it is not ſo conſpicuous. At this ſeaſon the breaſt and belly of the male becomes of a bright yellow; the other ſex is alſo brighter on thoſe parts than in the winter; and each have a white ſtreak from the baſe of the bill paſſing down the neck on the edge of the black.

It is more than probable this and the Yellow Wagtail have been ſometimes confounded. In the *Britiſh Zoology* mention is made of that bird remaining all the year in Hampſhire; probably taken from Mr. White's *Natural Hiſtory of Selborne*, p. 38, who ſays, "Wagtails, both white and yellow, remain with us all winter." And in the *Naturaliſt's Calendar* it ſays, p. 8, "Grey and White Wagtails appear January the 2d;" again, p. 12, "Yellow Wagtails appear from January 25th to April 14th." It ſeems evident, therefore, this author did not diſcriminate the two ſpecies. In the *Linnæan Tranſactions*, p. 126, Vol. I. Mr. Markwick ſays, "The White, Grey, and Yellow Wagtails are often ſeen in the middle of winter, although the generality of them diſappear in autumn." From this account it ſhould ſeem this bird was found in thoſe parts during ſummer. We have, however, never been able to have ocular proof of this bird remaining with us in ſummer, and are inclined to believe the

confuſion

confuſion of the ſpecies has given riſe to that idea, at leaſt in the ſouth of England.

Dr. Latham, in his *Synopſis*, ſays he has been informed theſe birds breed in Cumberland, where they firſt appear in April; that young ones have been ſhot in June; and that they depart again in October. The neſt is ſaid to be made of dried fibres and moſs, lined with hair, feathers, or wool; the eggs from ſix to eight in number, of a dirty white, marked with yellow ſpots.

This ſpecies with us is ſolitary, frequenting watery places, eſpecially ſmall ſhallow ſtreams, for the ſake of inſects, on which it feeds.

Albin ſeems to have given a figure of this bird for the female Yellow Wagtail.

WAGTAIL-SPRING, or SUMMER. Vide *Wagtail-yellow.*

WAGTAIL-WATER. Vide *Wagtail-white.*

WAGTAIL-WHITE.

Motacilla alba. *Lin. Syſt.* i. p. 331. 11. *Gmel. Syſt.* ii. p. 960. *Raii Syn.* p. 75. A. 1. *Will.* p. 171. t. 42. *Ind. Orn.* ii. p. 501. 1. *Brif.* iii. p. 461. 38.—*Ib.* 8vo. i. p. 437.

La Lavandiere. *Buf.* v. p. 251. t. 14. f. 1.

White Wagtail. *Br. Zool.* i. N° 142. t. 55.—*Ib.* fol. 104. *Arct. Zool.* ii. p. 396. B. *Will. Angl.* p. 237. *Albin.* i. t. 49. *Lath. Syn.* iv. p. 395. 1.—*Sup.* p. 178. *Lewin, Br. Birds*, iii. t. 95. *Walc. Syn.* ii. t. 226. *Pult. Cat. Dorſet.* p. 8. *Don. Br. Birds*, i. t. 5.

Motacilla

Motacilla alba. *Gmel. Syſt.* i. p. 961.
Collared Wagtail. *Lath. Syn.* iv. p. 396.

PROVINCIAL.

Water-wagtail. Diſhwaſher. Waſherwoman.

The weight of this ſpecies is near ſix drams; length ſeven inches and a half.

The bill is black; irides duſky. The forehead, cheeks, and ſide of the neck, white; the back of the head, upper and under ſide of the neck, chin, and breaſt, black; back duſky, daſhed with cinereous, in ſome nearly black; lower part of the breaſt and belly white; quill-feathers duſky, two or three next the body excepted, which are black, deeply bordered on their exterior webs with white; the greater coverts black, tipped with white; the others black, daſhed with aſh-colour; the tail-feathers are black, except the two outmoſt on each ſide, which are black at the baſe, the other part white.

The female is duſky where the male is black, with more cinereous on the back and rump.

This is the ſummer plumage; in the autumn the black feathers on the chin and throat fall off, and are replaced by white ones, leaving only a black patch, ſomewhat in form of a creſcent, on the breaſt.

The young birds have no black on the throat till the returning ſpring. In this ſtate it has been deſcribed

deſcribed as a variety, but is in fact the conſtant winter plumage, regaining this black mark about the month of March.

The White Wagtail is a very active bird, and continually in motion, running after flies. In winter they change their abode, but do not quit the kingdom. As the weather becomes ſevere they haunt marſhes ſubject to the flow of the tide. In ſuch places on the coaſt we have ſeen them in abundance when none were to be found inland.

Early in the ſpring they return to their uſual ſummer ſituations, and from the number that are ſometimes ſeen together at this time attending ſheepfolds and new-ploughed fields, it ſhould ſeem they were gregarious in their flights. In the breeding ſeaſon they ſeem to prefer pleaſure-grounds that are conſtantly mowed, on which they run unincumbered, and where the inſects have not ſufficient cover to evade their ſight.

The neſt is found in various places; ſometimes on the ground in a heap of ſtones, the hole of a wall, or on the top of a pollard-tree. It is compoſed of moſs, dried graſs and fibres, put together with wool, and lined with feathers or hair. The eggs are four or five in number, white, ſpotted all over with light brown and aſh-colour, weighing about forty grains. They exactly reſemble that of the Cuckoo, which bird frequently makes choice of its neſt to depoſit her egg in. It ſings very prettily early in the ſpring, and frequently gives the

alarm

alarm on the appearance of a Hawk, which it purſues in company with the Swallows.

WAGTAIL-WINTER. Vide *Wagtail-grey.*

WAGTAIL-YELLOW.

Motacilla flava. *Lin. Syſt.* i. p. 331. 12. *Gmel. Syſt.* ii. p. 963. *Raii Syn.* p. 75. A. 2. *Will.* p. 172. t. 68. *Ind. Orn.* ii. p. 504. 8.

Motacilla verna. *Briſ.* iii. p. 468.—*Ib.* 8vo. i. p. 438.

Bergeronette de Printemps. *Buf.* v. p. 265. t. 14. f. 1.

Yellow Wagtail. *Br. Zool.* i. N° 143.—*Ib.* fol. 105. *Arct. Zool.* ii. p. 396. F. *Will. Angl.* p. 238. t. 68. *Edw.* t. 258. *Lath. Syn.* iv. p. 400. 6.—*Ib. Sup.* p. 179. *Lewin, Br. Birds,* iii. t. 97. *Walc. Syn.* ii. t. 228. *Don. Br. Birds,* i. t. 15. *Pult. Cat. Dorſet.* p. 8.

PROVINCIAL.

Spring, or Summer Wagtail.

Weight about five drams; length ſix inches and a half.

The bill is black; irides hazel. The upper part of the head and back of the neck pale olive-green; back, ſcapulars, rump, and wing coverts, of the ſame colour, but darker: the whole under parts, from chin to vent, full bright yellow; the coverts of the ears like the upper part of the head; over the

the eye a yellow ſtreak; quill-feathers duſky, thoſe next the body, and the greater coverts, edged with yellowiſh white; tail duſky, the two middle feathers daſhed with olive, two outer feathers on each ſide white full half way from the end of the ſecond feather, and running obliquely upwards leaves the outer web of the firſt entirely white; legs black; hind claw very long, and but little curved.

The female is leſs bright in colour, the yellow underneath in ſome appearing almoſt white at a little diſtance.

There appears no doubt but many authors have confounded this ſpecies with the Grey Wagtail, which we have remarked more fully in the hiſtory of that bird.

The male of this ſpecies, it is ſaid, poſſeſſes a few black ſpots on the throat; but ſuch a mark we have never obſerved in more than a hundred ſpecimens. It muſt therefore be rare, if not a miſtake in deſcribing the Grey Wagtail for this. If no other mark of diſtinction were to be found but the length and ſtraitneſs of the hind claw in this, it would be ſufficient to know it from the Grey Wagtail, which is very ſhort and crooked. The tail of this bird is alſo an inch ſhorter, and has only two feathers on each ſide, partly white. The under parts of the male are of a much fuller yellow, and the upper parts never poſſeſs any of the cinereous colour.

The Yellow Wagtail viſits us about the time the other departs, and migrates again in September.

It

It frequents arable land, eſpecially in the more champaign parts; ſometimes uncultivated ground interſperſed with furze; is alſo partial to bean fields: in all ſuch places it breeds, and does not ſeem to regard water ſo much as either the other ſpecies. The neſt is always placed on the ground, compoſed of dried ſtalks and fibres, lined with hair. The eggs are four or five in number, not very unlike thoſe of the Sedge Warbler, of a pale brown, ſprinkled all over with a darker ſhade, in ſome very obſcurely, weighing about twenty-ſeven grains. It has all the actions and notes of the other ſpecies; the cry is more ſhrill than the White, and leſs ſo than the Grey Wagtail, but does not ſeem to have much of a ſong.

Said to be found in Siberia and Ruſſia in ſummer, and to continue in France the whole year.

WARBLER. A genus of birds, the characters of which are,

Bill ſlender and weak.
Noſtrils a little depreſſed and ſmall.
Tongue cloven.
Toes, the extreme one joined at the under part to the middle one at the baſe.

WARBLER-DARTFORD.

Motacilla provincialis. *Gmel. Syſt.* ii. p. 958. 67.
Sylvia dartfordienſis. *Ind. Orn.* ii. p. 517. 31.
Le Pitchou de Provence. *Buf.* v. p. 158.

Dartford Warbler. *Br. Zool.* i. N° 161. t. 56. *Lath. Syn.* iv. p. 437. 27.—*Ib. Sup.* p. 181. *Don. Br. Birds*, t. 10. *Lewin, Br. Birds*, iii. t. 106. *Walc. Syn.* ii. t. 237.

This ſpecies is rather larger than the common Wren, and much longer by reaſon of the tail, which is one half its length nearly; the weight is about two drams and a half; length five inches and a half.

Bill black, at the baſe of the upper mandible whitiſh; the upper mandible a little curved; irides and eyelids yellow. The whole upper parts are of a duſky brown; cheeks dark cinereous; throat, neck, and breaſt, fine deep ferruginous; ſides the ſame, but not ſo bright; middle of the belly white; quills duſky, ſlightly edged with dark cinereous on the outer webs, thoſe next the body and coverts with dark ferruginous brown; at the bend of the wing, under the *alula ſpuriaræ*, is a ſpot of white; the tail is conſiderably cuneiform; the outer feather is tipped with white, and edged with the ſame on the exterior web; the next ſlightly tipped with white; the remainder of theſe and all the others duſky; the middle ones edged with cinereous; legs yellowiſh.

In ſome the throat is ſpeckled with white.

The female and young birds are of a lighter colour and more rufous.

The Dartford Warbler, ſo called from having been

been firſt diſcovered in England near that place; it is, however, a ſcarce ſpecies, rarely noticed in this country Dr. Latham ſeems to have been the firſt diſcoverer of this bird in England, and communicated it to Mr. Pennant, who firſt publiſhed it in his *Britiſh Zoology*, a pair having been killed on Bexley Heath, near Dartford, on the 10th of April 1773. Since that time, Dr. Latham informs us, ſeveral were ſhot in the winter of 1783 on a common near Wandſworth in Surrey, now in the Leverean Muſeum; from which circumſtance that author very juſtly obſerves, that if it is found here only as a winter migrant, he cannot reconcile the circumſtance of its breeding in France (which has been ſaid to be the caſe), as all migratory birds go northward to breed, not to a warmer climate. In the month of September 1796 we obſerved many of theſe birds about Falmouth in Cornwall frequenting the furzy hills, and killed ſeveral from that time to the twenty-fourth of December, when a ſudden fall of ſnow, that covered the ground for ſome time, drove them from that part. Many of theſe birds, on their firſt appearance, were in their neſtling feathers, from which ſome hopes were entertained of their breeding in thoſe parts; but with the moſt diligent ſearch not one was to be found the following ſummer; nor indeed did they ever return after the ſnow had driven them away. Where they could go to from that moſt ſouthern part of England, unleſs they croſſed the ſea (which

is very improbable), is impoſſible to gueſs, except they ſought ſome more ſheltered ſituation further in the peninſula of that country towards the land's-end; and even in that caſe it was natural to expect them in their return northward; for undoubtedly they muſt breed in ſome more northern parts.

As yet the neſt and eggs are unknown.

It is a ſhy bird, concealing itſelf amongſt the thickeſt furze on the leaſt alarm, and creeping from buſh to buſh; is ſaid to perch on the top of a buſh in the ſpring, and from thence are ſeen darting into the air after flies, and returning again to the ſame ſpot. Their food is ſmall inſects of all kinds; in the ſtomach of one we diſſected was the *larvæ* of a large Cimex.

The ſhortneſs of the wing and length of tail give it a ſingular manner of flying, which is in ſhort jerks, with the tail thrown up. The note is a weak but ſhrill piping noiſe ſeveral times repeated.

WARBLER-GRASSHOPPER.

Alauda trivialis. *Gmel. Syſt.* p. 796. 5.
Sylvia Locuſtella. *Ind. Orn.* ii. p. 515. 25.
Locuſtella avicula. *Raii Syn.* p. 70. A. 7. *Will.* p. 151.
La Locuſtelle. *Buf.* v. p. 42.
Titlark that ſings like a Graſshopper. *Will. Angl.* p. 207.
Graſshopper Lark, Warbler. *Br. Zool.* i. N° 156.

—*Ib.* fol. 95. t. Q f. 5. *Arct. Zool.* ii. p. 419. L. *White, Hiſt. Selb.* p 45.

Graſshopper Warbler. *Lath. Syn.* iv. p. 429. 20. *Lewin, Br. Birds,* iii. t. 98.

This ſpecies is leſs than the Whitethroat; length five inches and a half; weight about three drams and a quarter. The bill is duſky above, whitiſh beneath; irides light hazel. The whole upper parts of the bird are olivaceous-brown; the middle of each feather duſky, except on the back of the neck, which gives it a pretty ſpotted appearance; eyelids, chin, throat, and belly, yellowiſh white; breaſt, ſides, and thighs, inclining to brown, the two laſt faintly ſtreaked with duſky; under tail coverts very pale brown, marked down the ſhafts with long pointed ſtreaks of a duſky colour; quills and tail duſky brown, lighter on their exterior edges, tinged with olive; the tail is much cuneiform, and the feathers ſomewhat pointed; legs very pale brown; claws light horn-colour; hind claw ſhort and crooked.

In ſhape the Graſshopper Warbler very much reſembles the Sedge Warbler; is rather inferior in ſize, and at once diſtinguiſhed by its ſpotted back.

It is not a plentiful ſpecies, but probably appears leſs ſo by its habit of concealing itſelf amongſt furze and thick hedges, diſcovering their place of concealment only by their ſingular Cricket-like note, which

which is ſo exactly like that of the Grilla-Talpa as ſcarce to be diſtinguiſhed.

We have found it in Hampſhire, in South Wales, and in Ireland, but no where ſo plentiful as on Malmſbury Common in Wiltſhire, to which place the males come about the ſecond week in April. At this time only they expoſe themſelves upon the top branches of the furze, and are continually making their ſingular chirping notes, their only ſong. In this ſituation we have killed ſeveral.

As ſoon as the females arrive, which is in about ten days after, the males no longer expoſe themſelves, and are almoſt ſilent till about the duſk of the evening, when they are inceſſantly crying; poſſibly to decoy the larger ſpecies of Graſshoppers, or the Grilla-Talpa, which begin their chirping with the ſetting ſun. The female very much reſembles the other ſex; is ſo ſhy as to be obtained with difficulty. On the eighteenth of May we found the neſt of this bird in a patch of thick brambles and furze, with two eggs; but as they had not been incubated, probably more would have been laid. The neſt is of a flimſy texture, like that of the Whitethroat, compoſed of dried ſtalks and gooſegraſs, lined with fibrous roots. The eggs are of a ſpotleſs bluiſh white, weighing twenty-one grains. From the ſcarcity of the bird, and the artful manner in which it conceals its neſt, it is rarely found; nor has any author noticed it.

WARBLER-HEDGE.

Motacilla modularis. *Lin. Syſt.* i. p. 329. 3. *Gmel. Syſt.* ii. p. 952. *Ind. Orn.* ii. p. 511. 13.

Curruca Eliotæ. *Raii Syn.* p. 79. A. 6. *Will.* p. 157.

Magnanima Aldr. *Will.* p. 157.

Corruca ſepiaria. *Briſ.* iii. p. 394. 12.—*Ib.* 8vo. p. 420.

Traine-buiſſon Mouchet, *ou* Fauvette d'hyver. *Buf.* v. p. 151.

Hedge Sparrow. *Br. Zool.* i. N° 150.—*Ib.* fol. t. S. 1. f. 3. 4. *Arct. Zool.* ii. p. 418. H. *Will. Angl.* p. 215. *Albin.* iii. t. 59.—*Ib. Song Birds*, t. p. 81. *Lath. Syn.* iv. p. 419. 9. *Lewin, Br. Birds*, iii. t. 102. *Walc. Syn.* ii. t. 232. *Pult. Cat. Dorſet.* p. 9.

PROVINCIAL.

Titling. Dunnock.

This well-known ſpecies, commonly called Hedge Sparrow, needs little deſcription. The length is five inches and three quarters; weight near ſix drams.

Bill duſky; irides light hazel. Head and neck brown, mixed with aſh-colour; back and wing coverts darker brown, edged with rufous-brown; throat and breaſt dull aſh-colour; belly dirty white; ſides and vent tawny-brown.

The

The female has leſs aſh-colour about the head and breaſt.

The Hedge Warbler is found in all parts of England; has a pleaſing ſong, which it begins with the new year, if the weather is mild; breeds early, making a neſt in March, compoſed of green moſs and wool, and lined with hair, which is placed in ſome low evergreen ſhrub, thick bruſh, or cut hedge; frequently builds in faggot piles. The eggs are four or five in number, blue; their weight about twenty-eight grains.

This bird is one of the few of the Warbler tribe that remains with us the whole year. Its food is inſects and worms, but, like the Redbreaſt, will in defect of theſe pick up crumbs of bread; and ſeems to prefer ſituations near the habitation of man.

The Cuckoo frequently makes choice of this bird's neſt for the purpoſe of depoſiting its egg.

Is ſaid to be migratory in France, leaving that country in ſpring, a very few excepted.

WARBLER-SEDGE.

Motacilla ſalicaria. *Lin. Syſt.* i. p. 330. 8. *Gmel. Syſt.* ii. p 955.

Sylvia ſalicaria. *Ind. Orn.* ii. p. 516. 26.

Curruca arundinacia. *Briſ.* iii. p. 378. 5.—*Ib.* 8vo. i. p. 415.

Avis conſimilis Stoparolæ, et Magnanimæ. *Raii Syn.* p. 81. 6. *Will.* p. 153. *Will. Angl.* p. 217.

Salicaria. *Raii Syn.* p. 81. 11. *Will.* p. 158.

La Fauvette de roſeaux. *Buf.* v. p. 142.
Willow Lark, *or* Sedge Bird. *Br. Zool.* i. N° 155.
Sedge Warbler. *Arct. Zool.* ii. p. 419. M. *White, Hiſt. Selb.* pp. 67. 71. 74. *Lewin* iii. t. 62. *Lath. Syn.* iv. p. 432. 21.—*Sup.* p. 150. *Lewin, Br. Birds*, iii. t. 105. *Walc. Syn.* ii. t. 236. *Don. Br. Birds*, ii. t. 48. *Pult. Cat. Dorſet.* p. 9.

PROVINCIAL.

Sedge-Wren. Leſſer-reed-Sparrow.

The weight of this ſpecies is about three drams; length five inches and a half. Bill duſky above, whitiſh beneath; irides hazel. Crown of the head and whole upper parts of a yellowiſh brown, plain on the back and ſides of the neck, rump, and upper tail coverts; the reſt obſcurely marked with duſky; the coverts of the wings more duſky, edged with olivaceous brown; quills the ſame, but ſlightly edged; over the eye a whitiſh ſtroke; all the under parts, from the chin to the under tail coverts, yellowiſh white, darkeſt on the breaſt and ſides; tail like the quills, a little cuneiform, which, when ſpread, gives it a rounded ſhape; legs duſky.

The Sedge Warbler comes to us about the middle of April, and leaves us again in September. It has a variety of notes, which it delivers in a hurried manner, and which partakes of that of the Sky Lark and the Swallow, as well as the chatter of the

the Houſe Sparrow. Is frequent by the ſides of rivers and watery places, where ſedge and reeds grow, amongſt which it makes a neſt compoſed of a little moſs intermixed with dried ſtalks, and lined with dried graſs, occaſionally a few hairs; ſometimes it is faſtened between two or three reeds; others we have found in a tuft of ruſhes on the ground, or very near it, faſtened round the bottom of them; at other times in a low buſh, or on the tump of a willow. The eggs are five or ſix in number, of a light brown colour, mottled with darker ſhades of the ſame; their weight from twenty-four to twenty-eight grains.

The ſong of this bird has been erroneouſly given to the Reed Bunting by various authors, whereas that bird has no notes that deſerve the name of ſong; but as they frequent the ſame places to breed, and that is conſpicuous on the upper branches, while this little warbler, concealed in the thickeſt part, is heard aloud, the ſong has been confounded. It has been juſtly remarked, that, if it is ſilent, a ſtone thrown into the buſh will ſet it ſinging inſtantly; it will alſo ſing of a moon-light night.

The ſimilitude in colour, ſize, manners, and habits, of this and the Reed Wren is ſo great that they have been moſt times confounded; but on compariſon that bird will be found not to poſſeſs the broad white ſtreak over the eye, nor any of the feathers on the back and wing coverts duſky. The neſt and eggs are alſo eſſentially different, as may

be found by comparing the deſcriptions. This is alſo a much more plentiful ſpecies, at leaſt is not ſo local; is to be met with in moſt parts of England, whereas the other is confined to certain tracts, eſpecially to ſuch where quantities of reeds grow. They are, however, frequently found together, for wherever the Reed Wren is the Sedge Warbler alſo inhabits; but the reverſe is not the caſe.

WASHER-WOMAN. Vide *Wagtail-white.*

WATER-CRAKE. Vide *Ouzel-water.*

WATER-CRAW. Vide *Ouzel-water.*

WATER-HEN. Vide *Gallinule.*

WATER-HEN-SPOTTED. Vide *Gallinule-ſpotted.*

WAXEN-CHATTERER. Vide *Chatterer-Bohemian.*

WHEATEAR.

Motacilla Œnanthe. *Lin. Syſt.* i. p. 332. 15. *Gmel. Syſt.* ii. p. 966. *Raii Syn.* p. 75. A. 1. *Will.* p. 168. t. 41.

Sylvia Œnanthe. *Ind. Orn.* ii. p. 529. 79.

Vitiflora. *Briſ.* iii. p. 449. 33.—*Ib.* 8vo. i. p. 434.

Le Cul blanc, Vitrec, *ou* Motteux. *Buf.* v. p. 237.

Wheat-ear. *Br. Zool.* i. N° 157.—*Ib.* fol. 102. t. S. 1. f. 5. 6. *Arct. Zool.* ii. p. 420. P. *Will. Angl.* p. 133. t. 41. *Albin.* i. t. 55. (M.) 3. t. 54. (F.) *Edw.* pref. p. 12. *Lath. Syn.* iv. p. 465.

WHE

p. 465. 75.—*Sup.* p. 182. *Lewin, Br. Birds*, iii. t. 110. *Walc. Syn.* ii. t. 241. *Pult. Cat. Dorſet.* p. 9.

PROVINCIAL.

Fallow-Finch, *or* Fallow-Smich. White tail. Snorter.

This ſpecies weighs about ſix drams and a half; length near ſix inches and a half. The bill is black, and conſiderably broad at the baſe, where it is beſet with briſtles; irides hazel. From the noſtril a black ſtreak through the eye, taking in the coverts of the ear; over each eye a white ſtroke meeting on the forehead; the upper part of the head and back cinereous-grey; rump, upper and under tail coverts, white; quill-feathers duſky, moſtly edged with light ruſt-colour; coverts black, tipped with ruſty yellow; under part of the neck buff; breaſt and belly yellowiſh white; tail white, the two middle feathers black at the end for about an inch, the reſt tipped half an inch with the ſame; legs and claws black.

The female is about a dram heavier than the male; has all the markings of that ſex, but leſs vivid; the white on the forehead and over the eye, as well as the black ſtreak, is very obſcure; and the cinereous grey on the back is mixed with brown.

The Wheatear is a migrative ſpecies, appearing with

with us the latter end of March; ſome few probably remain the whole year, as we have now and then ſeen it in the month of February. It principally frequents rabbit-warrens, or ſuch parts as are encloſed with ſtone-walls; makes its neſt in a deſerted rabbit-burrow, or in an old ſtone quarry; ſometimes in a heap of ſtones, or the hole in a wall; but moſt times on the ground. It is compoſed of moſs and dried ſtalks and fibres put together with wool, and lined with hair or wool. The eggs are five or ſix in number, of a uniform pale blue colour, weighing about forty-three grains. The number that breed in this country muſt be very conſiderable, but ſo diſperſed that few are ſeen at that ſeaſon in the ſame ſituation. In September they begin to retire, and ſeem to aſſemble from all parts to the Suſſex and Dorſet downs contiguous to the coaſt, preparatory to their departure. The quantity taken annually about Eaſtbourn is prodigious; Mr. Pennant ſays 1840 dozen. Theſe are caught in a ſingular manner, by placing two turfs on edge; at each end a ſmall horſe-hair nooſe is fixed to a ſtick, which the bird, either in ſearch of food or to evade a ſtorm of rain, attempts to get under, and is caught. Upon inquiry of the ſhepherds, whoſe trade it is, we have been informed fifty or ſixty of theſe traps have had a bird in them of a morning; ſometimes ſeveral mornings together, and then for a day or two ſcarce one is to be ſeen; and yet they are never obſerved to come in flocks;

flocks; and it is the general opinion that they come in the night.

Theſe birds uſually ſell for a ſhilling a dozen; and it is a common cuſtom in thoſe parts where they are taken to viſit the traps, take the bird out, and leave a penny in each as a reward for the ſhepherd. It is eſteemed a great delicacy, not much inferior to the Ortolan; are ſometimes ſent to the London poulterers ready picked.

This bird ſings very prettily, and not unfrequently on wing, hovering over the female; in the courting ſeaſon diſplays its tail in a ſingular manner.

A variety has been deſcribed by ſome authors to have a mixture of whitiſh and fulvous on the upper parts, and very ſmall grey ſpots on the lower part of the neck; and the two middle feathers of the tail wholly black. Such is called by Mr. Pennant, in his Appendix to the *Britiſh Zoology*, Grey Wheat-Ear.

WHEEL-BIRD. Vide *Goatſucker*.

WHEWER. Vide *Widgeon*.

WHILK. Vide *Scoter*.

WHIM. Vide *Widgeon*.

WHIMBREL.

Scolopax Phœopus. *Lin. Syſt.* i. p. 243. 4. *Gmel. Syſt.* ii. p. 657.

Numenius Phœopus. *Ind. Orn.* ii. p. 711. 6.

Numenius minor. *Briſ.* v. p. 317. t. 27. f. 1.

Arquata

Arquata minor. *Raii Syn.* p. 103. A. 2. *Will.* p. 217.

Corlieu, *ou* petit Courlis. *Buf.* viii. p. 27.

Whimbrel. *Br. Zool.* ii. N° 177.—*Ib.* fol. 119. *Arct. Zool.* ii. p. 462. B. *Will. Angl.* p. 294. *Edw.* t. 307. *Lath. Syn.* v. p. 123. *Lewin, Br. Birds*, iv. t. 154. *Walc. Syn.* ii. t. 134. *Don. Br. Birds*, iii. t. 72.

PROVINCIAL.

Curlew-knot. Curlew-Jack. Half-Curlew. Stone-Curlew.

The weight of this ſpecies of Curlew is about fourteen ounces; length eighteen inches. The bill is above three inches in length, arcuated; upper mandible duſky, under part whitiſh at the baſe, in ſome of a reddiſh fleſh-colour. The head, neck, and breaſt, pale brown down the middle of each feather, margined with white, lighteſt on the forehead, and darkeſt on the crown of the head; chin and belly white; the irides are duſky, eyelids white; ſides of the body barred with duſky; the upper parts of the body, ſcapulars, and wing coverts, duſky, margined with pale brown; quill-feathers duſky; the borders of the internal webs barred with white; the tail is duſky aſh-colour, the middle feathers darkeſt, marked with ſix or ſeven duſky bars; legs duſky.

The Whimbrel has all the manners of the Curlew,

lew, and indeed is ſo very like in plumage, that in ſome parts it has obtained the name of Jack-Curlew, from a ſuppoſition that it is the male of that bird; but it is by no means ſo plentiful a ſpecies.

It is a migrative bird, viſiting our coaſts in Auguſt, and continuing the winter, keeping together in ſmall flocks of five or ſix; has been ſuſpected to breed on the coaſt of Suſſex and Kent, eſpecially about Romney Marſh; but that has not been aſcertained with certainty.

WHIN-CHAT.

Motacilla Rubetra. *Lin. Syſt.* i. p. 332. 16. *Gmel. Syſt.* ii. p. 967. *Ind. Orn.* ii. p. 525. 58.

Rubetra major, ſive Rubicola. *Briſ.* iii. p. 432. 26. t. 24. f. 1.—*Ib.* 8vo. i. p. 430.

Œnanthe ſecunda. *Raii Syn.* p. 76. A. 3. *Will.* p. 234.

Grand Traquet, *ou* Tarier. *Buf.* v. p. 224.

Whin-Chat. *Br. Zool.* i. N° 158.—*Ib.* fol. 103. t. S. 2. f. 3. 4. *Will. Angl* p. 234. *Hayes, Br. Birds*, t. 39. *Lath. Syn.* iv. p. 454. 54. *Don. Br. Birds*, t. 60. *Lewin, Br. Birds*, iii. t. 109. *Waic. Syn.* ii. t. 140. *Pult. Cat. Dorſet.* p. 9.

PROVINCIAL.

Furze-Chat.

This

WHITE-GAME. Vide *Ptarmigan.*

WHITE-NUN. Vide *Smew.*

WHITE-TAIL. Vide *Wheatear.*

WHITE-THROAT.

Motacilla Sylvia. *Lin. Syſt.* i. p. 330. 9? *Gmel. Syſt.* ii. p. 956.

Sylvia cinerea. *Ind. Orn.* ii. p. 514. 23.

Parus cinereus. *Briſ.* iii. p. 549. 4.—*Ib.* 8vo. i. p. 463. *Buf.* v. p. 409. iii.

Ficedulæ affinis, an Spipola prima. *Raii Syn.* p. 77. A. 6. *Will.* p. 171.

Firſt Spipola of Aldrovandus. *Will. Angl.* p. 210.

VAR.

Currucca cinerea ſive cineraria. *Briſ.* iii. p. 376. t. 21. f. 1?—*Ib.* 8vo. i. p. 415.

Spipola Aldr. *Raii Syn.* p. 77. A. 1? *Will.* p. 153. —*Ib. Angl.* p. 210.

Fauvette griſe, *ou* la Griſette. *Buf.* v. p. 132.

White-Throat. *Br. Zool.* i. N° 160.—*Ib.* fol. 104. t. S. f. 4. *Arct. Zool.* ii. p. 422. S. *Albin.* iii. t. 58. *Lewin, Br. Birds,* iii. t. 104. *Hiſt. Selb.* p. 103. *Walc. Syn.* ii. t. 235. *Lath. Syn.* iv. p. 428. 19. *Pult. Cat. Dorſet.* p. 9.

PROVINCIAL.

Nettle-creeper.

This ſpecies of Warbler weighs about four drams; length five inches and three quarters. The

bill duſky brown above, whitiſh beneath; irides yellowiſh. The whole upper parts, from head to tail, cinereous-brown; coverts of the wings darkeſt, bordered with brown, inclining to rufous; quills duſky, ſlightly edged with cinereous-brown; under parts, from chin to tail, greyiſh white, darkeſt on the breaſt and thighs; in ſome the breaſt has a roſy tinge; tail like the quills; outer feather white, except at the baſe of the inner web; legs pale brown.

The female is like the male.

This is a very common ſpecies, viſits all parts of the kingdom which are incloſed about the middle of April, and conſtantly enlivens our hedges with its ſong, at which time it erects the feathers on the crown of the head. The neſt is made of gooſe-graſs lined with fibres, and ſometimes a few long hairs, but is of ſo flimſy a texture that it can afford no warmth to the eggs or young: this is generally placed in ſome low buſh amongſt nettles or other luxuriant herbs.

The eggs are four or five in number, of a greeniſh white, ſpeckled all over with light brown or aſh-colour; weight about thirty grains.

It is proper in this place to remark, that we have more than once killed a bird ſomewhat reſembling the above; weight and length the ſame; irides not ſo yellow; the whole upper parts rufous-brown; the coverts of the wings and quills next the body deeply margined with bright rufous; throat and belly ſilvery white; breaſt inclining to brown,

darker on the ſides; outer feathers of the tail like the above. In Latham's *Synopſis* a variety of the White-throat is mentioned, as deſcribed by Buffon; but that author conſidered it a different ſpecies: a plate of it is alſo given in the *Planches Enluminées*. We confeſs this bird has puzzled us not a little, more eſpecially as the neſt from which we ſhot one of them contained four white eggs, except a few grey ſpecks at the point of the larger end; the ſhape nearly round; their weight rather more than thoſe given above. The neſt was the ſame; it was placed very low in a hedge of white-thorn; another was found on bean-ſtalks, on which the male was killed. Poſſibly this may prove hereafter to be a diſtinct ſpecies, and the ſame as Dr. Latham ſpeaks of being found in Provence.

The common White-throat feeds on inſects and berries, frequenting our gardens in the ſummer for the ſake of cherries and currants. The birds here given have, it ſhould ſeem, been made two diſtinct ſpecies by ſome authors, while others have made ſeveral varieties of the White-throat; ſo that we are not a little at a loſs in reſpect to the quotations, ſome of which we give with doubts.

WHITE-THROAT-LESSER.

Sylvia Sylviella. *Ind. Orn.* ii. p. 515. 24.

Leſſer Whitethroat. *Lath. Syn. Sup.* p. 185. t. 113. *Don. Br. Birds*, iv. t. 86.

This ſpecies of Warbler was firſt noticed b

Mr. Lightfoot in Buckinghamſhire, and communicated to Dr. Latham, who firſt gave it to the world as a Britiſh ſpecies in the Supplement to his *Synopſis*.

It is leſs than the Common White-throat; length five inches and a quarter; weight about three drams and a quarter.

The bill is duſky; irides yellowiſh, with a daſh of pearl-colour. The upper part of the head, taking in the eyes, is dark aſh-colour; all the other parts above cinereous-brown; quills and tail duſky, edged with aſh-colour; from throat to vent, including the under tail coverts, ſilvery white; the exterior feather of the tail whitiſh almoſt to the baſe; the outer web quite white; legs duſky lead-colour.

This and the preceding ſpecies have doubtleſs been confounded; nor is it eaſy to determine which of them is the *Sylvia* of Linnæus. We obſerved the arrival of this bird for ſeveral years together in Wiltſhire to be from the twenty-firſt of April to the tenth of May. It is not uncommon in the north of that county, and is eaſily diſcovered by its ſhrill note, which is ſcarcely to be called a ſong, as it is only a repetition of the ſame whiſtling note ſeveral times in a hurried manner; beſides which it has a ſoft pleaſing ſong not to be heard unleſs very near. It conceals itſelf in the thickeſt hedges, and when the foliage is complete is very difficult to be ſhot. In ſuch ſituations the neſt is

placed not very diſtant from the ground, compoſed of *goſſe-graſs*, neatly but ſlightly put together, with a ſmall quantity of wool, very much like that of the Common White-throat. The eggs are four or five in number, of a bluiſh white, ſpeckled with brown and aſh-colour at the larger end, and ſometimes a few diſtant ſpots all over; their weight about twenty-five grains.

The great diſtinction between this and the Common White-throat and its varieties is, that this is inferior in ſize; the bill is ſhorter, the under as well as upper mandible is duſky; the legs darker; the whole under parts of the plumage much whiter; and the upper parts do not poſſeſs the leaſt appearance of ruſous-brown, which in the other is more or leſs invariably found, eſpecially on the wing coverts.

The Leſſer White-throat does not appear to be a plentiful ſpecies in this country, and is confined to the eaſtern parts of the kingdom, from Glouceſterſhire and Wiltſhire, in both which counties we have found them, and is probably in part of Somerſetſhire, but not in Devonſhire or Cornwall.

WHITWALL. Vide *Oriole and Woodpecker-great-ſpotted.*

WIERANGLE. Vide *Shrike-cinereous.*

WIGEON.

Anas Penelope. *Lin. Syſt.* p. 202. 27. *Gmel. Syſt.* ii. p. 527. *Raii Syn.* p. 146, A. 3. *Will.* p. 288. t. 72. *Ind. Orn.* ii. p. 860. 71.

Anas

Anas fiſtularis. *Briſ.* vi. p. 391. 21. t. 35. f. 2.
—*Ib.* 8vo. ii. p. 464.
Canard ſiffleur. *Buf.* ix. p. 169. t. 10. 11.
Wigeon, Whewer, Whim. *Br. Zool.* ii. N° 268.
—*Ib.* fol. 157. Addend. t. Q. 157. *Arct. Zool.* ii. p. 574. K. *Will. Angl.* p. 375. t. 72. *Albin.* ii. t. 99. *Lath. Syn.* vi. p. 518. 63. *Lewin, Br. Birds*, vii. t. 251. *Walc. Syn.* i. t. 71. *Pult. Cat. Dorſet.* p. 21.

This ſpecies of Duck weighs about twenty-four ounces; length twenty inches. The bill is narrow, of a bluiſh lead-colour, tipped with black; irides hazel. The top of the head, from the bill and chin, yellowiſh cream-colour; the reſt of the head and upper part of the neck bay, ſpeckled with with black; breaſt vinaceous; belly white; upper and under tail coverts black, edged with rufous above and with white beneath; back, ſcapulars, and ſides under the wings, black and white in fine undulated lines; coverts of the wings duſky brown, with pale edges; quills duſky; ſecondaries green on the outer webs, tipped with black; thoſe next the body have the margin of the outer webs more or leſs white; the tail is cuneiform; the two middle feathers pointed, and conſiderably longer than the reſt; theſe are duſky, daſhed with cinereous, the reſt cinereous, edged with rufous-brown; legs and feet duſky lead-colour, the latter ſmall.

This bird, like all the Duck tribe, is ſubject to

variety, according to age: in some the forehead is almost white, and the feathers on the back and sides mottled with brown; the wing coverts mixed with white.

The female is brown; the middle of each feather darkest; fore part of the neck and breast pale brown; wings and belly like the male. This species has a small *bagpipe* in the lower part of the *trachea*, peculiar to the male sex.

The Wigeon is found in most parts of Europe; breeds in the more northern parts. Visits England in the autumn, when great numbers are caught in our decoys for the table, being esteemed an excellent bird. It also frequents our rivers and salt-water inlets in small flocks.

WIGEON-PHEASANT-TAILED. Vide *Duck-long-tailed.*

WIGEON-PIED. Vide *Garganey.*

WIGEON-RED-HEADED. Vide *Pochard.*

WILLOCK. Vide *Razor-bill, Puffin, and Guillemot.*

WIND-HOVER. Vide *Kestrel.*

WINDLE. Vide *Redwing.*

WINNARD. Vide *Redwing.*

WINTER-MEW. Vide *Gull-winter.*

WIWALL. Vide *Woodpecker-greater-spotted.*

WOOD-CHAT.

Lanius rutilus. *Ind. Orn.* i. p. 70. 12.

Lanius Collurio. *Gmel. Syst.* i. p. 301. 12. γ.

Laniu

Lanius minor cinerascens. *Rati Syn.* p. 19. A. 6. *Will.* p. 54. iv. t. 10. f. 2?

Lanius femina. *Will.* p. 54. III.

Lanius rufus. *Brisſ.* ii. p. 147. 3.—*Ib.* 8vo. i. p. 199.

Pie-griesche rousse. *Buf.* i. p. 301.

Another sort of Butcher-bird. *Will. Angl.* p. 89. 4.

Wood-chat. *Br. Zool.* i. N° 73.—*Ib.* fol. 74. t. C. 1. *Albin.* ii. t. 16. *Lewin, Br. Birds,* i. t. 32. *Lath. Syn.* p. 169. 17. *Walc. Br. Birds,* i. t. 31. *Pult. Cat. Dorset.* p. 4. *Don. Br. Birds,* iv. t. 84.

This species of Shrike is said to be about the size of the Red-backed Shrike, or Butcher-bird; but as it has never come under our inspection, we must borrow the description of other ornithologists.

The bill is horn-colour; the feathers round the base whitish; head and hind part of the neck bright bay; from the base of the bill over the forehead, and through the eyes passing downwards on each side the neck, is a streak of black; back dusky; upper tail coverts grey; wing coverts dusky; quills black, near the bottom of which a white spot; throat, breast, and belly, yellowish white; the tail is black, the margins and tips whitish, except the two middle ones; legs black. Latham says the scapulars are white. The female is reddish on the upper parts of the body, the under parts dirty

white; every where tranſverſely ſtriated with brown.

The above is nearly the deſcription of Mr. Pennant and Dr. Latham, and which appears to be borrowed from earlier authors, as neither of them ſeems to have ſeen the bird. By the laſt mentioned author two varieties are given: one with the upper parts of the body rufous, under parts white; wings wholly brown, with a ſmall ſpot of white at the baſe of the quills: the other differs only in having the head black, and the tail rather longer.

We cannot help entertaining ſome doubts whether this may not hereafter prove to be no other than a variety of the Redbacked Shrike, or the young male in ſome of the intermediate ſtages between the neſtling and adult plumage, which with us rarely if ever appear, as thoſe birds leave us in autumn before they are maturely feathered. If the black ſtreak through the eye, and the grey on the rump, ſhould firſt appear, it would only want the white ſcapulars to make it the Woodchat; and that mark all authors have not deſcribed. Gmelin has given it as a mere variety of that bird.

WOODCOCK.

Scolopax ruſticola. *Lin. Syſt.* i. p. 243. 6. *Gmel. Syſt.* ii. p. 660. *Ind. Orn.* ii. p. 713. 1.

Scolopax. *Raii Syn.* p. 104. A. 1. *Will.* p. 213. t. 53. *Briſ.* v. p. 292. 1.—*Ib.* 8vo. ii. p. 284.

La Becaſſe. *Buf.* vii. p. 462. t. 25.

Woodcock.

Woodcock. *Br. Zool.* ii. N° 178. t. 65.—*Ib.* fol. 119. *Artt. Zool.* ii. p. 470. A.—*Ib. Sup.* p. 68. *Will. Angl.* p. 289. t. 53. *Albin.* i. t. 70. *Borlaſ. Cornw.* p. 245. t. 24. f. 12. *Lath. Syn.* v. p. 129. 1. *Lewin, Br. Birds,* iv. t. 156. *Walc. Syn.* ii. t. 136. *Pult. Cat. Dorſet,* p. 14.

This well-known ſpecies of Snipe needs little deſcription.

The length is fifteen inches; weight from twelve to fifteen ounces.

The bill is about three inches long, furrowed along the ſide of the upper mandible. Forehead cinereous; the reſt of the upper parts of the bird is a mixture of ferruginous, black, and grey, diſpoſed in bars; the under parts are yellowiſh white, with duſky ſtreaks; the tail conſiſts of twelve feathers, black on the inner web, the outer bordered with rufous; tips above cinereous, beneath white.

This bird is ſubject to great variety; ſome are very dark coloured and ſmall, others are large and the plumage much more rufous. Theſe have been thought two different ſexes, but without foundation.

The Woodcock appears firſt in the beginning of October, ſome few the latter end of September, but never in quantity till the middle of November. They generally come to us with northerly or eaſterly winds, when the more northern counties become

become frozen; and if the froſt in thoſe countries where they breed is ſuddenly ſevere large flights are ſometimes met with on our coaſts, where they remain for one day to recruit their ſtrength, and then diſperſe. With us it is not near ſo plentiful as formerly, when the art of ſhooting flying was leſs practiſed; a great many, however, are yet to be found in the more uncultivated parts of Devonſhire, Cornwall, and Wales, as well as in the north of Scotland; but no where ſo plentiful as in the large tracts of woods in Ireland. In ſevere weather they accumulate from the moors and inland counties to the woods in the weſt of England.

It is amongſt the few winter birds that occaſionally breed with us, many inſtances of which are recorded. The young birds have been killed in Auguſt, and eggs taken in June, both of which we can ſpeak of from our own knowledge. In the year 1795 we were favoured with two eggs from the Rev. Mr. Wheatear of Haſtings, who informed us they were found in a wood near Battel in Suſſex, and that there were four in the neſt. Theſe are larger than thoſe of a Pigeon, of a yellowiſh white, ſpotted and blotched with rufous-brown and aſh-colour, moſt ſo at the larger end.

The food of this bird is inſects and worms, for which it bores with its bill into moiſt places, feeding principally at night. They quit the woods in the duſk of the evening, and then only make a noiſe ſomething like the Snipe, but not ſo ſhrill.

At this time numbers were formerly caught in nets placed acroſs the glades of a wood, now much out of practice, except in the weſtern counties. Sometimes a net is placed by the ſide of a hedge, near to a wood, where they are taken either morning or evening as they fly from or to the wood. They are alſo taken in ſpringes placed on the ground in ſwampy places, where they are known to feed; and ſo much is their fleſh ſought after that ſince the introduction of mail-coaches to all parts the price of them in the country has increaſed more than double. We have known more than a hundred taken to London by one coach from the ſouth of Wales.

It is found in all parts of the old continent, from north to ſouth, in ſome places remaining the whole year, only changing their ſituation in the breeding ſeaſon from the plains to the mountains. We have been informed it viſits ſome parts of America in the winter; but we ſuſpect the Little Woodcock of that country has been confounded for it, as we have ſeen that ſpecies in plenty, during ſummer, about the temperate parts, which changes its ſituation with the ſeaſon to the warmer provinces.

WOODCOCK-SEA. Vide *Godwit-common.*

WOOD-CRACKER. Vide *Nuthatch.*

WOODPECKER. A genus of birds, the characters of which are,

Bill ſtrait, ſtrong, angular, and cuneated at the point.

Noſtrils

Noſtrils covered with reflected briſtles.
Tongue ſcarce longer than the bill, ſlimmiſh; the end horny and jagged, middle.
Toes, two forward, two backward.
Tail conſiſting of ten ſtiff ſharp-pointed feathers.

WOODPECKER-GREAT-BLACK.

Picus martius. *Lin. ſyſt.* i. p. 173. 1. *Gmel. Lin.* i. p. 424. *Ind. Orn.* i. p. 224. 1.

Picus niger. *Briſ.* iv. p. 21. 6.—*Ed. 8vo.* ii. p. 47.

——— *maximus.* *Raii Syn.* p. 42. 1. *Will.* p. 92. t. 21.

Le Pic noir. *Buf.* vii. p. 41. t. 2.

Great Black Woodpecker. *Albin.* ii. t. 27. *Will. Angl.* p. 135. *Arct. Zool.* ii. p. 276. A. *Lewin, Br. Birds,* ii. t. 46. *Lath. Syn.* ii. p. 552. 1. —*Ib. Sup.* p. 104. *Don. Br. Birds,* i. t. 13. *Walc. Syn.* i. t. 45. *Pult. Cat. Dorſet.* p. 6.

We give this bird as a Britiſh ſpecies with conſiderable doubt; but as Dr. Latham ſays he has been informed it is ſometimes met with in the ſouth, and in particular Devonſhire, a deſcription of it to the Engliſh ornithologiſt may be acceptable. It muſt, however, be obſerved, that author does not ſpeak of it as ſuch from his own knowledge, nor have we been able to aſcertain the fact.

It

It is about the ſize of a Jackdaw; length ſeventeen inches and a half; irides pale yellow. Bill near two inches and a half long, of a dark aſh-colour, and whitiſh on the ſides. The whole bird is black, except the crown of the head, which is vermilion; the two middle feathers of the tail are longer than the reſt; the legs are lead-coloured, covered with feathers on the fore part half their length.

The female has only the hind part of the head red, and the whole plumage has a tinge of brown. In ſome the red on the head is wholly wanting; but this marking in both ſexes is ſubject to much variation. This ſpecies is found in many parts of the European continent, but no where ſo plentiful as in Germany. It has all the habits of the Green Woodpecker, and is ſaid to be a great deſtroyer of bees. Makes its neſt deep in ſome tree which it has excavated for the purpoſe, and lays two or three white eggs, which ſeems peculiar to the whole genus.

WOODPECKER-GREEN.

Picus viridis. *Lin. Syſt.* i. p. 175. 12. *Gmel. Syſt.* i. p. 433. *Raii Syn.* p. 42. A. 2. *Will.* p. 93. t. 21. *Ind. Orn.* i. p. 234. 27. *Briſ.* iv. p. 9. 1.—*Ib.* 8vo. ii. p. 44.

Pic verd. *Buf.* vii. p. 7. t. 1.

Green Woodpecker. *Br. Zool.* i. N° 84.—*Ib.* fol. p. 78. t. E. *Arct. Zool.* ii. p. 277. B. *Albin.*

[illegible]

[illegible]

This species weighs about six ounces; length thirteen inches.

The bill is dusky, two inches long; the tongue near six inches; irides white. The feathers on the crown of the head dusky, tipped with cinereous; the eyes surrounded with black; beneath which, in the male is a crimson spot bordered with black, which in the female is wholly black; the neck, back, lesser covers of the wings, and scapulars, are green; the rump pale yellow; quill-feathers dusky, the greater spotted on each web with white, the lesser very faintly spotted on the exterior webs, and deeply bordered with green; the coverts of the ears and whole under parts are of a very pale yellow-green; the tail-feathers are stiff, pointed, alternately barred with dusky and green, tipped with black, except the outer feathers; legs ash-colour; claws much hooked.

This species is not uncommon about most of the wooded

wooded parts of England, its food is entirely inſects. The formation of the whole of this tribe is admirably adapted to their mode of life. The bill, which is ſtrong, and formed like a wedge at the point, enables them to force their way through the ſap of a tree, when by inſtinct it is diſcovered to be decayed at heart. With this inſtrument it diſlodges the *larvæ* of a numerous tribe of the coleopterous inſects, as well as that ſtinking caterpillar the *larva* of the Goat Moth (*Phalena Coſſus*), of which they frequently ſmell.

The tongue is no leſs wonderfully formed for inſinuating into all the ſmaller crevices to extract the hidden treaſure, by transfixing the larger inſects, or by adheſion withdraw the ſmaller; for, like the Wryneck, it is furniſhed with a glutinous ſubſtance for that purpoſe. Nor can we leſs admire the ſhort and ſtrong formation of the legs, and the hooked claws, ſo well calculated to enable them to climb and affix themſelves againſt the body of a tree, either to rooſt, or perforate a hole; to aſſiſt which the ſtiff tail is of infinite uſe. The jarring noiſe ſo frequently heard in woods in the ſpring is occaſioned by one or other of this genus, which, from frequent obſervation, we have no doubt is uſed as a call by both ſexes to each other. It is curious to obſerve them try every part of a dead limb till they have diſcovered the moſt ſonorous, and then the ſtrokes are reiterated with ſuch velocity that the head

head is ſcarcely perceived to move, the ſound of which may be diſtinctly heard half a mile.

Dr. Plot was miſtaken when he conceived this noiſe to be produced by the Nuthatch, who ſays, in deſcribing that bird, "That by putting their bill into a crack in the bough of a tree can make ſuch a violent ſound as if it was rending aſunder."

Woodpeckers are commonly ſeen climbing up a tree, but never down, as ſome have aſſerted. The hole which they make is as perfect a circle as if deſcribed by a pair of compaſſes. For the places of nidification the ſofter woods are attacked, the elm, aſh, and particularly the aſp, but rarely the oak. Theſe are only perforated where they have ſymptoms of decay; and the excavations are frequently deep, to give ſecurity to their eggs. This ſpecies lays four or five white eggs, weighing about two drams, which are placed on the rotten wood, without any neſt. The young birds have the appearance of crimſon on their heads, but not ſo bright as in adults.

Ants and their eggs are a favourite repaſt of this ſpecies, for which they are frequently ſeen on the ground ſearching the emmit hills. The tongue is here made uſe of inſtead of the bill, ſimilar to that of the Wryneck.

Its note is harſh, and its manner of flying undulated.

WOODPECKER-HAIRY.

Picus villoſus. *Lin. Syſt.* i. p. 175. 16. *Gmel. Syſt.* i. p. 435. *Ind. Orn.* i. p. 232. 19.

Picus varius virginianus. *Briſ.* iv. p. 48. 17. — *Ib.* 8vo. ii. p. 54.

Pic chevelu de Virginie. *Buf.* vii. p. 174.

Hairy Woodpecker. *Arct. Zool.* ii. N° 164. *Lewin, Br. Birds,* ii. t. 48. *Lath. Syn.* ii. p. 572. 18.—*Ib. Sup.* p. 108. *Walc. Syn.* i. t. 47.

This ſpecies is rather leſs than the Spotted Woodpecker; weight about two ounces; length eight inches and three quarters.

The bill is an inch and a quarter long, of a horn-colour. The head is black; on each ſide the head are two white ſtreaks, one over the eye, the other along the lower jaw, both ariſing from the baſe of the bill; acroſs the hind part of the head is a red band; the upper parts of the body are black, divided down the middle of the back with a liſt of white hair-like feathers; the wings are ſpotted with white; under parts of the body wholly white; the four middle feathers of the tail are black; the next on each ſide obliquely marked with white at the tip; the laſt but one white, with the baſe black; outer one wholly white; legs and claws greyiſh-brown.

The female wants the red on the head, in other reſpects like the male.

The Hairy Woodpecker is ſaid to be not uncommon in the north of England. Dr. Latham mentions having ſeen a pair in the collection of the Ducheſs of Portland, which were ſhot near Halifax in Yorkſhire. It is common in America. Said to be a great deſtroyer of apple-trees, by pecking holes in them.

WOODPECKER-SPOTTED-GREATER.

Picus major. *Lin. Syſt.* i. p. 176. 17. *Gmel. Syſt.* i. p. 436. *Ind. Orn.* i. p. 228. 13.

Picus varius major. *Raii Syn.* p. 43. A. 4. *Will.* p. 94. t. 21. *Briſ.* iv. p. 34. 13.—*Ib.* 8vo. ii. p. 51.

L'Epeiche, *ou* Pic varié. *Buf.* vii. p. 57.

Greater Spotted Woodpecker. *Br. Zool.* i. N° 85. —*Ib.* fol. p. 79. t. E. *Arct. Zool.* ii. N° 162. *Will. Angl.* p. 137. t. 21. *Albin.* i. t. 19. *Hayes, Br. Birds,* t. 9. *Lewin, Br. Birds,* t. 47. *Lath. Syn.* ii. p. 564. 12. — *Sup.* p. 107. *Walc. Syn.* i. t. 48. *Pult. Cat. Dorſet.* p. 6.

The weight of this ſpecies is about two ounces and three quarters; length nine inches. The bill is duſky, an inch and a quarter long; irides reddiſh brown. The forehead dirty white; crown of the head black; hind part of a deep crimſon; the cheeks are white, beneath which is a black line from the lower mandible to the back of the head, from the middle of which another line of the ſame colour

colour runs down on each ſide, and almoſt meets at the upper part of the breaſt; on each ſide the back of the neck is a white ſpot; the back, rump, coverts of the tail, and leſſer coverts of the wings, are black; the ſcapulars, and ſome of the greater coverts adjoining, are white; the quill-feathers are black, marked with white ſpots on each web; the throat and under parts are dirty white; vent bright crimſon; the four middle feathers of the tail are black, the reſt more or leſs white towards the point, marked with black ſpots; legs lead-colour.

The female differs only in having that part of the head black which in the male is red.

The Spotted Woodpecker is leſs frequent in England than the Green. Its habits and manners nearly the ſame, except that it rarely deſcends to the ground in ſearch of food, and that it more frequently makes that jarring noiſe than either of the other ſpecies, eſpecially when diſturbed from the neſt, which we had the opportunity of obſerving. It was with difficulty the bird was made to quit her eggs; for notwithſtanding a chiſel and mallet were uſed to enlarge the hole, ſhe did not attempt to fly out till the hand was introduced, when ſhe quitted the tree at another opening. The eggs were five in number, perfectly white and gloſſy, weighing about one dram, or rather more. Theſe were depoſited two feet below the opening, on the decayed wood, without the ſmalleſt appearance of a neſt.

As ſoon as the female had eſcaped, ſhe flew to a decayed branch of a neighbouring tree, and there began the jarring noiſe before mentioned, which was ſoon anſwered by the male from a diſtant part of the wood, but ſoon joined his mate, and both continued theſe vibrations, trying different branches till they found the moſt ſonorous.

WOODPECKER-SPOTTED-LESSER.

Picus minor. *Lin. Syſt.* i. p. 176. 19. *Gmel. Syſt.* i. p. 437. *Ind. Orn.* i. p. 229.

Picus varius tertius. *Raii Syn.* p. 43. 6. *Will.* p. 94. t. 21.

Picus varius minor. *Briſ.* iv. p. 41. 15.—*Ib.* 8vo. ii. p. 53.

La petite Epeiche. *Buf.* vii. p. 62.

Leſſer Spotted Woodpecker. *Br. Zool.* i. N° 87. t. 37.—*Ib.* fol. 79. t. E. *Arct. Zool* p. 278. E. *Albin.* i. t. 20. *Lewin, Br. Birds*, t. 49. *Lath. Syn.* ii. p. 566. 14.—*Sup.* p. 107. *Walc. Syn.* i. t. 50. *Pult. Cat. Dorſet.* p. 6. *Don. Br. Birds*, ii. t. 36.

The weight of this ſmall ſpecies is not quite five drams; length five inches and a half.

The bill is lead-colour; irides red-brown. Forehead, breaſt, and belly, dirty white; the cheeks and ſides of the neck white, bounded by black above and below; crown of the head crimſon, bounded

bounded on each ſide with black, which, joining behind, runs down in a peak on the back of the neck; the back and ſcapulars are barred with black and white; the quill-feathers, and ſome of the larger coverts, are black, ſpotted with white, the leſſer coverts wholly black; the four middle feathers of the tail are black, the reſt more or leſs white at the ends, ſpotted with black; the outer one almoſt white; legs lead-colour.

The female reſembles the other ſex in plumage, except on the crown of the head, which is white where the male is red; neither ſex have any red at the vent.

The male ſometimes has a few duſky ſtreaks on the ſides of the breaſt.

Young male birds poſſeſs the red on the head before they leave their neſt; a circumſtance not uſual in birds that have ſuch vivid colours.

This is by much the ſcarceſt ſpecies; it poſſeſſes all the habits of the preceding; the note is alſo the ſame, but not ſo ſtrong.

The eggs are white, and weigh about thirty-three grains, five of which we took out of a decayed tree, depoſited on the rotten wood, without any neſt, at a conſiderable diſtance below the entrance. The aperture correſponded with the ſize of the bird, but did not appear recently made. It is probable, however, it is able to perform this work for itſelf; and inſtinct points out the inſecurity in making choice of a larger opening to their

place of incubation, as they would then be liable to be diſlodged by the larger ſpecies, the Daw and the Stare. We have received it from Glouceſter-ſhire by the names of Hickwall and Crank-bird; have alſo ſeen it in Wiltſhire, where we took its eggs.

WOODPECKER-SPOTTED-MIDDLE.

Picus medius. *Lin. Syſt.* i. p. 176. 18. *Gmel. Syſt.* i. p. 436. *Ind. Orn.* i. p. 229. 14.

Picus varius minor. *Raii Syn.* p. 43. 5.

———— major. *Will.* t. 21.

Picus varius. *Briſ.* iv. p. 38. 14. t. 2. f. 1.—*Ib.* 8vo. ii. p. 52.

Middle Spotted Woodpecker. *Br. Zool.* i. N°86. t. 37. *Arct. Zool.* ii. p. 278. D. *Lewin, Br. Birds*, t. 48. *Lath. Syn.* ii. p. 565. 13. —*Ib. Sup.* p. 107. *Walc. Syn.* i. t. 49. *Pult. Cat. Dorſet.* p. 6.

This bird, which we conſider as the young of the Greater Spotted Woodpecker, but deſcribed by Linnæus and others as a diſtinct ſpecies, differs only from that bird in having the whole crown of the head red; the black marking about the head and neck not ſo large, and the white on the cheeks more duſky; that on the breaſt inclining to brown; the length and weight nearly the ſame. It ſhould ſeem that the young males of the Greater Spotted Woodpecker have at firſt the whole crown of the

head

head red; but that in two or three months after they leave the neſt the red feathers on the top of head are diſplaced by black ones. In this ſtate of change we have now one before us, which was killed in November. The back of the head is crimſon; the crown black, ſpotted with crimſon feathers; and what ſtrengthens this opinion is, that theſe birds are only to be met with from July to the latter end of November, as far as we have been able to collect; at leaſt we have never been able to procure ſuch at any other time.

Dr. Latham ſeems to favour this opinion. Buffon alſo appears to conſider it as a variety only of the greater ſpotted ſpecies.

The reaſon why this ſhould appear ſcarcer is, that few Woodpeckers are killed when the trees are in full foliage, and by the time the leaves are fallen theſe have aſſumed their full plumage.

WOODSPITE. Vide *Woodpecker-green.*

WREN-COMMON.

Motacilla Troglodytes. *Lin. Syſt.* i. p. 337. 46. *Gmel. Syſt.* ii. p. 993. *Raii Syn.* p. 80. A. 11. *Will.* p. 164. t. 42.

Sylvia Troglodytes. *Ind. Orn.* ii. p. 547. 148.

Regulus. *Brif.* iii. p. 425. 24.—*Ib.* 8vo. i. p. 428.

Troglodyte, Roitelet. *Buf.* v. p. 352. t. 16. f. 1.

Wren. *Br. Zool.* i. N° 154.—*Ib.* fol. 102. t. 42. *Arct. Zool.* ii. N° 322. *Albin.* i. t. 53. B.—

Ib. Song Birds, t. p. 64. *Will. Angl.* p. 229. t. 42. *Hayes, Br. Birds*, t. 38. *Lath. Syn.* iv. p. 506. 143. *Lewin, Br. Birds*, iii. t. 111. *Wale. Syn.* ii. t. 242. *Pult. Cat. Dorſet.* p. 9.

This ſpecies of Warbler weighs about two drams and three quarters; length near four inches and a quarter.

The bill is ſlender, full half an inch in length, a little curved, and of a duſky brown colour; irides dark hazel. The head, neck, and upper parts of the body, are of a deep reddiſh brown, obſcurely marked with tranſverſe duſky lines; over the eye a light-coloured ſtreak; quills and tail duſky brown; the former ſpotted on the outer webs with light brown, the latter croſſed with duſky black lines; the under parts light rufous-brown; the ſides and thighs croſſed with darker lines; under tail coverts obſcurely ſpotted with black and white; legs pale brown.

This pretty little warbler, like the Redbreaſt, frequently approaches the habitation of man, and enlivens the ruſtic garden with its ſong the greater part of the year. It begins to make a neſt early in the ſpring, but frequently deſerts it before it is lined, and ſearches for a more ſecure place. It is frequently made under the thatch of out-buildings, againſt the ſide of a moſſy tree, or againſt an impending bank that ſecures it from the rain; ſometimes in a low thick buſh. But what is remarkable,

the

the materials of the neſt are generally adapted to the place: if built againſt the ſide of a hay-rick, it is compoſed of hay; if againſt the ſide of a tree covered with white moſs, it is made of that material; and with green moſs if againſt a tree covered with the ſame, or in a bank. Thus inſtinct directs it for ſecurity. The lining is invariably feathers.

The eggs are ſeven or eight in number, white, ſparingly marked with ſmall reddiſh ſpots, moſt commonly at the larger end; their weight about twenty grains.

The Wren does not begin the bottom of its neſt firſt, which is uſual in moſt birds, but firſt, as it were, traces the outlines if againſt a tree, which is of an oval ſhape, and by that means faſtens it equally ſtrong to all parts, and afterwards encloſes the ſides and top, leaving only a ſmall hole near the top for entrance. If the neſt is placed under a bank, the top is firſt begun, and well ſecured in ſome ſmall cavity, by which the fabric is ſuſpended.

The food of this ſpecies is inſects, which it finds in ſufficient abundance to ſupport life even in the ſevereſt winters.

WREN-GOLDEN-CRESTED.

Motacilla Regulus. *Lin. Syſt.* i. p. 338. 48. *Gmel. Syſt.* ii. p. 995.

Sylvia Regulus. *Ind. Orn.* ii. p. 548. 152.

Regulus criſtatus. *Raii Syn.* p. 79. A. 9. *Will.* p. 163. t. 42. *Briſ.* iii. p. 579. 17.—*Ib.* 8vo. i. p. 472.

Roitelet,

Roitelet, Poul, Souci. *Buf.* v. p. 363. t. 16. f. 2.
Golden-creſted Wren. *Br. Zool.* N° 153.—*Ib.* fol. 101. t. S. f. 3. *Arct. Zool.* ii. N° 321.—*Ib. Sup.* p. 64. *Will. Angl.* p. 227. *Edw.* t. 254. 1. *Albin.* i. t. 53. *Hayes, Br. Birds,* t. 38. *Lath. Syn.* iv. p. 508. 145. *Lewin, Br. Birds,* iii. t. 112. *Walc. Sys.* ii. t. 243. *Pult. Cat. Dorſet.* p. 9. *Don. Br. Birds,* i. t. 4.

This elegant little ſpecies of Warbler is the ſmalleſt Britiſh bird.

Its weight ſeldom exceeds eighty grains; length three inches and three quarters.

The bill is ſlender and black; irides hazel. The crown of the head is ſingularly beautiful; the creſt is compoſed of a double ſeries of feathers ariſing from each ſide, and almoſt meet at their points; the exterior are black, the interior bright yellow; between which, on the crown, the feathers are ſhorter, and of a fine deep orange; the forehead, chin, and round the eyes, whitiſh; the hind part of the head, neck, and back, green; the two firſt daſhed with aſh-colour; quills duſky, edged with green; at the baſe of the ſecondary quills is a black bar, above which the coverts are tipped with white, forming a narrow bar of that colour; above that the ſmaller coverts are black, tipped with white, which form two other ſmall obſcure bars; the under parts of the body are browniſh white, lighteſt

on the belly, a little tinged with yellow; the tail is ſomewhat forked, the feathers duſky, edged with yellowiſh green.

The female has the head rather leſs creſted, and the crown is bright yellow where the male is orange.

The young birds do not poſſeſs the creſt and yellow feathers till autumn.

The neſt is not made with an opening on one ſide, as deſcribed by ſome, but is in form and elegance like that of the Chaffinch, compoſed of green moſs, interwoven with wool, and invariably lined with ſmall feathers, with which it is ſo well bedded as to conceal the eggs. It is ſometimes placed againſt the body of a tree covered with ivy, but moſt times underneath a thick branch of a fir. The eggs are from ſeven to ten in number, of a browniſh white, rather darker at the larger end; their weight nine or ten grains.

This minute ſpecies braves the ſevereſt winters of our climate, being equally found in all parts in that rigorous ſeaſon; and is by no means ſo ſcarce a bird as it appears, but from its ſmallneſs is ſeldom noticed. It would in all probability be much more plentiful, but from ſome cauſe we have not been able to diſcover, the female is frequently deſtroyed at the time of incubation, and the neſt with eggs left to decay. For a further account of this beautiful little bird ſee the Introduction.

In Cornwall it is called Wood Titmouſe, in Devonſhire Tidley Goldfinch.

WREN-GREEN. Vide *Wren-wood.*

WREN-GROUND. Vide *Wren-yellow.*

WREN-REED.

Motacilla arundinacea. *Gmel. Syſt.* i. p. 992.
Sylvia arundinacea. *Ind. Orn.* ii. p. 510. 12.
Paſſer arundinaceus minor. *Raii Syn.* p. 47. A. 3? *Will.* p. 99?
Bouſcarle de Provence. *Buf.* v. p. 134?
Leſſer Reed-Sparrow. *Will. Angl.* p. 144?
Reed Wren. *Lath. Syn. Sup.* p. 184. *Lewin, Br. Birds,* iii. t. 114.

The length of this ſpecies of Warbler is ſcarce five inches and a half; weight nearly three drams.

The bill is about half an inch in length, duſky above, yellowiſh beneath, and broad at the baſe; at the corner of the mouth are three ſtrong briſtles; irides hazel.

The plumage of the whole upper parts of the bird are of a plain olive-brown; the under parts yellowiſh white, lighteſt on the throat and down the middle of the belly; the ſides a little inclining to rufous-brown; from the bill to the eye is an obſcure lightiſh ſtreak; eyelids lighter, but no ſtroke over the eye; the tail is cuneiform; the feathers, like thoſe of the quills, duſky brown, edged with the ſame colour as the back; legs duſky brown.

This bird has been in general confounded with the Sedge Warbler; its form, ſize, manners, and habits, are alike, and are both migrative ſpecies;

ſo that it is difficult to diſcover which of theſe birds moſt authors mean by their deſcriptions. The Reed Wren, however, may at once be diſtinguiſhed from the other by the baſe of the bill being broader; in having no light ſtroke over the eye, which in the other is broad and conſpicuous; and in the whole upper parts being of one plain colour.

Its neſt and eggs are alſo different. The neſt is compoſed of long graſs and the ſeed branches of reeds, and lined with the finer parts of the latter; it is very deep, and conceals the bird when ſitting. This is generally faſtened by long graſs to ſeveral reeds which are drawn together for that purpoſe, and generally placed over the water. The eggs are four or five in number, rather larger than thoſe of the Sedge Warbler, of a greeniſh white, blotched all over with duſky brown.

This ſpecies is much more local than the Sedge bird, but are ſometimes found together. Their notes are ſimilar. We have found both ſpecies all along the coaſts of Kent and Suſſex, from Sandwich to Arundel, amongſt the reedy pools and ditches, eſpecially on Romney Marſh; but in Wiltſhire and Somerſetſhire, where the Sedge Warbler is found in abundance throughout the banks of the Avon, not a ſingle Reed Wren is to be found.

The neſt of this bird being deep gives ſecurity to the eggs, which would otherwiſe be thrown out by

by the wind. We have ſeen the bird ſitting on her neſt when the wind blew hard, and at every guſt forced it almoſt to the ſurface of the water.

It makes its firſt appearance with us the latter end of April, or beginning of May, and departs in September. Is ſaid to be found in the fens of Lincolnſhire, and the banks of the river Coln in Buckinghamſhire.

WREN-WILLOW. Vide *Wren-yellow.*

WREN-WOOD.

Sylvia Sylvicola. *Lin. Tranſ.* iv. p. 35.

Regulus non criſtatus major. *Will.* p. 164.—*—Ib. Angl.* p. 228. *White, Hiſt. Selb.* p. 55.

Wood Wren. *Lin. Tranſ.* ii. p. 245. t. 24.

Green Wren. *Albin.* ii. t. 86. 6?

This ſpecies of Warbler weighs about two drams forty grains; length five inches and a quarter. The bill is duſky; irides hazel. The upper part of the head, back, ſcapulars, wing coverts, and upper coverts of the tail, are of a lively yellow-green; over the eye is a bright brimſtone-coloured ſtreak; the cheeks and throat are yellow; the upper part of the breaſt white, tinged with yellow; the lower part, the belly, and under tail coverts, pure white; the quill-feathers are duſky, edged on their exterior webs with yellow-green; the tail a little forked, coloured like the quills, except the two outmoſt feathers, which want the yellow margin; the legs yellowiſh brown.

The

The female is rather larger, weighing about three drams; the plumage exactly like the male.

The Wood Wren is a migrative ſpecies, appearing with us firſt about the latter end of April, the females ten days or a fortnight later, and depart again in September. The reaſon this bird has been ſo little noticed as a diſtinct ſpecies is, from its great ſimilitude to the Yellow Wren, for which it has been confounded, and probably likewiſe for the Leſſer Pettychaps. But its ſuperior ſize to the latter, and the pure whiteneſs of the under tail coverts, are characteriſtic marks of diſtinction; which part in both the others is tinged with yellow; the colour of the upper parts is alſo much more vivid, and the ſtroke above the eye brighter yellow. It differs alſo from thoſe birds in manners and habits: this is found to inhabit woods only in the breeding ſeaſon. The neſt is placed on the ground, in form like both that of the other birds, being oval, with a ſmall hole near the top, conſtructed of dry graſs, a few dead leaves, and a little moſs; and invariably lined with finer graſs and a few long hairs. The other ſpecies always line their neſts with feathers.

The eggs are ſix in number, white, ſprinkled all over with purpliſh ſpots. In ſome theſe markings are confluent, inclining to ruſt-colour; their weight from eighteen to twenty-two grains.

This is not an uncommon ſpecies, having met with it in moſt parts of the ſouth of England and Wales, and as far weſtward as Cornwall. It ſeems

partial

partial to oak and beech woods, where it may be found by its ſingular note, which ſeems to expreſs the word *twee*, drawn out to ſome length, and repeated five or ſix times ſucceſſively, terminating with the ſame notes delivered in a hurried manner, at which time it ſhakes its wings. This ſeems to be the extent of its ſong, the latter part of which is chiefly left out after the breeding ſeaſon.

It is alſo found in Germany, where it is likewiſe a migrative bird. Dr. Beckſtein deſcribes it in the 27th volume of *Naturforſcher*, under the title Der Laubrogelchen (Little leaf-bird). In the Leverian Muſeum, caſe N° 271, we obſerved it inſcribed Green Wren.

WREN-YELLOW.

Motacilia Trochilus. *Lin. Syſt.* i. p. 338. 49. *Gmel. Syſt.* ii. p. 995.

Sylvia Trochilus. *Ind. Orn.* ii. p. 550. 155.

Aſilus. *Brif.* iii. p. 479. 45.—*Ib.* 8vo. i. p. 441. *Raii Syn.* p. 80. A. 10. *Will.* p. 164.—*Ib. Angl.* p. 228.

Le Pouillot. *Buf.* v. p. 344.

Yellow Wren. *Br. Zool.* i. N° 151.—*Ib.* fol. 101. t. S. f. 2. S. 2. f. 1. *Arct. Zool.* ii. N° 319. *White, Hiſt. Selb.* pp. 28. 55. *Hayes, Br. Birds,* t. 38. *Lath. Syn.* iv. p. 512. 147. *Don. Br. Birds,* i. t. 14. *Lewin, Br. Birds,* iii. t. 113. *Pult. Cat. Dorſet.* p. 9.

VAR.

Motacilla acredula. *Lin. Syſt.* i. p. 338. 49. β.
Ficedula Carolinenſis. *Briſ.* iii. p. 486. 48.—*Ib.* 8vo. i. p. 443.
Œnanthe fuſco-lutea minor. *Raii Syn.* p. 186. 39.
Le Fiquier brun et jaune. *Buf.* v. p. 295.
Yellow Wren. *Edw.* t. 278. f. 2.
Scotch Wren. *Br. Zool.* ii. N° 152. *Arct. Zool.* ii. p. 420. N.—*Ib. Sup.* p. 64. *Lath. Syn.* iv. p. 513.

PROVINCIAL.

Willow Wren. Ground Wren. Ground Huckmuck.

This ſpecies weighs about two drams and three quarters; length five inches and a quarter. The bill is duſky above, yellowiſh beneath; irides hazel. The whole upper parts of the plumage are of a greeniſh yellow-brown; the under parts are white, tinged with yellow; on the breaſt are a few yellow ſtreaks; quills duſky brown, edged with yellow; coverts yellow, thighs the ſame; tail like the quills, ſlightly edged the ſame; over the eye a faint yellowiſh ſtroke; legs light brown.

This is a plentiful ſpecies in ſome parts. Frequents wooded and encloſed ſituations, eſpecially where willows abound. Is frequently found with the Wood Wren, but does not extend ſo far to the weſt in England, as it is rarely met with in Cornwall. It comes to us early in April, and ſoon be-

gins its uſual ſong, which is ſhort, with little variety. About the latter end of the ſame month, or beginning of May, it makes a neſt of an oval ſhape, with a ſmall opening near the top, compoſed of moſs and dried graſs, and lined with feathers. This is placed in the hollow of a ditch, or in a low buſh cloſe to the ground.

The eggs are ſix or ſeven in number, white, ſpotted with light ruſt-colour towards the larger end; others are ſprinkled all over; their weight ſixteen or eighteen grains.

The plumage of this bird is ſo like that of the Leſſer Pettychaps that, were it not for its ſuperior ſize, it would be ſcarcely diſtinguiſhable. The colour of the legs, however, is an unerring mark of diſtinction. It has alſo been frequently confounded with the Wood Wren; but each have their eſſential characters; we therefore refer our readers for further particulars to thoſe ſpecies. The egg of this bird is readily diſtinguiſhed from that of the Leſſer Pettychaps, the ſpots in this being of a ruſt-colour, thoſe in the other invariably dark purple.

WRYNECK. This genus conſiſts of only a ſingle ſpecies, no other having as yet been diſcovered in any part of the world. The characters are as follows:

Bill roundiſh, ſlightly incurvated, and not very ſtrong.

Noſtrils bare of feathers, and ſomewhat concave.

Tongue

Tongue three times the length of the bill, ſlender, and armed at the point with a horny ſubſtance.

Tail conſiſts of ten flexible feathers.

Toes placed two backwards, two forwards.

WRYNECK.

Yunx torquilla. *Lin. Syſt.* i. p. 172. *Gmel. Syſt.* i. p. 423. *Raii Syn.* p. 44. A. 8. *Will.* p. 95. t. 22. *Ind. Orn.* i. p. 223. 1. *Briſ.* iv. p. 4. t. 1. f. 1.—*Ib.* 8vo. ii. p. 43.

Le Torcol. *Buf.* vii. p. 84. t. 3.

Wryneck. *Br. Zool.* i. N° 83.—*Ib.* fol. 80. t. F. G. *Arct. Zool.* ii. p. 267. B. *Will. Angl.* p. 138. t. 22. *Albin.* i. t. 21. *Lewin, Br. Birds,* ii. t. 43. *Lath. Syn.* ii. p. 548. t. 24. —*Ib. Sup.* p. 103. *Walc. Syn.* i. t. 44. *Pult. Cat. Dorſet.* p. 6. *Don. Br. Birds,* iv. t. 83.

PROVINCIAL.

Long-tongue. Emmet-hunter.

The Wryneck ſeems to be the link between the Woodpecker and the Cuckow; it has the tongue and feet of the former, but not the ſtrong angular bill or ſtiff tail; it has alſo the feet, and ſomewhat the bill, of the latter; as well as the ſame number of flexible feathers in the tail.

The weight of this beautiful bird is about ten drams; the length ſeven inches; the bill is three quarters of an inch long, nearly ſtrait, and ſharp-pointed; tongue cylindric, two inches and a quarter in length; irides light hazel, inclining to yellow.

The upper part of the head and neck, back, rump, and upper tail coverts, are aſh-colour, marked with numerous fine duſky, tranſverſe, undulated lines; from the crown of the head a black liſt runs down half way the back, diſpoſed in four indiſtinct lines; the wing coverts like the back, but inclining to rufous, and ſpotted with grey; the greater quills duſky, marked on their exterior webs with quadrangular ruſt-coloured ſpots; the chin and fore part of the neck yellowiſh white, croſſed with duſky lines; lower part of the breaſt, belly, and ſides, white, with ſmall triangular duſky ſpots; under tail coverts rufous-white, croſſed with duſky lines; the tail-feathers are broad, and rounded at their ends, and of the ſame colour as the back, with four diſtinct black bars; legs brown.

The pen or the pencil can only give a very inadequate idea of the elegant markings of this bird.

The name of Wryneck ſeems to have been given it from the ſingular manner of turning its head over its ſhoulder alternately, at which time the black liſt on the back of the neck gives it a twiſted appearance; it alſo erects all the feathers on the crown of the head in a terrific manner. It has been called the Emmet-hunter with great propriety, that inſect being its chief food.

The bill of this bird ſeems to be of little uſe in procuring ſubſiſtence, the tongue being the chief inſtrument uſed for ſuch purpoſes.

We were enabled to examine the manners of

this bird minutely by taking a female from her neſt, and confining her in a cage for ſome days. A quantity of mould with emmets and their eggs were given it; and it was curious to obſerve the tongue darted forward and retracted with ſuch velocity, and with ſuch unerring aim, that it never returned without an ant or an egg adhering to it, not transfixed by the horny point, as ſome have imagined, but retained by a peculiar tenacious moiſture, by nature provided for that purpoſe. While it is feeding the body is motionleſs, the head only is turned to every ſide, and the motion of the tongue is ſo rapid that an ant's egg, which is of a light colour, and more conſpicuous than the tongue, has ſomewhat the appearance of moving towards the mouth by attraction, as a needle flies to a magnet. The bill is rarely uſed except to remove the mould in order to get more readily at theſe inſects; where the earth is hollow the tongue is thruſt into all the cavities to rouſe the ants; for this purpoſe the horny appendage is extremely ſerviceable, as a guide to the tongue. We have ſeen the Green Woodpecker take its food in a ſimilar manner; and moſt probably every ſpecies of that genus does the ſame.

The Wryneck makes a noiſe very much like the ſmaller ſpecies of Hawks, which it frequently repeats in the ſpring, ſoon after its arrival, which is generally about the middle of April, ſometimes earlier in that month. Appearing at the ſame time

with the Cuckow, it has been termed that bird's ſervant or attendant.

If this bird is ſurpriſed in its neſt, it has a ſingular action of defence: ſtretching itſelf at full length, and erecting the feathers on the head, it ſuddenly riſes, making at the ſame time a ſhort hiſſing noiſe, ſomething like that of a cock Turkey. This was frequently done in the cage when it was approached.

It is not uncommon in the ſouthern and eaſtern parts of the kingdom, but is more ſcarce weſtward, and rarely if ever found in Cornwall. It chiefly inhabits woods, or thickly-encloſed countries, where trees abound, or orchards; is ſeldom obſerved to climb trees, although the feet ſeem calculated for that purpoſe.

It makes no neſt, but depoſits its eggs, to the number of nine or ten, in the hole of a decayed tree, on the rotten wood; theſe are perfectly white, weighing about thirty-ſeven grains.

WYNKERNEL. Vide *Gallinule-ſpotted.*

Y.

YAFFLE, or YAFFLER. Vide *Woodpecker-green.*
YARWHELP. Vide *Godwit-common.*
YARWIP. Vide *Godwit-common.*
YELLOW-BIRD. Vide *Oriole.*
YELLOW-BIRD-LITTLE. Vide *Wren-yellow.*
YELLOW-HAMMER. Vide *Bunting-yellow.*
YELPER. Vide *Avoſet.*

APPENDIX.

SANDPIPER-LITTLE.

Tringa pucilla. *Lin. Syſt.* i. p. 252. 20.

A female of this ſpecies, lately killed on the ſouth coaſt of Devonſhire, differs ſomewhat from that deſcribed in the *Britiſh Zoology*; and as it is an extremely rare bird in England we judge it proper to give a deſcription of it in this place.

Weight ſix drams; length ſix inches.

Bill duſky, three quarters of an inch long, very ſlender, a little bending downward, and rather larger near the point than in the middle; irides duſky. The forehead, crown of the head, back of the neck, back and ſcapulars, dark cinereous, duſky down the ſhafts, except on the neck, which is rather lighter coloured than the reſt; from the bill to the eye a duſky brown ſtreak; above that an obſcure dirty white one; chin and throat white; fore part of the neck and upper breaſt pale cinereous-brown; lower breaſt, belly, vent, and under tail coverts, pure white; primores, ſecondaries, and the greater coverts immediately impending them, duſky, very ſlightly tipped with white, moſt ſo on the coverts, and the primores margined with white on the outer webs, except the two firſt; the ſhaft of the firſt quill is white, the others duſky brown; ſpurious wing, and ſmaller coverts near it, duſky; thoſe along the ridge of the wing duſky and cinereous; the reſt of the coverts and tertials cinereous, like the back, a few of the former edged with pale rufous-brown; the rump and upper tail coverts duſky, the feathers

ſlightly tipped with cinereous; the tail is cuneiform, compoſed of twelve feathers, the ſix middle ones are cinereous, the two middlemoſt inclining to duſky; the three outer feathers on each ſide pure white; legs light olive-brown, three quarters of an inch long; claws black; outer toe very ſlightly connected at the baſe by a membrane to the middle one.

This bird was ſhot for a Jack Snipe on a ſalt marſh not very remote from the ſea in the month of November.

SANDPIPER-YELLOW-LEGGED.

Tringa Flavipes.

This ſpecies is about the ſize of the Grey Sandpiper; length eleven inches and a half.

Bill one inch and a half long, black half way from the end, baſe yellow; irides light yellow. Head and neck cinereous, ſtreaked with duſky; the upper parts of the body cinereous-brown, the middle of each feather duſky; wing coverts the ſame; beneath the ſpurious wing a ſmall patch of white; primary quill-feathers duſky, the firſt with a white ſhaft; ſecondaries white half way from their tips; the under parts white; rump the ſame; the tail and its upper coverts cinereous-brown; under tail coverts ſpeckled with duſky; legs orange yellow.

This bird was bought in the market at Bath in the winter, a drawing of which was taken by Mr. Robins of that place, painter of natural hiſtory, from whom we obtained the deſcription, with a drawing.

It appears to be a nondeſcript ſpecies, or at leaſt is eſſentially different from any one of the genus hitherto deſcribed.

A LIST OF BRITISH BIRDS,

SYSTEMATICALLY ARRANGED INTO

ORDINES, GENERA, AND SPECIES.

DIVISION I. LAND BIRDS.
DIVISION II. WATER BIRDS.

DIV. I.

ORDER I. Rapacious.

GENUS I.

FALCON.		FALCO.
Golden Eagle,	-	Chryſaetos.
Sea E.	-	Oſſifragus.
Ringtail E.	-	Fulvus.
Cinereous E.	-	Albicilla.
Oſprey,	-	Haliæetus.
Common Buzzard,	-	Buteo.
Honey B.	-	Apivorus.
Moor B.	-	Æruginoſus.
Goſhawk,	-	Palumbarius.
		Kite,

Kite,	-	Milvus.
Gentil Falcon,	-	Gentilis.
Peregrine F.	-	Peregrinus.
Spotted F.	-	Verſicolor.
Rough-legged F.	-	Lagopus.
Grey F.	-	Griſeus.
Jerfalcon,	-	Iſlandicus.
Lanner,	-	Lanarius.
Hen-harrier,	-	Cyaneus.
Keſtril,	-	Tinnunculus.
Sparrow-Hawk,	-	Niſus.
Hobby,	-	Subbuteo.
Merlin,	-	Æſalon.
Aſh-coloured F.	-	Cineraceus.

GENUS II.

OWL.		STRIX.
Great-eared O.	-	Bubo.
Long-eared O.	-	Otus.
Short-eared O.	-	Brachyotus.
White O.	-	Flammea.
Tawny O.	-	Stridula.
Little O.	-	Paſſerina.

GENUS III.

SHRIKE.		LANIUS.
Great Cinereous Shrike,	-	Excubitor.
Red-backed Shrike,	-	Collurio.
Wood Chat,	-	Rutilus.

ORDER

ORDER II. Pies.

GENUS IV.

CROW.	CORVUS.
Raven, -	Corax.
Crow, -	Corone.
Rook, -	Frugilegus.
Hooded Crow, -	Cornix.
Jackdaw, -	Monedula.
Jay, -	Glandarius.
Magpie, -	Pica.
Red-legged Crow, -	Graculus.
Nutcracker, -	Caryocatactes.

GENUS V.

ROLLER.	CORACIAS.
Garrulus Roller, -	Garrula.

GENUS VI.

ORIOLE.	ORIOLUS.
Golden Oriole, -	Galbula.

GENUS VII.

CUCKOW.	CUCULUS.
Common Cuckow, -	Canorus.

GENUS VIII.

WRYNECK.	YUNX.
Common Wryneck, -	Torquilla.

GENUS

GENUS IX.

WOODPECKER.	PICUS.
Great Black W.	Martius.
Greater Spotted W.	Major.
Middle Spotted W.	Medius.
Leſſer Spotted W.	Minor.
Hairy W.	Villoſus.
Green W.	Viridis.

GENUS X.

KINGSFISHER.	ALCEDO.
Common K.	Iſpida.

GENUS XI.

NUTHATCH.	SITTA.
European N.	Europæa.

GENUS XII.

BEE-EATER.	MEROPS.
Common Bee-eater,	Apiaſter.

GENUS XIII.

HOOPOE.	UPUPA.
Common H.	Epops.

GENUS XIV.

CREEPER.	CERTHIA.
Common C.	Familiaris.

ORDER III. Paſſerine.

GENUS XV.

STARE.	STURNUS.
Common S.	Vulgaris.

GENUS

GENUS XVI.

THRUSH.	TURDUS.
Miſſel Thruſh,	Viſcivorus.
Throſtle,	Muſicus.
Fieldfare,	Pilaris.
Redwing,	Iliacus.
Blackbird,	Merula.
Ring Ouzel,	Torquatus.
Water O.	Cinclus.
Roſe-coloured Thruſh,	Roſeus.

GENUS XVII.

CHATTERER.	AMPELIS.
Waxen Chatterer	Garrulus.

GENUS XVIII.

GROSBEAK.	LOXIA.
Common Croſsbill,	Curviroſtra.
Hawfinch,	Coccothrauſtes.
Pine Groſbeak,	Enucleator.
Greenfinch,	Chloris.
Bulfinch,	Pyrrhula.

GENUS XIX.

BUNTING.	EMBERIZA.
Common Bunting,	Miliaria.
Yellow B.	Citrinella.
Reed B.	Schæniclus.

Snow

Snow Bunting,	-	Nivalis.
Tawny B.	-	Glacialis.
Mourning B.	-	Montana.
Green-headed B.	-	Turdalis.
Cirl B.	-	Cirlus.

GENUS XX.

FINCH.		FRINGILLA.
House Sparrow,	-	Domeſtica.
Tree Sparrow,	-	Montana.
Chaffinch,	-	Cælebs.
Brambling,	-	Montifringilla.
Goldfinch,	-	Carduelis.
Siſkin,	-	Spinus.
Common Linnet,	-	Linota.
Greater Redpole,	-	Cannabina.
Leſſer Redpole,	-	Linaria.
Twite,	-	Montium.

GENUS XXI.

FLYCATCHER.		MUSCICAPA.
Spotted Flycatcher,	-	Griſola.
Pied F.	-	Atricapilla.

GENUS XXII.

LARK.		ALAUDA.
Sky Lark,	-	Arvenſis.
Wood L.	-	Arborea.
Red L.	-	Rubra.

Leſſer-

Leſſer-crested Lark,	- -	Criſtatella.
Field L.	- -	Minor.
Tit L.	- -	Pratenſis.
Pipit L.	- -	Trivialis.
Rock L.	- -	Petroſus.

GENUS XXIII.

WAGTAIL.		MOTACILLA.
White Wagtail,	-	Alba.
Grey W.	-	Boarula.
Yellow W.	-	Flava.

GENUS XXIV.

WARBLER.		SYLVIA. (Motacilla, *Lin.*)
Nightingale,		Luſcinia.
Greater Pettychaps,	-	Hortenſis.
Leſſer Ditto,	-	Hippolais.
Hedge W.	-	Modularis.
White-throat,	-	Cinerea.
Leſſer White-throat,	-	Sylviella.
Graſshopper W.	-	Locuſtella.
Sedge W.	-	Salicaria.
Dartford W.	-	Dartfordienſis.
Redbreaſt,	-	Rubecula.
Wheat-ear,	-	Oenanthe.
Stone Chat,	-	Rubicola.
Whin Chat,	-	Rubetra.
Common Wren,	-	Troglodytes.
Gold-crested W.	-	Regulus.

Yellow

Yellow Wren,	-	Trochilus.
Wood W.	-	Sylvicola.
Reed W.	-	Arundinacea.
Redſtart,	-	Phœnicurus.
Blackcap,	-	Atricapilla.

GENUS XXV.

TITMOUSE,		PARUS.
Great Titmouſe,	-	Major.
Marſh T.	-	Paluſtris.
Colemouſe,	-	Ater.
Blue T.	-	Cæruleus.
Long-tailed T.	-	Caudatus.
Bearded T.	-	Biarmicus.
Creſted T.	-	Criſtatus.

GENUS XXVI.

SWALLOW.		HIRUNDO.
Chimney Swallow,	-	Ruſtica.
Martin,	-	Urbica.
Sand Martin,	-	Riparia.
Swift,	-	Apus.

GENUS XXVII.

GOATSUCKER.		CAPRIMULGUS.
European Goatſucker,	-	Europæus.

ORDER IV. Columbine.

GENUS XXVIII.

DOVE.		COLUMBA.
Ring Dove,	-	Palumbus.

Rock

Rock Dove,	-	Œnas.
Turtle D.	-	Turtur.

ORDER V. Gallinaceous.

GENUS XXIX.

PHEASANT.		PHASIANUS.
Common Pheaſant,	-	Colchicus.

GENUS XXX.

GROUS.		TETRAO.
Wood Grous,	-	Urogallus.
Black Gr.	-	Tetrix.
Red Gr.	-	Scoticus.
Ptarmigan,	-	Lagopus.

GENUS XXXI.

PARTRIDGE.		PERDIX. (Tetrao, *Lin.*)
Common Partridge,	-	Cinerea.
Guernſey P.	-	Rufa.
Quail,	-	Coturnix.

GENUS XXXII.

BUSTARD.		OTIS.
Great Buſtard,	-	Tarda.
Little B.	-	Tetrax.
Thick-kneed B.	-	Œdicnemus.

DIV. II.

WATER BIRDS.

ORDER VI. Grallæ.

GENUS XXXIII.

SPOONBILL.	PLATALEA.
White Spoonbill,	Leucorodia

GENUS XXXIV.

HERON.	ARDEA.
Common Heron,	Cinerea.
Crane,	Grus.
White Stork,	Ciconia.
Great White H.	Alba.
Night H.	Nycticorax.
African H.	Caspica.
Bittern,	Stellaris.
Little B.	Minuta.
Egret,	Garzetta.
Gardenian H.	Gardeni.

GENUS XXXV.

IBIS.	TANTALUS.
Gloſſy Ibis,	Igneus.

GENUS XXXVI.

CURLEW.	NUMENIUS. (Scolopax, *Lin.*)
Common Curlew,	Arquata.
Whimbrel,	

Whimbrel,	-	Phæopus.
Pygmy Curlew,	-	Pygmeus.

GENUS XXXVII.

SNIPE.		SCOLOPAX.
Woodcock,	-	Ruſticola.
Great Snipe,	-	Major.
Brown Sn.	-	Griſea.
Common Sn.	-	Gallinago.
Jack Sn.	-	Gallinula.
Jadreka Sn.	-	Limoſa.
Spotted Sn.	-	Totanus.
Common Godwit,	-	Œgocephala.
Red Godwit,	-	Lapponica.
Cinereous G.	-	Caneſcens.
Cambridge G.	-	Cantabrigienſis.
Greenſhank,	-	Glottis.
Redſhank,	-	Calidris.

GENUS XXXVIII.

SANDPIPER.		TRINGA.
Ruff,	-	Pugnax.
Lapwing,	-	Vanellus.
Gambet,	-	Gambetta.
Grey S.	-	Squatarola.
Aſh-coloured S.	-	Cinerea.
Green S.	-	Ocropus.
Common S.	-	Hypoleucos.
Spotted S.	-	Macularia.
Red S.	-	Iſlandica.

Little Sandpiper,	-	Puſilla.
Wood S.	-	Glareola.
Purre,	-	Cinclus.
Dunlin,	-	Alpina.
Knot,	-	Canutus.
Turnſtone,	-	Interpres.
Greenwich S.	-	Grenovicenſis.
Black S.	-	Lincolnienſis.
Brown S.	-	Fuſca.
Purple S.	-	Nigricans.
Yellow-legged S.	-	Flavipes.

GENUS XXXIX.

PLOVER.		CHARADRIUS.
Golden Plover,	-	Pluvialis.
Dotterel,	-	Morinellus.
Ringed P.	-	Hiaticula.
Sanderling,	-	Calidris.
Long-legged P.	-	Himantopus.
* Cream-coloured P.	-	Curſor.

GENUS XL.

OYSTER-CATCHER.		HÆMATOPUS.
Pied Oyſter-catcher,	-	Oſtralegus.

GENUS XLI.

RAIL.		RALLUS.
Water Rail,	-	Aquaticus.

* Dr. Latham has removed this bird in his *Index Ornithologicus* into a new genus called Curſorius, under the ſpecific title of Europæus.

GENUS

GENUS XLII.

GALLINULE.		GALLINULA. (Rallus & Fulica, *Lin*.)
Crake G.	-	Crex.
Common G.	-	Chloropus.
Spotted G.	-	Porzana.

ORDER VII. Pinnatipedes.

GENUS XLIII.

PHALAROPE.		PHALAROPUS.
Red Phalarope,	-	Hyperboreus.
Grey Ph.	-	Lobatus.

GENUS XLIV.

COOT.		FULICA.
Common Coot.	-	Atra.
Greater C.	-	Aterrima.

GENUS XLV.

GREBE.		PODICEPS. (Colymbus, *Lin*.)
Crested Grebe,	-	Cristatus.
Dusky Grebe,	-	Obscurus.
Eared Gr.	-	Auritus.
Red-necked Gr.	-	Rubricolis.
Little Gr.	-	Minor.
Black-chin Gr.	-	Hebridicus.
Sclavonian Gr.	-	Cornutus.

ORDER VIII. Palmipedes.

GENUS XLVI.

AVOSET.		RECURVIROSTRA.
Scooping Avoſet,	-	Avocetta.

GENUS XLVII.

AUK.		ALCA.
Great Auk,	-	Impennis.
Razor-billed A.	-	Torda.
Black-billed A.	-	Pica.
Puffin,	-	Arctica.
Little A.	-	Alle.

GENUS XLVIII.

GUILLEMOT.		URIA. (Colymbus, *Lin.*)
Fooliſh Guillemot,	-	Troile.
Leſſer G.	-	Minor.
Black G.	-	Grylle.

GENUS XLXIX.

DIVER.		COLYMBUS.
Northern Diver,	-	Glacialis.
Imber D.	-	Immer.
Speckled D.	-	Stellatus.
Black-throated D.	-	Arcticus.
Red-throated D.	-	Septentrionalis.

GENUS.

GENUS L.

TERN.		STERNA.
Sandwich Tern,	-	Boyſii.
Common T.	-	Hirundo.
Leſſer T.	-	Minuta.
Black T.	-	Fiſſipes.
Brown T.	-	Obſcura.

GENUS LI.

GULL.		LARUS.
Arctic Gull,	-	Paraſiticus.
Black-headed G.	-	Ridibundus.
Black-toed G.	-	Crepidatus.
Brown-headed G.	-	Erythropus.
Common G.	-	Canus.
Great Black-backed G.	-	Marinus.
Leſs Black-backed G.	-	Argentatus.
Herring G.	-	Fuſcus.
Laughing G.	-	Atricilla.
Red-legged G.	-	Cinerarius.
Skua G.	-	Catarractes.
Tarrock G.	-	Tridactylus.
Wagel G.	-	Nævius.
Winter G.	-	Hybernus.
Kittiwake.	-	Riſſa.

GENUS LII.

PETREL.		PROCELLARIA.
Fulmar Petrel,	-	Glacialis.
Shearwater P.	-	Puffinus.
Stormy P.	-	Pelagica.

GENUS LIII.

MERGANSER.		MERGUS.
Gooſander,	-	Merganſer.
Dun-diver,	-	Caſtor.
Red-breaſted M.	-	Serrator.
Smew,	-	Albellus.
Minute M.	-	Minutus.

GENUS LIV.

DUCK.		ANAS.
Whiſtling Swan,	-	Cygnus (ferus.)
Grey-lag Gooſe,	-	Anſer.
Red-breaſted G.	-	Ruficollis.
White-fronted G.	-	Albifrons.
Bean G.	-	Segetum.
Bernacle G.	-	Erythropus.
Brent G.	-	Bernicla.
Eider D.	-	Molliſſima.
Velvet D.	-	Fuſca.
Scoter,	-	Nigra.
Common Wild D.	-	Boſchas.
Shieldrake,	-	Tadorna.
Scaup D.	-	Marila.
Shoveler,	-	Clypeata.
Red-breaſted Sh.	-	Rubens.
Ferruginous D.	-	Ferruginea.
Bimaculated D.	-	Glocitans.
Gadwall,	-	Strepera.

Wigeon,

Wigeon,	-	Penelope.
Pochard,	-	Ferina.
Pintail D.	-	Acuta.
Long-tailed D.	-	Glacialis.
Golden-eye,	-	Clangula.
Morillon,	-	Glaucion.
Tufted D.	-	Fuligula.
Garganey,	-	Querquedula.
Teal,	-	Crecca.
Harlequin D.	-	Hiſtrionica.
King D.	-	Spectabilis.

GENUS LV.

PELICAN.		PELECANUS.
Corvorant,	-	Carbo.
Shag,	-	Graculus.
Gannet,	-	Baſſanus.

EXPLANATION

OF SOME

TECHNICAL TERMS

Uſed in ORNITHOLOGY by LINNÆUS and others, and in this WORK.

Cere. Cera. The naked ſkin that covers the baſe of the bill in the Hawk kind.

Lore. Lorum. The ſpace between the bill and eye, which in ſome birds, eſpecially of the Grebe kind, is naked.

Orbit. Orbita. The ſkin that ſurrounds the eye, which in ſome birds is bare of feathers, as in the Heron.

Iris, or Irides. That part of the eye ſurrounding the pupil, frequently of a different colour.

Nictitating membrane. A membrane capable of extending over the eye to preſerve it, and yet ſufficiently tranſparent

	tranſparent to admit the rays of light. Conſpicuous in the Owl genus.
Alula ſpuria. Baſtard wing.	A ſmall joint at the baſe of the greater quill-feathers furniſhed with three or four ſtiff feathers.
Vibriſſæ pectinatæ.	Stiff hairs that grow on each ſide the mouth in a regular form, as in the Goat-ſucker.
Primary quill-feathers, or primaries. Primores.	The largeſt feathers of the wing, ariſing from the firſt joint.
Secondary quill-feathers. Secondaries. Secondariæ.	The quill-feathers that ariſe from the ſecond joint.
Greater wing-coverts. Tectrices ſecundæ.	Thoſe feathers which immediately cover the baſe of the quills.
Leſſer wing-coverts. Tectrices primæ.	Thoſe ſmaller feathers which cover the wing in ſeveral rows above the greater coverts.
Under wing-coverts.	Thoſe feathers covering the wing underneath.

Upper

Upper tail-coverts. **Uropygium.**	Thoſe feathers immediately covering the baſe of the tail-feathers.
Under tail-coverts.	Thoſe feathers covering the baſe of the tail beneath.
Vent feathers.	Thoſe which lie between the vent and under tail-coverts.
Scapular feathers.	Thoſe which ariſe from the ſhoulders, and cover the ſides of the back.
Rectrices.	The tail.
Nucha.	The hind part of the head.
Roſtrum ſubulatum.	A ſtrait and ſlender bill.
Roſtrum cultratum.	A bill with ſharp edges.
Unguiculatum.	A bill with a nail at the end, as in the Ducks.
Lingua ciliata.	The tongue when edged with briſtles, as in the Ducks.
Integra.	When quite plain.
Lumbriciformis.	When the tongue is long, round, and ſlender, like a worm, as in Woodpeckers.
Nares Lineares.	When the noſtrils are very narrow.
Marginatæ.	When a rim round the noſtrils.

Fin-

Fin-footed. Pes lobatus. Pinnatus.	Such as the Grebes.
Scalloped.	When the membranes of the toes are indented, as in the Phalarope and Coot.
Pes ſcanſorius.	When formed for climbing, like the Woodpecker.
Pes tridactylus.	Such as want the back-toe.
Semi-palmated. Pes Semipalmatus.	When the webs reach only half way the toes.
Ungue poſtico ſeſſili.	When the hind claw adheres to the leg without any toe, as in the Petrel.
Digitis 4 omnibus palmatis.	When all four toes are connected together by a web, as in the Corvorant.
Pes compedes.	When the legs are placed ſo far behind as to be rendered almoſt uſeleſs in walking, as in the Grebes and Divers.

CATALOGUE of the PRINCIPAL AUTHORS referred to in this WORK.

Albin.	Natural Hiſtory of Birds, with coloured plates, engraven from life, by Eleazer Albin. 3 vols. 4to. London, 1738.
Arct. Zool.	Arctic Zoology. By Thomas Pennant, Eſq. 2 vol. 4to. 1784.
Borlaſe, Corn.	Natural Hiſtory of Cornwall. By William Borlaſe, A. M. fol. London, 1758.
Br. Zool.	Britiſh Zoology. By Thomas Pennant, Eſq. 4to. fourth edit. London, 1776.
Briſ.	Ornithologie. Par M. Briſſon. 4to. Paris, 1760.
Brown, Ill.	New Illuſtrations of Zoology. By Peter Brown. London, 1776.
Buf.	Hiſtoire Naturelle des Oiſeaux. Par M. Buffon. 4to. Paris 1770, 1783.
Don. Br. Birds.	Natural Hiſtory of Britiſh Birds. By E. Donovan, F. L. S. 8vo. London, 1799.
Edw.	Natural Hiſtory of Birds, and other rare and undeſcribed Animals. By George Edwards. 4to. 7 vols. London, 1743, 1763.
Fauna Suec.	Caroli Linnæi Fauna Suecica Siſtens Animalia Sueciæ Regni. 8vo. Stockholmiæ, 1761.

Gen.

Gen. of Birds. Genera of Birds. By Thomas Pennant, Eſq. 2 vol. 4to. 1784.

Gmel. Syſt. Syſtema Naturæ. By Jo. Fred. Gmelin. Leipſiæ, 1788.

Ind. Orn. Index Ornithologicus, ſive Syſtema Ornithologiæ, complectens, Avium Diviſionem in Claſſes, Ordines, Species, ipſarumque varietates, adjectis ſynonimis, locis, deſcriptionibus, &c. Studio et opera Joannis Latham, S. R. S.

Lath. Syn. A General Synopſis of Birds. By John Latham. 4to. 3 vol. London, 1785.

Lewin, Br. Birds. The Birds of Great Britain. By Wm. Lewin. 4to. London, 1789.

Lin. Mantis. Caroli à Linné, Eq. Aur. Mantiſſa Plantarum altera. 8vo. Holmiæ, 1771.

Lin. Syſt. Caroli à Linné Syſtema Naturæ, &c. 8vo. 3 vols. edit. 12. Holmiæ, 1766.

Lin. Tranſ. Tranſactions of the Linnæan Society. 4to. London.

Morton, Northamp. Natural Hiſtory of Northamptonſhire. By John Morton, A. M. folio. London, 1712.

Oſt. Menag. Portraits of rare and curious Birds, with their Deſcriptions, from the Menagery of Oſterley Park. By W. Hayes and Family. 4to. London, 1794. 1802.

Phil. Tranſ. Philoſophical Tranſactions of the Royal Society of London. 4to. London.

Plot. Oxf.	Natural Hiſtory of Oxfordſhire. Folio. Oxford, 1708.
Plot. Staf.	Natural Hiſtory of Staffordſhire. Folio. Oxford, 1686.
Pult. Cat. Dorſet.	Catalogues of the Birds, Shells, and ſome of the more rare Plants of Dorſetſhire, from the new and enlarged edition of Mr. Hutchins's Hiſtory of that County. By Richard Pulteney, M. D. F. R. S. Lond. and Edinb. and F. L. S. London, 1799.
Raii Syn.	Joannis Raii Synopſis Methodica Avium et Piſcium, &c. 8vo. Londini, 1713.
Sib. Scot.	Scotia Illuſtrata, ſive Prodromus Hiſtoriæ Naturalis, &c. Authore Roberto Sibbaldo, M. D. Eq. Aur. Folio. Edinburgi, 1684.
Walc. Syn.	Synopſis of Britiſh Birds. By John Walcot, Eſq. 4to. London, 1789.
White, Hiſt. Selb.	Natural Hiſtory and Antiquities of Selborne in the County of Southampton. By Gilb. White. 4to. London, 1789.
Will. Orn.	The Ornithology of Francis Willughby, Eſq. F. R. S. Folio. London, 1678.

THE END.

Printed by T. Bensley, Bolt Court, Fleet Street, London.



